

# Reproductive Physiology

## Lecture 1

# Hypothalamic-Pituitary-Gonadal axis

**DR. MOHAMMED ALOTAIBI**

ASSISTANT PROFESSOR OF PHYSIOLOGY

COLLEGE OF MEDICINE

DEPARTMENT OF PHYSIOLOGY

EMAIL: [MFALOTAIBI@KSU.EDU.SA](mailto:MFALOTAIBI@KSU.EDU.SA)

TEL: 0114672629

# OBJECTIVES



## **By the end of this lecture, you should be able to:**

1. Define hormones
2. Characterize hypothalamic-pituitary relationship
3. Name the hypophysiotropic hormones and outline the effects that each has on anterior pituitary function
4. Name anterior pituitary gonadotropic hormones and outline the effects that each has on the gonads
5. Describe the negative and positive feedback mechanisms in the hypothalamic-pituitary-gonadal axis and their importance in the control of reproductive function


# Hypothalamic-Pituitary-Gonadal axis



**Hormone:** Chemical substance (messengers) produced by ductless glands that are transported in the circulation to the target cells where they regulate the metabolic processes.

**The anterior pituitary secretes six hormones:**

1. Adrenocorticotrophic hormone (ACTH)
2. Thyroid-stimulating hormone (TSH)
3. Growth hormone (GH)
4. Prolactin (PRL)
5. **Follicle-stimulating hormone (FSH)**
6. **Luteinizing hormone (LH)**



## Control of *male sexual functions* by hormones from the hypothalamus and anterior pituitary gland:



**GnRH** is a peptide secreted by the arcuate nuclei of the hypothalamus through the hypothalamic-hypophyseal portal system to the anterior pituitary gland and stimulates the release of **gonadotropins** (LH and FSH).

**GnRH** is secreted intermittently for few minutes every 1 to 3 hrs.

- The secretion of LH by the anterior pituitary is also cyclical following the **pulsatile** release of GnRH.


# Testosterone



Testosterone is secreted by **leydig cells** in the interstitium of the testis, by **LH** stimulation from the AP and its release is directly proportional to the amount of LH. Mature leydig cells are found in a child's testis few weeks after birth & then becomes quiescent until puberty when they appear again.

## **Inhibition of anterior pituitary secretion of LH and FSH by testosterone – negative feedback control of testosterone secretion:**

Most of the inhibitory effects result from a direct effect of testosterone in the hypothalamus to decrease secretion of GnRH which causes a decreased secretion of both LH & FSH.



# Regulation of spermatogenesis by FSH and testosterone



- **FSH** binds with specific FSH receptors attached to the ***Sertoli cell*** in the seminiferous tubules, which causes these cells to grow & secrete spermatogenic substances. Also testosterone & dihydrotestosterone diffuses into the seminiferous tubules from the Leydig cells and affect the spermatogenesis. Both FSH & testosterone are necessary to initiate spermatogenesis.

## **Negative feedback control of seminiferous tubule activity – role of the hormone inhibin:**

- When the seminiferous tubules fail to produce sperms, secretion of FSH from the AP increases. Conversely, when spermatogenesis proceeds rapidly, pituitary secretion of FSH diminishes. This is due to the secretion of inhibin hormone from the Sertoli cells which strongly inhibits the AP- FSH and has a slight inhibitory effect on the hypothalamus to inhibit GnRH secretion.



# Regulation of the *female monthly rhythm*, interplay between the ovarian and hypothalamic-pituitary hormones



- Secretion of AP hormone is controlled by “releasing hormones” formed in the hypothalamus and transported to the AP gland by the hypothalamic-hypophyseal portal system.
- **Intermittent, pulsatile** secretion of GnRH by the hypothalamus stimulates pulsatile release of LH from the AP:

GnRH is secreted in pulses lasting 5 to 25 minutes every 1 to 2 hrs. The pulsatile release of GnRH causes intermittent output of LH secretion about every 90 minutes.

## **Hypothalamic centers for release of GnRH:**

The neural activity that causes pulsatile release of GnRH occurs in the mediobasal hypothalamus, in the arcuate nuclei which regulates most of the female sexual activity.

# Negative feedback effects of estrogen and progesterone in decreasing both LH and FSH secretion



- Estrogen in small amounts has strong effect to inhibit the production of LH & FSH. This inhibitory effect of estrogen is increased when progesterone is available. These feedback effects seems to operate mainly on the AP directly & to lesser extent on the hypothalamus to inhibit the secretion of GnRH.


## **Hormone inhibin from the corpus luteum inhibits FSH & LH secretion:**

- The hormone inhibin secreted by the granulosa cells of the **ovarian corpus luteum** (in the same way that **sertoli cells** in males) inhibits the secretion of FSH & to lesser extent LH.

# Positive feedback effect of estrogen before ovulation (The pre-ovulatory LH surge)



- AP secretes increased amount of LH for 1 to 2 days before ovulation. FSH surge is much smaller in the pre-ovulatory than LH surge.



# Positive feedback effect of estrogen before ovulation (The pre-ovulatory LH surge)



## The possible causes of LH secretion could be:

- estrogen has special **positive feedback** effect of stimulating pituitary secretion of LH & to a lesser extent FSH
- the granulosa cells of the follicle begin to secrete small increasing amount of progesterone about 1 day before ovulation which stimulate LH secretion


# Feedback oscillation of the hypothalamic-pituitary-ovarian system



## Postovulatory secretion of the ovarian hormones, and depression of the pituitary gonadotropins

During the postovulatory phase (between ovulation & beginning of menstruation) the corpus luteum secretes large quantities of progesterone & estrogen & inhibin which all together cause negative feedback effect on AP & hypothalamus to inhibit both FSH & LH secretion.

(lowest level 3-4 days before the onset of menstruation)



# Feedback oscillation of the hypothalamic-pituitary-ovarian system



## Follicular growth phase

- 2 to 3 days before menstruation, corpus luteum regress & secretion of estrogen, progesterone & inhibin decreases.
- This decrease removes the negative feedback effect on AP hormones. Therefore a day after menstruation FSH secretion begins to increase (2 folds) while LH secretion is slightly increases.
- These hormones causes growth of the follicle. During the first 11 to 12 days of this follicle growth, the rate of secretion of FSH & LH decreases slightly because of the negative feedback effect of estrogen on the AP.

# Feedback oscillation of the hypothalamic-pituitary-ovarian system



## Pre ovulatory surge of LH & FSH causes ovulation

- About 12 days of the monthly cycle, the high secretion of **FSH & LH** start to increase due to high level of **estrogens**, that is causing positive feedback on the anterior pituitary and leading to pre-ovulatory LH surge & FSH surge.



***The End***

***Thank You***