

Heart Failure

Khalid Alhabib
Professor of Cardiology





Figure 1: Unadjusted mortality at 1 year, by region and cause


Definition

- ▶ Heart failure is a complex clinical syndrome


Can result from:

- ▶ structural or functional cardiac disorder
- ▶ impairs the ability of the ventricle to **fill** with or **eject** blood


- ▶ Characterized by:
 - ▶ signs and symptoms of intravascular and interstitial volume overload and/or
 - ▶ manifestations of inadequate tissue perfusion
- 

- ▶ Heart failure may result from an acute insult to cardiac function, such as a large myocardial infarction, valvular diseases, myocarditis, and cardiogenic shock
 - ▶ More commonly, from a chronic process
- 

Common Causes

- ▶ Coronary artery disease
 - ▶ Hypertension
 - ▶ Valvular heart disease
 - ▶ Dilated cardiomyopathy
- 

Nomenclature


- ▶ Heart failure vs.
 - ▶ Cardiomyopathy
 - ▶ LV dysfunction
 - ▶ Pulmonary edema
- 


Classification

- ▶ Left vs. Right
- ▶ Systolic vs. Diastolic
- ▶ High output vs. low output

HFrEF vs HFpEF

Heart Failure Syndrome

- ▶ The initial manifestations of hemodynamic dysfunction are a reduction in stroke volume and a rise in ventricular filling pressures under conditions of increased systemic demand for blood flow
 - ▶ This stimulates a variety of interdependent compensatory responses involving the cardiovascular system, neurohormonal systems, and alterations in renal physiology
- 



Modified Framingham clinical criteria for the diagnosis of heart failure

Major
Paroxysmal nocturnal dyspnea
Orthopnea
Elevated jugular venous pressure
Pulmonary rales
Third heart sound
Cardiomegaly on chest x-ray
Pulmonary edema on chest x-ray
Weight loss ≥ 4.5 kg in five days in response to treatment of presumed heart failure
Minor
Bilateral leg edema
Nocturnal cough
Dyspnea on ordinary exertion
Hepatomegaly
Pleural effusion
Tachycardia (heart rate ≥ 120 beats/min)
Weight loss ≥ 4.5 kg in five days
Diagnosis
The diagnosis of heart failure requires that 2 major or 1 major and 2 minor criteria cannot be attributed to another medical condition.

From Senni, M, Tribouilloy, CM, Rodeheffer, RJ, et al, *Circulation* 1998; 98:2282; adapted from McKee, PA, Castelli, WP, McNamara, PM, Kannel, WB. *N Engl J Med* 1971; 85:1441.

- ▶ **FACTORS THAT MAY PRECIPITATE ACUTE DECOMPENSATION OF CHRONIC HEART FAILURE**

Discontinuation of therapy (patient noncompliance or physician initiated)

Initiation of medications that worsen heart failure (calcium antagonists, β -blockers, nonsteroidal anti-inflammatory drugs, antiarrhythmic agents)

Iatrogenic volume overload (transfusion, fluid administration)

Dietary indiscretion

Pregnancy

Exposure to high altitude

Arrhythmias

Myocardial ischemia or infarction

Worsening hypertension

Worsening mitral or tricuspid regurgitation

Fever or infection

Anemia

Events usually leading to rapid deterioration

- Rapid arrhythmia or severe bradycardia/conduction disturbance
- Acute coronary syndrome
- Mechanical complication of acute coronary syndrome (e.g. rupture of interventricular septum, mitral valve chordal rupture, right ventricular infarction)
- Acute pulmonary embolism
- Hypertensive crisis
- Cardiac tamponade
- Aortic dissection
- Surgery and perioperative problems
- Peripartum cardiomyopathy

Events usually leading to less rapid deterioration

- Infection (including infective endocarditis)
- Exacerbation of COPD/asthma
- Anaemia
- Kidney dysfunction
- Non-adherence to diet/drug therapy
- Iatrogenic causes (e.g. prescription of an NSAID or corticosteroid; drug interactions)
- Arrhythmias, bradycardia, and conduction disturbances not leading to sudden, severe change in heart rate
- Uncontrolled hypertension
- Hypothyroidism or hyperthyroidism
- Alcohol and drug abuse

Evaluation

Evidence for Congestion (Elevated Filling Pressure)

- Orthopnea
- High Jugular Venous Pressure
- Increasing S_3
- Loud P_2
- Edema
- Ascites
- Rales (Uncommon)
- Abdominojugular Reflux
- Valsalva Square Wave

Evidence for Low Perfusion

- Narrow Pulse Pressure
- Pulsus Alterations
- Cool Forearms and Legs
- May Be Sleepy, Obtunded
- ACE Inhibitor-Related
 - Symptomatic Hypotension
- Declining Serum Sodium Level
- Worsening Renal Function

Low Perfusion at Rest?

Congestion at Rest?

	No	Yes
No	Warm and Dry A	Warm and Wet B
Yes	Cold and Dry L	Cold and Wet C


NYHA Classification

Class I	No limitations of activities; no symptoms with ordinary activities
Class II	Slight or mild limitation of activity; comfortable with rest or mild exertion
Class III	Marked limitation of activity; comfortable only at rest
Class IV	Any physical activity brings on discomfort, and symptoms occur at rest

ACC/AHA

At Risk for Heart Failure

Heart Failure



STAGE A

At high risk for HF but without structural heart disease or symptoms of HF.

e.g., Patients with:
-hypertension
-atherosclerotic disease
-diabetes
-obesity
-metabolic syndrome
or
Patients
-using cardiotoxins
-with FHx CM

THERAPY

GOALS

- Treat hypertension
- Encourage smoking cessation
- Treat lipid disorders
- Encourage regular exercise
- Discourage alcohol intake, illicit drug use
- Control metabolic syndrome

DRUGS

- ACEI or ARB in appropriate patients (see text) for vascular disease or diabetes

STAGE B

Structural heart disease but without signs or symptoms of HF.

e.g., Patients with:
-previous MI
-LV remodeling including LVH and low EF
-asymptomatic valvular disease

THERAPY

GOALS

- All measures under Stage A

DRUGS

- ACEI or ARB in appropriate patients (see text)
- Beta-blockers in appropriate patients (see text)

DEVICES IN SELECTED PATIENTS

- Implantable defibrillators

STAGE C

Structural heart disease with prior or current symptoms of HF.

e.g., Patients with:
-known structural heart disease and
-shortness of breath and fatigue, reduced exercise tolerance

THERAPY

GOALS

- All measures under Stages A and B
- Dietary salt restriction

DRUGS FOR ROUTINE USE

- Diuretics for fluid retention
- ACEI
- Beta-blockers

DRUGS IN SELECTED PATIENTS

- Aldosterone antagonist
- ARBs
- Digitalis
- Hydralazine/nitrates

DEVICES IN SELECTED PATIENTS

- Biventricular pacing
- Implantable defibrillators

STAGE D

Refractory HF requiring specialized interventions.

e.g., Patients who have marked symptoms at rest despite maximal medical therapy (e.g., those who are recurrently hospitalized or cannot be safely discharged from the hospital without specialized interventions)

THERAPY

GOALS

- Appropriate measures under Stages A, B, C
- Decision re: appropriate level of care

OPTIONS

- Compassionate end-of-life care/hospice
- Extraordinary measures
 - heart transplant
 - chronic inotropes
 - permanent mechanical support
 - experimental surgery or drugs

Investigations to consider in all patients

Transthoracic echocardiography is recommended to evaluate cardiac structure and function, including diastolic function (Section 4.1.2), and to measure LVEF to make the diagnosis of HF, assist in planning and monitoring of treatment, and to obtain prognostic information.

A 12-lead ECG is recommended to determine heart rhythm, heart rate, QRS morphology, and QRS duration, and to detect other relevant abnormalities (*Table 5*). This information also assists in planning treatment and is of prognostic importance. A completely normal ECG makes systolic HF unlikely.

Measurement of blood chemistry (including sodium, potassium, calcium, urea/blood urea nitrogen, creatinine/estimated glomerular filtration rate, liver enzymes and bilirubin, ferritin/TIBC) and thyroid function is recommended to:

- (i) Evaluate patient suitability for diuretic, renin–angiotensin–aldosterone antagonist, and anticoagulant therapy (and monitor treatment)
- (ii) Detect reversible/treatable causes of HF (e.g. hypocalcaemia, thyroid dysfunction) and co-morbidities (e.g. iron deficiency)
- (iii) Obtain prognostic information.

A complete blood count is recommended to:

- (i) Detect anaemia, which may be an alternative cause of the patient's symptoms and signs and may cause worsening of HF
- (ii) Obtain prognostic information.

Measurement of natriuretic peptide (BNP, NT-proBNP, or MR-proANP) should be considered to:

- (i) Exclude alternative causes of dyspnoea (if the level is below the exclusion cut-point—see *Figure 1*—HF is very unlikely)
- (ii) Obtain prognostic information.

A chest radiograph (X-ray) should be considered to detect/exclude certain types of lung disease, e.g. cancer (does not exclude asthma/COPD). It may also identify pulmonary congestion/oedema and is more useful in patients with suspected HF in the acute setting.

Investigations to consider in selected patients

CMR imaging is recommended to evaluate cardiac structure and function, to measure LVEF, and to characterize cardiac tissue, especially in subjects with inadequate echocardiographic images or where the echocardiographic findings are inconclusive or incomplete (but taking account of cautions/contraindications to CMR).

Coronary angiography is recommended in patients with angina pectoris, who are considered suitable for coronary revascularization, to evaluate the coronary anatomy.


Myocardial perfusion/ischaemia imaging (echocardiography, CMR, SPECT, or PET) should be considered in patients thought to have CAD, and who are considered suitable for coronary revascularization, to determine whether there is reversible myocardial ischaemia and viable myocardium.

Left and right heart catheterization is recommended in patients being evaluated for heart transplantation or mechanical circulatory support, to evaluate right and left heart function and pulmonary arterial resistance.

Exercise testing should be considered:


- (i) To detect reversible myocardial ischaemia
- (ii) As part of the evaluation of patients for heart transplantation and mechanical circulatory support
- (iii) To aid in the prescription of exercise training
- (iv) To obtain prognostic information.


Therapy



- ▶ 56 Y/O gentleman
- ▶ Diagnosed dilated cardiomyopathy
- ▶ LVEF 25%
- ▶ NYHA class II
- ▶ O/E B/P 112/68 HR 82 bpm
- ▶ JVP 7 cm water,
- ▶ Soft S3 and grade 2 PSM
- ▶ Chest clear,
- ▶ No LL edema and warm extremities

EMEDU






06/01/05 14:28:29

P80 4MHz S611

B5




CN15
16cm
DB72
V 84




MI < 0.4

06/01/05 14:25:43

P80 4MHz S611



CN0
16cm
DB48
V 62



TIS<0.4

	Starting dose (mg)	Target dose (mg)
ACE inhibitor		
Captopril ^a	6.25 t.i.d.	50 t.i.d.
Enalapril	2.5 b.i.d.	10–20 b.i.d.
Lisinopril ^b	2.5–5.0 o.d.	20–35 o.d.
Ramipril	2.5 o.d.	5 b.i.d.
Trandolapril ^b	0.5 o.d.	4 o.d.
Beta-blocker		
Bisoprolol	1.25 o.d.	10 o.d.
Carvedilol	3.125 b.i.d.	25–50 b.i.d.
Metoprolol succinate (CR/XL)	12.5/25 o.d.	200 o.d.
Nebivolol ^c	1.25 o.d.	10 o.d.
ARB		
Candesartan	4 or 8 o.d.	32 o.d.
Valsartan	40 b.i.d.	160 b.i.d.
Losartan ^{b,c}	50 o.d.	150 o.d.
MRA		
Eplerenone	25 o.d.	50 o.d.
Spirolactone	25 o.d.	25–50 o.d.

Treatments (or combinations of treatments) that may cause harm in patients with symptomatic (NYHA class II–IV) systolic heart failure

Recommendations

Thiazolidinediones (glitazones) should not be used as they cause worsening HF and increase the risk of HF hospitalization.

Most CCBs (with the exception of amlodipine and felodipine) should not be used as they have a negative inotropic effect and can cause worsening HF.

NSAIDs and COX-2 inhibitors should be avoided if possible as they may cause sodium and water retention, worsening renal function and worsening HF.

The addition of an ARB (or renin inhibitor) to the combination of an ACE inhibitor AND a mineralocorticoid antagonist is NOT recommended because of the risk of renal dysfunction and hyperkalaemia.

Symptomatic Heart Failure + Reduced Ejection Fraction

Detect Co-morbidities and Precipitating Factors

Non-cardiovascular

Anaemia
Pulmonary disease
Renal dysfunction
Thyroid dysfunction
Diabetes

Cardiovascular

Ischaemia/CAD
Hypertension
Valvular dysfunction
Diastolic dysfunction
Atrial fibrillation
Ventricular dysrhythmias
Bradycardia

Diuretic + ACEI (or ARB)
Titrate to clinical stability

β -Blocker

Persisting signs and symptoms?

Yes

No

ADD aldosterone antagonist OR ARB

Persisting symptoms?

Yes

No

QRS >120 ms?

LVEF <35%?

Yes

No

Yes


No

**Consider:
CRT-P or CRT-D**

**Consider: digoxin,
hydralazine/nitrate, LVAD,
transplantation**

Consider ICD

**No further treatment
indicated**



Risk factor modification	Understand the importance of smoking cessation Monitor blood pressure if hypertensive Maintain good glucose control if diabetic Avoid obesity
Diet recommendation	Sodium restriction if prescribed Avoid excessive fluid intake Modest intake of alcohol Monitor and prevent malnutrition
Exercise recommendations	Be reassured and comfortable about physical activity Understand the benefits of exercise Perform exercise training regularly
Sexual activity	Be reassured about engaging in sex and discuss problems with healthcare professionals Understand specific sexual problems and various coping strategies
Immunization	Receive immunization against infections such as influenza and pneumococcal disease
Sleep and breathing disorders	Recognize preventive behaviour such as reducing weight of obese, smoking cessation, and abstinence from alcohol Learn about treatment options if appropriate
Adherence	Understand the importance of following treatment recommendations and maintaining motivation to follow treatment plan

Acute Heart Failure




Two Minute Assessment of Hemodynamic Profile

		Congestion at rest?	
		NO	YES
Low perfusion at rest?	NO	Warm & Dry A	Warm & Wet B
	YES	Cold & Dry L	Cold & Wet C


<u>Evidence for low perfusion</u>	<u>Evidence for Congestion</u>
Narrow pulse pressure ⁺	Orthopnea ⁺
Cool extremities ⁺	Elevated JVP [±]
May be sleepy, obtunded	Edema (25%)
Suspect from ACEI hypotension and low Serum Sodium	Pulsatile hepatomegaly
One cause of worsening renal fn	Ascites
	Rales (rare in chronic HF)
	Louder S3
	P2 radiation leftward
	Abdomino-jugular reflex
	Valsalva square wave

** Most helpful*





For Heart Failure Diagnosis



Optimal NT-proBNP Cut-points

“Rule in”

Age strata	Optimal cut-point	Sensitivity	Specificity	PPV	NPV	Accuracy
All <50 years (n=183)	450 pg/ml	97%	93%	76%	99%	95%
All 50-75 years (n=554)	900 pg/ml	90%	82%	82%	88%	85%
All >75 years (n=519)	1800 pg/ml	85%	73%	92%	55%	83%
Overall average		92%	84%	88%	66%	93%

“Rule out”

	Optimal cut-point	Sensitivity	Specificity	PPV	NPV	Accuracy
Rule out	300 pg/ml	99%	62%	55%	99%	83%

Suspected acute heart failure

History/examination
(including blood pressure and respiratory rate)

Chest X-ray

Echocardiogram or NP (or both)

Blood chemistry

ECG

Oxygen saturation

Full blood count

**Simultaneously
assess for**

Ventilation/
systemic
oxygenation
inadequate?^a

Life-threatening
arrhythmia/
bradycardia?^b

Blood pressure
<85 mmHg
or shock^c

Acute
coronary
syndrome^d

Acute
mechanical
cause/severe
valvular disease^e

**Urgent
action
if present**

- Oxygen
- NIV
- ETT and
invasive
ventilation

- Electrical
cardioversion
- Pacing

- Inotrope/
vasopressor
- Mechanical
circulatory
support
(e.g. IABP)

- Coronary
reperfusion
- Antithrombotic
therapy

- Echocardiography
- Surgical/
percutaneous
intervention

ECG = electrocardiogram; ETT = endotracheal tube; IABP = intra-aortic balloon pump; NIV = non-invasive ventilation; NP = natriuretic peptide.

^aFor example, respiratory distress, cyanosis, SpO₂ <90% or PaO₂ <60 mmHg (8.0 kPa).