


Pulmonary Embolism

No.11

[Editing file](#)



Objectives :

- ★ Prevalence of PE
- ★ Risk factors
- ★ Clinical features
- ★ Pathophysiology
- ★ Massive PE
- ★ Diagnostic workup
- ★ Treatment

Color index

Original text
Females slides
Males slides
Doctor's notes ⁴³⁸
Doctor's notes ⁴³⁹
Doctor's notes ⁴⁴²
New text in slides ⁴⁴²
Text book
Important
Golden notes
Extra


Pulmonary Embolism

الانصمام الرئوي

Definition

- The majority of pulmonary emboli arise from the propagation of lower limb deep vein thrombosis.
- Rare causes include:
 - Septic emboli (from endocarditis affecting the tricuspid or pulmonary valves)
 - Tumour (especially choriocarcinoma)
 - Fat following fracture of long bones such as the femur
 - Air and amniotic fluid, which may enter the mother's circulation following delivery

Saddle shaped thrombi
U shaped



PE is **FATAL**: PE caused by **Fat, Air, Thrombus, Amniotic fluid**, and Less common, i.e., bacterial, tumor, and cement.

Epidemiology

50,000 individuals die from PE each year in USA

The incidence of PE in USA is 500,000 per year

The annual incidence of pulmonary embolism in the population is 1 per 1000 people, but this increases sharply with age. 1.4 per 1000 people aged 40-49
11.3 per 1000 aged 80 years or over

Total Incidence 630,000

89%
Survival >1hr
563,000

11%
Death within 1 hr
67,000¹

71%
Dx not made²
400,000

29%
Dx made, therapy instituted
163,000

70%
Survival 280,000

30%³
Death 120,000

92%
Survival 150,000

8%
Death 120,000

Massive pulmonary embolism⁴

- It is a catastrophic entity which often results in acute right ventricular failure and death.
- Frequently undiscovered until autopsy.
- Fatal PE typically leads to death within one to two hours of the event.

1- Death within 1h is caused by Massive PE.

2- Because PE symptoms aren't specific.


3- If you send them home without treatment there's a high risk they will get another fatal PE.

4- Massive means major hemodynamic effect (shock & hypotension), not reserved to the size.

Risk Factors

◀ Risk factors for venous thrombosis (Virchow Triad)

1. **Stasis:** eg. HF or any obstruction such as tumors, paraplegia.
2. **Alterations in the coagulation-fibrinolytic system:** (increase the chances of clot formation). eg. Deficiency in protein S or C or the presence of factor leiden 5 (mutated form of Factor 5), COVID-19 increase risk of thrombosis.
3. **Injury to venous intima:** eg. Autoimmune diseases causing Vasculitis, Smokers



- **50%** of venous thromboembolism events are associated with a **transient risk factor**, **20%** are associated with cancer.
- The remainder **30%** are associated with minor or no risk factors and are thus **classified as unprovoked**.

◀ Source of emboli

1

Deep venous thrombosis (>95%)¹

2

Other veins: Renal, Uterine, Right cardiac chambers.

◀ Risk factors of DVT : ★

- General anesthesia usually they are at risk after 1-4 weeks, If one year passed = no risk.
- Lower limb or pelvic injury or surgery²
- Congestive heart failure
- Prolonged immobility³
- Pregnancy
- Oral contraceptives pills
- Malignancy
- Obesity
- Advanced age
- Coagulation problems
- Postpartum

just read it


Strong risk factor (odds ratio > 10)	Moderate risk factor (odds ratio 2-9)	weak risk factor (odds ratio <2)
<ul style="list-style-type: none"> • Hip or leg fracture • Hip or leg joint replacement • Major general surgery • Major trauma • Spinal cord injury⁴ <p>Because of the prophylaxis (heparin), we don't see pulmonary embolism cases after these surgeries.</p>	<ul style="list-style-type: none"> • Arthroscopic knee surgery • Central venous lines (femoral central lines) • Congestive heart or respiratory failure⁵ • Hormone replacement therapy • Oral contraceptive therapy • Malignancy • Paralytic stroke • Postpartum • Previous VTE • Thrombophilia (high platelet count) 	<ul style="list-style-type: none"> • Bed rest >3 days • Immobility due to sitting • Increasing age • Laparoscopic surgery • Obesity • Pregnancy • Varicose veins

- Above the knee (between pelvis & knee): increased risk.
- Under the knee: decreased risk.

- 1- When you get a pt with PE it's better to look for signs of DVT. Most of them in the lower limbs (**thigh**), but under the knee the risk is low.
- 2- Thrombi of lower legs usually don't detach and travel to the rest of the body, mostly hips and pelvic thrombi are the ones that do that.
- 3- As in stroke patient or long distance travel especially for elderly.
- 4- Muscles become flaccid with weak contractions → unable to pump up blood. Ex: paraplegia
- 5- Respiratory failure due to polycythemia (develop due to hypoxia).


Pathophysiology

- Emboli block a portion of pulmonary vasculature, leading to increased pulmonary vascular resistance, pulmonary artery pressure, and right ventricular pressure. If it is severe (large blockage), acute cor pulmonale may result.
- Blood flow decreases in some areas of the lung. Dead space is created in areas of the lung in which there is ventilation but no perfusion. The resulting hypoxemia and hypercarbia drive respiratory effort, which leads to tachypnea.
- If the size of the dead space is large (large PE), clinical signs are more overt (SOB, tachypnea).
 - If there is a clot in the pulmonary artery, the right ventricle cannot pump blood due to the resistance. This resistance will result in an increase in diameter (D shape) and a bulge of the septum to the left side. This reduces the left ventricular EDV, which in turn reduces stroke volume, resulting in a decrease in cardiac output (Frank-Starling mechanism). Because the heart is a closed chamber, the left and right sides of the heart pump the same amount of blood. The right ventricle has a pressure of 20/50 mmHg and the left ventricle has a pressure of 120/80 mmHg. Because of the resistance, they pump the same amount of blood despite the pressure difference. Because the resistance of the lungs is very low (negligible), it does not require strong muscles to pump blood to the pulmonary circulation, as opposed to the systemic circulation, which has high resistance. This is why we refer to the right ventricle as the (volume chamber) and the left ventricle as the (pressure chamber). In contrast to the left ventricle, the right ventricle has thin walls. A sudden increase in resistance causes the left ventricle to dilate (loses its ability to contract properly).
 - The effects of an embolus depend on the extent to which it obstructs the pulmonary circulation, the duration over which that obstruction accumulates, and the pre-existing state of the patient, which has been defined only imprecisely. In general, a patient who has pre-existing cardiopulmonary disease or who is old, frail or debilitated will be more sensitive to the effects of pulmonary embolism than a patient who was well until the embolic event occurred. Most emboli are multiple. As both the extent and chronicity of obstruction vary so widely, pulmonary embolism can produce widely differing clinical pictures. Disregarding chronic thromboembolic pulmonary hypertension, it is convenient to classify pulmonary embolism into three main types (table 3). The first and most common presentation is dyspnoea with or without pleuritic pain and haemoptysis (acute minor pulmonary embolism). The second presentation is haemodynamic instability, which is associated with acute massive pulmonary embolism. The third and least common presentation mimics heart failure or indolent pneumonia, especially in the elderly (subacute massive pulmonary embolism).



- Massive PE causes an increase in Pulmonary Vascular Resistance [PVR], PAP, and right ventricular pressure → right ventricular outflow obstruction → decrease left ventricular preload → Decrease CO.
 - If it is severe (large blockage), acute cor pulmonale may result
 - Acute cor pulmonale:** This sudden right heart failure may be caused by a saddle embolus obstructing the pulmonary artery or sudden overload of a chronic cor pulmonale by pneumonia.
 - Chronic cor pulmonale:** This form of chronic right heart failure is a consequence of chronic pulmonary hypertension.

◀ Pulmonary Embolism Death Spiral: ★




- In patients without cardiopulmonary disease, occlusion of 25-30% of the vascular bed → increase in PAP.
- Hypoxemia ensues → stimulating vasoconstriction → increase in PAP.
- More than 50% of the vascular bed has to be occluded before PAP becomes substantially elevated.
- When obstruction approaches 75%, the RV must generate systolic pressure in excess of 50 mmHg to preserve pulmonary circulation.
- The normal RV is unable to accomplish this acutely and eventually fails.

Sign & Symptoms



Sudden onset dyspnea



Pleuritic chest pain¹
From lung infarction



Haemoptysis²

- Clinical clues cannot make the diagnosis of PE, their main value lies in suggesting the diagnosis.
- Note that only one-third of patients with PE will have signs and symptoms of a DVT.
- Syncope seen in large PE.

◀ Signs or symptoms observed in patients with thromboembolism (a study) Important to know these two things DVT and PE

Disease	Sign & Symptoms	Stein et al., % (n=117)	Anderson et al., % (n=131)
Deep vein thrombosis	Swelling ⁴	28	88
	Pain (unilateral)	26	56
	Tenderness	-	55
	Warmth	-	42
	Redness	-	34
	Homan's sign	4	13
	Palpable cord	-	6
PE	Dyspnea	73	77
	Tachypnea	70	70
	Chest pain	66	55
	Cough	37	-
	Tachycardia	30	43
	Cyanosis	1	18
	Hemoptysis ³	13	13
	Wheezing	9	-
	Hypotension	-	10
	Syncope	-	10
	↑Jugular venous pulse	-	8
	Temp > 38.5°C	7	-
	S3 gallop	3	5
Pleural friction rub	3	2	

1- Pleuritic chest pain is characterized by being: localized (the patient locate the pain with his finger), stabbing pain, occurs during inspiration.

2- Not common nor specific.

3- Hemoptysis occurs with wedge infarction, and with PE usually there is no infarction because of the blood dual supply (pulmonary artery and bronchial).

4- Not all patients will experience swelling. For example, if you suspect a patient has PE and examine his legs and find no swelling or tenderness, you do not rule out DVT. If you perform a Duplex, you may or may not find a DVT. If the duplex is normal, DVT cannot be ruled out because the clot may have traveled.

PE Probability Scores

I don't like to ask about these scores in the exam but they may ask you about them in OSCE

The diagnostic process starts with assessment of the clinical probability using a clinical prediction score that takes account of the individual patient's clinical features, the presence or absence of risk factors for VTE, and whether an alternative diagnosis is likely to explain the symptoms and signs. There are a number of clinical prediction scores available, and NICE recommends the modified two-level Wells score for DVT (Box 29.3) and the two-level Wells score for PE (Box 29.4); these categorize patients into groups that are likely or unlikely to have DVT or PE, respectively. Clinical prediction scores standardize clinical assessment and increase reproducibility among less experienced clinicians. However, in themselves, they do not confirm or exclude the diagnosis of VTE and further assessment is necessary.

1

Wells Rule MORE COMMONLY USED

- Using **traditional** score:
 - <2.0 points indicates low probability
 - 2.0- 6.0 points indicates moderate probability
 - >6.0 points indicates high probability
- Using **simplified** score
 - >4 points indicates that PE is likely
 - ≤4 points indicates that PE is unlikely.

Wells Rule	Points
Sign or symptom of DVT (minimum of leg swelling and pain with palpation)	3
Alternative diagnosis is less likely than PE	3
HR > 100 beats/min	1.5
Immobilization/surgery in previous 4 wk	1.5
History of DVT or PE	1.5
Hemoptysis	1
Active cancer (on treatment, treated in the last 6 months, or palliative)	1

2

Modified and simplified Geneva rules

Modified Geneva Rule	Points
Age ≥ 65 years	1
Previous DVT or PE	3
Surgery or fracture within 1 month	2
Active cancer	2
Unilateral lower limb pain	3
Pain on deep palpation of lower limb and unilateral edema	4
Hemoptysis	2
HR 75-94 beats/min	3
HR ≥ 95 beats/min	5

Simplified Geneva Rule	Points
Age > 65 years	1
Surgery or fracture within 1 month	1
Active cancer	1
Unilateral lower limb pain	1
Hemoptysis	1
Pain on deep palpation of lower limb and unilateral edema	1
HR 75-94 beats/min	1
HR > 94 beats/min	2

- Using **modified** score:
 - <3 points indicates low probability
 - 4-10 points indicates intermediate probability
 - >10 points indicates high probability.
- Using **simplified** score, ≤2 points indicates that PE is unlikely.

PE risk prediction scores

1 The Pulmonary Embolism Severity Index and the Simplified versions

The Pulmonary Embolism Severity Index (PESI) is a risk stratification tool to determine the mortality of patients with newly diagnosed pulmonary embolism (PE). It supports physicians in identifying those patients who could potentially be treated as outpatient

The Simplified Pulmonary Embolism Severity Index (sPESI) was designed to remove some of the more complicated elements of the Pulmonary Embolism Severity Index (PESI) and aid in the risk stratification of patients with pulmonary embolism (PE).

Parameter	Original version	Simplified version
Age	Age in years	1 point (if age >80)
Male sex	+ 10 points	-
History of Cancer	+ 30 points	1 point
History of heart failure	+ 10 points	-
History of chronic lung disease	+ 10 points	1 point
Pulse rate ≥ 110	+ 20 points	1 point
Systolic BP <100 mmHg	+ 30 points	1 point
RR > 30 breath/min	+ 20 points	-
Temperature <36 °C	+ 20 points	-
Altered mental status	+ 60 points	-
Arterial Hb saturation < 90%	+ 20 points	1 point

Original Ver:

Class 1: ≤ 65 points

very low mortality risk

Class2: 66-85 points

low mortality risk

Class3: 86-105 points

moderate mortality risk

Class4: 106-125 points

high mortality risk

Class5: >125 points

very high mortality risk

- Class I and II defined as low risk.

Simplified Ver:

0 point= low risk
30 day mortality risk 1%

≥ 1 point = high risk
30 day mortality risk 10.9%

2

Hestia criteria (Yes\ No Questions)

If any of the following Qs answered with [Yes]; Patient need to be admitted

- Is the patient hemodynamically unstable?
- Is the thrombolysis or embolectomy necessary?
- Active bleeding or high risk for bleeding?
- >24 h of O₂ supply to maintain O₂ saturation >90%?
- Is PE diagnosed during anticoagulant treatment?
- Severe pain needing I.V pain med for > 24h?
- Medical or social reasons for treatment in hospital >24h (infection, malignancy, no support system?)
- Does the patient have a creatinine clearance < 30mL/min?
- Does the patient have severe liver impairment ?
- Is the patient pregnant?
- Does the patient have a documented history of HIT?

Diagnosis

- The diagnosis of massive PE should be explored whenever oxygenation or hemodynamic parameters are severely compromised without explanation.
- The diagnosis of pulmonary embolism may be aided by asking three questions:
 1. Is the clinical presentation consistent with PE?
 2. Does the patient have risk factors for PE?
 3. Are there any alternative diagnoses that can explain the patient's presentation?

Diagnosis Methods

Important to know the order of diagnosis:
 1- CXR 2- ECG 3- ABG
 We do not do advance investigations right away unless it is unstable case and there is high suspicion of




- There is no single, uncomplicated diagnostic test for a PE. Chest x-ray, EKG, and ABG are the **best initial tests**. **Angiography** is the most accurate test, but can be fatal in 0.5% of cases. After doing an ABG, chest x-ray, and EKG, the “best next step” is **most often a CT angiogram**.
- When the Wells score indicates that a PE is unlikely, a negative D-dimer excludes the diagnosis without further investigation. However, imaging is essential in those who either have a raised D-dimer level or are categorized as likely to have a PE based on their Wells score.
- Some tests are suggestive of PE (CXR, ECG, ABG) we do them first to lead us to PE → Then we do the Diagnostic test which is (CT angio AKA Spiral CT) . Pulmonary Angio is also diagnostic but it's not used anymore because it's time consuming and there are some risks involved.

◀ **CXR:** You have to do basic investigation, CXR is important.

CXR usually normal.

- Atelectasis¹ or pleural effusion may be present.
- The **main usefulness is in excluding** alternative diagnoses, e.g. pneumonia or pneumothorax.
- Classic radiographic signs, such as Hampton hump or Westermark sign, are rarely present.
- Infarction is not that common because the lung has dual supply (supplied by pulmonary and aortic vessels).

Pulmonary infarct in right lower lobe



A: Wedge infarct
 B: Raised hemidiaphragm

Findings	COPD % (n=21)	No prior cardiopulmonary disease, %(n=117)
Atelectasis or pulmonary parenchymal abnormality	76	68
Pleural effusion Very common	52	48
Pleural-based opacity	33	35
Elevated diaphragm	14	24
Decreased pulmonary vascularity	38	21
Prominent central pulmonary artery	29	15
Cardiomegaly	19	12
Westermark's sign	5	7
Pulmonary edema	14	4

1- Atelectasis is the collapse of some alveoli, which results in a decrease in lung volume.

◀ ABG¹


- Significant hypoxemia is almost uniformly present when there is a hemodynamically significant PE.
- ABG levels are not diagnostic for PE
 - PaO₂ and PaCO₂ are low (the latter due to hyperventilation) and pH is high; thus, there is typically a respiratory alkalosis.
 - A more accurate test is to measure alveolar arterial gradient [A-a gradient] (PAo₂)-(Pao₂), (Alveolar gas equation). Normal gradient is between 5-10, A-a gradient is usually elevated 15 or above. A normal value makes PE less likely, but cannot be relied on to exclude the diagnosis. A-a gradient is usually elevated.

◀ ECG:²



The ECG is an advance level for you


Rt. Bundle Branch Block



Findings:

- Lead 1: S wave
- V1,2: Slurred R wave
- There are 2 types of RBBB:
 - 1- Complete RBBB: QRS duration is prolonged
 - 2- Incomplete RBBB: Normal QRS duration (<0.12s or 4 small squares)

S1 Q3 T3 Pattern



Findings:

- You see the S wave in lead 1 and the Q and T waves in lead 3
- This finding is helpful but not diagnostic for PE

Rt. Ventricular Strain



Findings:

- ST-segment depression
- T-wave inversion

T-wave inversion suggests ischemia and stress



Findings:

- T-wave inversion in V1,V2,V3

1. People suffering from pulmonary embolism may be tachypneic (they wash out CO₂). Boyle's law states (In a closed chamber, if one gas decreases the other will increase). An alveolus is a closed chamber. Normal O₂ saturation does not confirm that the patient is normal, as they can have normal or high PO₂ but low PCO₂ levels (normal range is about 35 to 45 mmHg). "YOU WILL NOT BE ASKED ABOUT IT"

2.a. Not diagnostic but has things that are suggestive.


2.b. Usually shows sinus tachycardia. The most common abnormality is nonspecific ST-T wave changes. Only 5% will show right axis deviation, RV hypertrophy or right bundle branch block.

Computed tomographic pulmonary angiography (CTPA) ★

(Previously Spiral CT) **(first-line diagnostic test)** The Gold Standard

- **Multidetector computed tomography (MDCT)** is considered the standard imaging modality for diagnosing PE.
- Technical advancements in CT in the past decade have allowed physicians to obtain images of higher quality, with exposure to lower levels of radiation, which demonstrate small subsegmental pulmonary artery embolisms.
- **Dual-energy computed tomography (DECT)** can provide both morphological and functional pulmonary information of the lung in a single contrast-enhanced examination. It uses X-rays with two different energy spectra to detect specific substances according to the material decomposition theory.
- Data suggest that a negative Spiral CT is an adequate criterion for excluding PE in patients with a non-high clinical probability of PE.
- Has been found to have good sensitivity (>90%) and specificity.
- Can visualize very small clots (as small as 2 mm); may miss clots in small subsegmental vessels (far periphery).
- **It is the standard of care in terms of diagnostic testing to confirm the presence of a PE after the x-ray, EKG, and ABG are done.**

- In combination with clinical suspicion, guides treatment
- CTA **cannot** be performed in patients with significant renal insufficiency because of the IV contrast that is required.



V/Q (ventilation perfusion scan)³ ★

- **This test is performed in two stages:**
 1. **perfusion phase:** in which technetium-labelled albumin aggregates are injected intravenously and blood flow to the lungs is assessed.
 2. **ventilation phase:** in which patients inhale radiolabelled xenon or technetium to assess air delivery to the lungs.
- ★ Traditionally, this was the most common test used when PE is suspected, but has been replaced by CT angiography (CTA) as the initial study of choice in many medical centers.
- ★ Plays an important role in diagnosis when there is a contraindication to CTA. (This technique has the advantage of a lower radiation dose and is preferred in those with renal impairment and allergies to intravenous contrast agents.) May be useful when the chest x-ray is clear and when there is no underlying cardiopulmonary disease.
- V/Q is USED first only in pregnancy.
- The lungs of a young woman who is postpartum, or has had trauma, or pelvic surgery are normal (no disease no fibrosis no COPD). When she inhales, you will expect all of her alveoli to be normal. However, when you inject the radioactive compound, you will find some defects (areas of clot will not have blood flowing to them), resulting in a mismatch between perfusion and ventilation.

1- Lung infarction is rarely seen.

2- The aorta is usually larger than the pulmonary artery. However, the pulmonary artery became bigger in size as a result of the clot. Because the pulmonary artery has thin walls, it can be easily widened.

3- Not used for COPD patients, also it's rarely used now (time consuming).

The use of ventilation perfusion scan in diagnosing pulmonary embolism

- Interpretation of results, can be either
 - normal
 - low probability
 - intermediate probability
 - or high probability (treatment guidelines based on PIOPED study).
 - A high-probability V/Q scan has a very high sensitivity for PE; treat with heparin.
 - If there is low or intermediate probability, clinical suspicion determines the next step. If clinical suspicion is high, pulmonary angiography is indicated. Alternatively, perform a lower extremity duplex ultrasound to avoid pulmonary angiography. If the duplex is positive, treatment for DVT is the same as for PE. If the duplex is negative/uncertain, then pulmonary angiography is indicated to exclude PE.
- A diagnosis of PE is made if there are mismatched defects in the perfusion scan, indicating impaired blood flow to the lungs, but normal ventilation because air entry to the lungs is normal.
- Normal scan virtually rules out PE—no further testing is needed but a scan is almost never “normal” in anyone.

Prospective Investigation of Pulmonary Embolism Diagnosis (PIOPED) results:


Scan category	PE present	PE absent	PE uncertain	No angiogram	Total
High probability	102	14	1	7	124
Intermediate probability	105	217	9	33	364
Low probability	39	199	12	62	312
Near normal or normal	5	50	2	74	131
Total	251	480	24	176	931

The use of ventilation perfusion scan in diagnosing pulmonary embolism

SKIPPED BY THE DOCTOR
Dr. You will not be asked about it

High probability

- =2 large segmental (>75% of a segment) perfusion defects without corresponding ventilation or radiographic abnormalities or substantially larger than matching ventilation or radiologic abnormalities.
- OR
- =2 moderate segmental (>25% and <75% of a segment) perfusion defects without matching ventilation or chest radiographic abnormalities plus one large unmatched segmental defect.
- OR
- =4 moderate segmental perfusion defects without matching ventilation or chest radiologic abnormalities.



Red arrows = clots

Intermediate probability

- Scans that do not fall into normal, very low, low, or high probability categories.

Low probability

- Non-Segmental perfusion defects.
- OR
- Single moderate mismatched segmental perfusion defect with normal chest radiograph.
- OR
- Any perfusion defect with a substantially larger abnormality on chest radiograph.
- OR
- Large or moderate segmental perfusion defects involving no more than four segments in one lung and no more than three segments in one lung region with matching or larger ventilation/radiographic abnormalities.
- OR
- More than three small segmental perfusion defects (<25% of a segment) with a normal chest radiograph.

Very Low probability


- Three or fewer small segmental perfusion defects with a normal chest radiograph.
- Normal.
- No perfusion defects present.

Diagnosis

◀ D-Dimer¹

- The D-dimer is a degradation product of fibrinolysis and is increased in patients with acute venous thromboembolism (PE and DVT) as well other non-thrombotic disorders
- D-dimer is a sensitive (90% to 98%). If results are normal and clinical suspicion is low, PE is very unlikely, but not specific diagnostic test it may also be elevated in MI, CHF, pneumonia, and postoperative state (or in any state of acute inflammation). Any cause of clot or increased bleeding can elevate the D-dimer level
- A low clinical probability score is useful for excluding of venous thromboembolism
- If pretest probability of PE is low, D-dimer test is a good noninvasive test to rule out PE. So **if D-dimer is negative, you can rule out a clot. But if it is positive, this does not help you.**


◀ MRA real time




Comparison in the sensitivity of Spiral CT, MRA and RT- MRA

Reader	CT	MRA	RT-MRA
1	72.1	79.1	97.7
2	69.8	81.4	97.7
Mean	71	80.3	97.7
K	0.86	0.84	1


◀ MRA with contrast²



◀ Echo : TTE is used



-the RV is pushing the septum to the left
-returns to normal size after treatment (thrombolytics)



1.a. Measured by ELISA, important to know that if D-dimer positive then most likely the patient has PE, if D-dimer negative its less likely.


1.b. If negative you can exclude PE, but if positive it doesn't tell you anything.

2. MRA is simply MRI with angio.


◀ Pulmonary angiogram:

The most accurate test with nearly 100% specificity and a false negative rate under 1%.

- Definitively diagnoses or excludes PE, but is invasive. Contrast injected into the pulmonary artery branch after percutaneous catheterization of the femoral vein.
- Consider when noninvasive testing is equivocal and risk of anticoagulation is high, or if the patient is hemodynamically unstable and embolectomy may be required. Angiography is rarely performed because it carries a 0.5% mortality.
- Pulmonary angiogram is not used now cause it's time consuming , CT angiogram is much better



◀ Suggested diagnostic strategy for venous thromboembolism¹:



If you suspect PE and there are no contraindications to anticoagulation, start treatment right away. If the results of the investigation are -ve, the treatment should be stopped. Starting early is better than starting late because starting late can increase mortality.

1- The boxes with green borders were covered in the dr slides with (SPIRAL CT SCAN OF THE CHEST).

* After you do Hx, ABG, ECG, D-dimer, CXR. Do CT angio

* If CT is +ve --> treat.

* If -ve --> Do ultrasound for the leg

If thrombus present --> treat (follow the left side of the algorithm)

If no thrombus (follow the right side of the algorithm)

Treatment¹

- The main principle of treatment for PE is anticoagulant. Anticoagulation prevents further clot formation, but does not lyse existing emboli or diminish thrombus size
- **Prompt** initiation of anticoagulation while awaiting investigations is prudent because of the high risk of early mortality with untreated pulmonary embolism

Anticoagulants

- Direct Oral Anticoagulants [DOACs]² are given at fixed dose and **do not necessitate routine laboratory monitoring**.
- Each DOAC has been deemed non-inferior to the Vitamin K Antagonist \ Low molecular weight Heparin [VKA/LMWH] combination for the prevention of symptomatic recurrent venous thromboembolism in patients with an acute venous thromboembolism.
- **DOACs have significantly fewer major bleeding events compared with VKAs**

DOACs: We don't use warfarin as first choice. Know the DOACs names!

Drug	Target	Peak effect	Half life	Renal clearance	Protein binding
Dabigatran	Factor IIa (Thrombin) Selective thrombin antagonist	1.5 h	14-17 h	> 80% ³	35%
Apixaban	Factor Xa Selective and direct inhibition of factor Xa	3 h	8-14 h	25%	85%
Edoxaban		4 h	8-11 h	35%	55%
Rivaroxaban		2-3 h	7-11 h	33%	90%

- Rivaroxaban 15 mg and 20 mg tablets should be taken with food for maximum absorption and efficacy.
- **What agents reverse anticoagulation?**
- Andexanet alfa reverses rivaroxaban, apixaban, and edoxaban.
- Idarucizumab reverses dabigatran.
- Prothrombin complex concentrate (PCC) reverses warfarin

Dosage and monitoring of anticoagulant therapy:

1

Warfarin

- 5 mg/d can be started on day 1 of therapy; there is no benefit from higher starting doses
- Platelet count should be monitored at least every 3 d during initial heparin therapy
- Therapeutic APTT should correspond to plasma heparin level of 0.2–0.4 IU/mL
- Requires regular monitoring of **INR** (individual's prothrombin time)

2

UFH, LMWH⁴

- is usually continued for 5–7 d
- Can be stopped after 4–5 d of warfarin therapy when INR is in 2.0– 3.0 range

Heparin is administered as either unfractionated (IV) or LMWH. The issue with UFH is that it is impossible to predict how much of it will be protein bound. A small-sized lady can require a high dose, whereas a large-sized man can require a low dose, because the lady's heparin may be protein-bound. The man, on the other hand, requires a small dose because the majority of the heparin will be absorbed by the blood. As a result, it is unpredictable and impractical. We make use of LMWH.

1- When PE is diagnosed, mortality is 10% in the first 60 minutes. Of those who survive the initial event, approximately 30% of patients will die of a recurrent PE, if left untreated. Most deaths are due to recurrent PE within the first few hours of the initial PE. Treatment with anticoagulants decreases the mortality to 2% to 8%.

2- **DOAC** They are anticoagulant better than warfarin and heparin because it has less adverse effect. The dose response is predictable and they have less risk of bleeding compared to heparin.

3- Avoid in renal failure.

4- Start heparin first due to its fast action then warfarin + monitor INR carefully. if INR is stable, continue warfarin and stop heparin.

Treatment

◀ Important drug interactions with warfarin:¹

Don't memories this when you prescribe a medication just open your phone and see if there is DDI

Drugs that decrease warfarin requirement

- Second- and third-generation cephalosporins
- Phenylbutazone
- Metronidazole
- Trimethoprim-sulfamet hoxazole
- Amiodarone
- Clofibrate
- Erythromycin
- Anabolic steroids
- Thyroxine

Drugs that increase warfarin requirement

- Barbiturates
- Carbamazepine
- Rifampin
- Penicillin
- Griseofulvin
- Cholestyramine

◀ Complications:



Heparin

Important to understand

Complication	Management
Bleeding	<ol style="list-style-type: none"> 1. Stop heparin infusion. 2. For severe bleeding, the anticoagulant effect of heparin can be reversed with intravenous protamine sulfate 1 mg/100 units of heparin bolus or 0.5 mg for the number of units given by constant infusion over the past hour; provide supportive care including transfusion and clot evacuation from closed body cavities as needed.
Heparin-induced thrombocytopenia and thrombosis ² aka HIT syndrome or White-clot syndrome	<ul style="list-style-type: none"> • Carefully monitor platelet count during therapy; Stop-heparin for platelet counts <75,000. • Replace heparin with direct inhibitors of thrombin like desirudin if necessary. These agents do not cause heparin-induced thrombocytopenia. • Avoid platelet transfusion because of the risk for thrombosis.
Heparin-induced osteoporosis (therapy >1mo)	<ul style="list-style-type: none"> • LMWHs may have lower propensity to cause osteoporosis as compared with unfractionated heparin; consider LMWH if prolonged heparin therapy is necessary

1- You don't need to know the names of the drugs just know that it has a lot of interactions. if the pt is using any of them ask to monitor INR after few days of initiation of treatment.

2- Characterized by activation of fibrinogen which in turn will form A mesh that will trap platelets. Lab investigation will show very rapid fall in Platelets count, that's why you have to test platelet count in the first week of administration.

Treatment



Warfarin

Complication	Management
Bleeding	<ol style="list-style-type: none"> 1. Stop therapy. 2. Administer vitamin K and fresh frozen plasma for severe bleeding; provide supportive care including transfusion and clot evacuation from closed body cavities as needed
Skin necrosis (rare)	<ul style="list-style-type: none"> • Supportive care
Teratogenicity	<ul style="list-style-type: none"> • Do not use in pregnancy or in patients planning to become pregnant.

Thrombolytics¹

Approved thrombolytics for pulmonary embolism :

1

Recombinant tissue-plasminogen activator²

100 mg as a continuous peripheral intravenous infusion administered over 2 h

2

Streptokinase

250,000 IU as loading dose over 30 min, followed by 100,000 U/h for 24 h

3

Urokinase

4400 IU/kg as a loading dose over 10 min, followed by 4400 IU/kg/h for 12-24 h



Indications in PE: ★

- When are thrombolytics the right answer?
 - Hemodynamic instability³.
 - Hypoxia on 100% oxygen.
 - Right ventricular dysfunction by echocardiography.
- For **massive PE use thrombolytic + anticoagulant**. Any other PE use anticoagulant only (Thrombolytic doesn't show any improvement in mortality in these cases).
- Diuretics and vasodilators should also be avoided, as they will reduce cardiac output.

Massive PE = [Shock] Hypotension + Hypoxemia
Submassive = RV Dilation without shock



Contraindications

Relative:

- Recent surgery within last 10 d
- Previous arterial punctures within 10 d
- Neurosurgery within 6 mo
- Bleeding disorder (thrombocytopenia, renal failure, liver failure)
- Ophthalmologic surgery within 6 wk
- Hypertension >200 mm Hg systolic or 110 mm Hg diastolic
- Placement of central venous catheter within 48 h
- Hypertensive retinopathy with hemorrhages or exudates
- Intracerebral aneurysm or malignancy
- Cardiopulmonary resuscitation within 2 wk
- Cerebrovascular disease
- Major internal bleeding within the last 6 mo
- Pregnancy and the 1st 10 d postpartum
- Infectious endocarditis
- Severe trauma within 2 mo
- Pericarditis

Absolute:

- Active internal bleeding


1- Why don't we give thrombolytic to a stable patient with PE ? because if you take a patient with a stable PE and you give them anticoagulant + thrombolytic you will not have the best outcome. Giving one anticoagulant will be better and it will decrease the risk of bleeding.

2- Recombinant tissue-plasminogen activator is the one we're using currently.


2- (e.g., hypotension [systolic BP <90 mm Hg] and tachycardia)

Treatment


Treatment phases : Skip doses



- BID: twice daily. INR: international normalized ratio > LMWH: low molecular weight heparin
- *LMWH is needed for 5-10 days before starting dabigatran or edoxaban
- ~ 30 mg daily if creatinine clearance is 30-50 mL/min or weight < 60 kg
- “Dose reduction may be considered after 6 months of therapy



Identify category of Venous Thromboembolism [VTE] event



Outpatient vs inpatient therapy :

- RCTs have compared outpatient versus inpatient management of pulmonary embolism and found no difference in outcomes in selected patients.

Conclusions:


- PE is common and under-recognized serious medical problem
- Early diagnosis and treatment is essential for good outcome
- High index of suspicion is needed in high risk patients



Figure 1: CT pulmonary angiography shows multiple pulmonary emboli in a 63-year-old male with chest pain for 7 days. 2D axial images show presence of thrombus in the main pulmonary arteries with extension to the segmental and subsegmental pulmonary arterial branches on both sides resulting in filling defects (A-C). Coronal reformatted image demonstrates multiple emboli on pulmonary arteries (D). Arrows indicate the thrombi in pulmonary arteries.

Interventional Cardiology (2017) Volume 9, Issue 5


Fig. 1. A 76-year-old female diagnosed with acute pulmonary thromboembolism.



A 76-year-old female diagnosed with acute pulmonary thromboembolism.
A. CT angiography showing a focal filling defect in the right upper lobar pulmonary artery (arrow) due to acute pulmonary thromboembolism and bilateral pleural effusion
B. The fusion image of CT angiography and a color-coded iodine map showing a thrombus in the right upper pulmonary artery as well as a corresponding wedge-shaped perfusion defect in the right upper lobe.

Korean J Radiol 2021 Sep;22(9):1555-1568. <https://doi.org/10.3348/kjr.2020.1512>

Fig. 3. A 58-year-old female with pulmonary artery sarcoma.



A 58-year-old female with pulmonary artery sarcoma.
A. A coronal contrast enhanced CT image showing an intraluminal mass with an eccentric filling defect in the right main pulmonary trunk.
B. On a coronal color-coded iodine (water) image from dual-energy CT, the mean iodine concentration within the region of interest is 2.6 mg/mL.
C. A coronal PET image showing focal fluorodeoxyglucose uptake in the intraluminal mass and right upper paratracheal lymph nodes.

Korean J Radiol 2021 Sep;22(9):1555-1568. <https://doi.org/10.3348/kjr.2020.1512>



- The most important side effect of all oral anticoagulants is a **dose-dependent increase in bleeding risk**
- **DRAW: Dabigatran, Rivaroxaban, Apixaban, and Warfarin** are the most important oral anticoagulants.
- RivaroXaban, apiXaban, and edoXaban are factor Xa inhibitors.
- **WARsaw** is an **EXTRAordinary Place To check out: WARfarin** affects the **EXTRinsic** pathway; therefore, PT should be regularly checked.





Fig. 29.7 Models for anticoagulant management of newly diagnosed venous thromboembolism (VTE). *The decision to stop or continue anticoagulation beyond 3 months is based on a balance of risks of VTE recurrence if anticoagulants are stopped, bleeding if anticoagulants continue and patient preference. INR, International Normalized Ratio; LMWH, low-molecular-weight heparin.




Features of Pulmonary thromboemboli:

	Acute massive PE	Acute small/medium PE	Chronic PE
Patho-physiology	<p>Major haemodynamic effects:</p> <ul style="list-style-type: none"> - ↓cardiac output - acute right heart failure 	Occlusion of segmental pulmonary artery → infarction ± effusion	<ul style="list-style-type: none"> - Chronic occlusion of pulmonary microvasculature - Right heart failure
Symptoms	<ul style="list-style-type: none"> - Faintness or collapse - Crushing central chest pain - Apprehension - Severe dyspnoea 	<ul style="list-style-type: none"> - Pleuritic chest pain - Restricted breathing - Haemoptysis 	<p>Exertional dyspnoea</p> <p>Late:</p> <ul style="list-style-type: none"> - Symptoms of pulmonary hypertension - Right heart failure
Signs	<p>Major circulatory collapse:</p> <ul style="list-style-type: none"> - Tachycardia - Hypotension - ↑JVP - RV gallop rhythm - Loud P2 - Severe cyanosis - ↓urinary output 	<ul style="list-style-type: none"> - Tachycardia - Pleural rub - Raised hemidiaphragm - Crackles - Effusion (often blood-stained) - Low-grade fever 	<p>Early:</p> <ul style="list-style-type: none"> - may be minimal <p>Later:</p> <ul style="list-style-type: none"> - RV heave, loud P2 <p>Terminal:</p> <ul style="list-style-type: none"> - signs of right heart failure
Chest X-ray	Usually normal; may be subtle oligoemia	<ul style="list-style-type: none"> - Pleuropulmonary opacities - Pleural effusion - Linear shadow - Raised hemidiaphragm 	<ul style="list-style-type: none"> - Enlarged pulmonary artery trunk - Enlarged heart - Prominent right ventricle
ECG	<ul style="list-style-type: none"> - S1Q3T3 anterior T-wave inversion - RBBB 	Sinus tachycardia	RV hypertrophy and strain
Arterial blood gas	<ul style="list-style-type: none"> - Markedly abnormal with ↓PaO₂ and ↓PaCO₂ - Metabolic acidosis 	May be normal or ↓PaO ₂ or ↓PaCO ₂	Exertional ↓PaO ₂ or desaturation on formal exercise testing
Alternative diagnosis	<ul style="list-style-type: none"> - Myocardial infarction - Pericardial tamponade - Aortic dissection 	<ul style="list-style-type: none"> - Pneumonia - Pneumothorax - Musculoskeletal chest pain 	Other causes of pulmonary hypertension

Summary

PE is a medical emergency! Early diagnosis and management is crucial for reducing mortality



Clinical features :

Sudden onset dyspnea

Pleuritic chest pain

Hemoptysis

Investigation :

- First step is to determine whether the patient is stable or not .
- The following investigative guidelines should be followed based on patient risk :

Summary from Davidson's




Fig. 29.3 Investigation of suspected PE without haemodynamic compromise. *In selected cases where clinical suspicion remains high it may be necessary to consider either bilateral lower limb ultrasonography or V/Q scan.

Pharmacological treatment :

Massive PE	Any other PE
<ul style="list-style-type: none"> • thrombolytics+anticoagulant • Recombinant tissue-plasminogen activator • Streptokinase • Urokinase 	<ul style="list-style-type: none"> • anticoagulants • Heparin • Warfarin • NOACs (eg. Dabigatran , Rivaroxaban)

Lecture Quiz

Q1: A 41-year-old woman is referred for assessment after suffering a second pulmonary embolus within a year. She has not been travelling recently, has not had any surgery, does not smoke and does not take the oral contraceptive pill. She is not currently on any medication as the diagnosis is retrospective and she is now asymptomatic. What should be the next step in her management ?

- A- Initiation of warfarin therapy
- B- ECG
- C- Thrombophilia screen
- D- Insertion of inferior vena cava filter
- E- Duplex scan of lower limb veins and pelvic ultrasound

Q2: A 60-year-old woman is receiving chemotherapy for metastatic breast cancer. A staging computed tomographic scan reveals multiple pulmonary emboli in segmental arteries. She is asymptomatic. Which of the following treatments is the best?

- A- Give dalteparin and transition her to warfarin.
- B- Give dalteparin and continue it at least as long as she is receiving chemotherapy.
- C- Give rivaroxaban.
- D- Give warfarin.
- E- No therapy.

Q3: A 47-year-old obese Asian man complains of a sharp pain on the left side of his chest with difficulty breathing. The pain started a few hours ago and does not radiate anywhere, the patient also reports feeling increasingly short of breath and became extremely anxious when he started coughing blood-stained sputum. He states he has been flying all week on business trips and is getting late for his next flight. The most likely diagnosis is ?

- A- Myocardial infarction
- B- Muscular injury
- C- Pneumothorax
- D- Pulmonary embolism

Q4: A 38-year-old previously healthy woman presents to the emergency department with a 7-day history of low-grade fever, shortness of breath on exertion, and nonproductive cough. Her heart rate is 90 beats/min, her oxygen saturation on room air is 98%, and her chest radiograph is normal. Which of the following tests would help to exclude the diagnosis of pulmonary embolism?

- A- Bilateral compression ultrasound examination of the lower extremities
- B- D-dimer
- C- Electrocardiogram
- D- Computed tomographic pulmonary angiogram
- E- Ventilation-perfusion lung scan

Q5: A 59-year-old obese woman underwent a coronary artery stent procedure. She is a well-controlled type 2 diabetic. The operation was successful. However, after 1 week during recovery, the patient complained of severe chest pain and shortness of breath. Her heart rate was 115 bpm and blood pressure 107/89 mmHg. Following resuscitation of the airway, breathing and circulation, an electrocardiogram (ECG) showed sinus tachycardia and right axis deviation. The most appropriate treatment is ?

- A- Warfarin
- B- Intravenous adrenaline
- C- Alteplase
- D- Salbutamol
- E- Intravenous heparin

Our Team

This work was originally done by **438 Medicine team.**

Edited by **439 Medicine team:**

Team Leaders

- *Shaden Alobaid*
- *Ghada Alabdi*

- *Hamad Almousa*
- *Naif Alsulais*



Member : Raghad albarrak

Edited for the second time by **442 Medicine team:**

Team Leaders

- *Mohammed Alrashoud*
- *Maha Alzahrani*
- *Shatha Alshabani*

- *Mohammed Alzeer*
- ***Refal Alamry***
- *Arwa Alghamdi*



Members: Nourah Alfawaz
Saif Alotaibi



Medicine.med442@gmail.com

Special thanks to Mohammed Alorayidh and Arwa Alghamdi for the amazing first page theme!