

Microbiology team

MSQ

Done by : Aljoharah AL-Rashodey

Amjad AL-Turki

- 1) Direct microscopy is used in the diagnosis of :
 - a) **M tuberculosis**
 - b) Septicemia
 - c) Staphylococcus aureus
 - d) Diphtheria

- 2) In treatment of tuberculosis all these drugs except :
 - a) ethambutol
 - b) pyrazinamide
 - c) isoniazide
 - d) rifampicin
 - e) **Penicillin**

- 3) BCG :
 - a) serum
 - b) **Live attenuated vaccine**
 - c) Killed bacteria vaccine
 - d) a protein from bacteria
 - e) a toxin

- 4) Which one of these mycobacteria cause AIDS :
 - a) M kansasii
 - b) M avium intracellulare
 - c) M scrofulaceum
 - d) **M tuberculosis**
 - e) M bovis

- 5) Duration of treating open TB :
- a) 2 months
 - b) 4 months
 - c) 6 months
 - d) 18 months
 - e) 2 years
- 6) The time estimated for inoculation of TB is :
- a) 2 years
 - b) 2 weeks
 - c) 4 weeks
 - d) 8 weeks
- 7) Which media is used for TB :
- a) Lowenstein_jensen
 - b) Mconkey
 - c) Blood agar
 - d) Z/N
- 8) Which of the following are endogenous (intracellular) :
- a) pneumoniae
 - b) agalctae
 - c) pyogenic
 - d) diphtheria
 - e) T.B.
- 9) Primary tuberculosis :
- a) Non infectious
 - b) Open lesion
 - c) In adults
- 10) Secondary tuberculosis :
- a) In apex of lung
 - b) Non infectious
 - c) In the middle lobe

11) The virulence factor of TB is :

- a) M serotype
- b) Extracellular organism
- c) Capsulated
- d) Multiply in macrophage

اللهم اني استودعك محافظته ومآقراته ومافهمته وماتعلمته فرده عند حاجتي إليه

اللهم لا سهل إلا ما جعلته سهلا وانت تجعل الصعب إذا شئت سهلا