

Respiratory Block

Immunology

Lecture 3

Immunology of Asthma.

In this document you will find some main points gathered from the 1st lecture..This document is NOT a replacement for the lecture..If you need additional information go back to the lecture or use a book as a reference so you understand everything correctly.

Hopefully all the information is correct and Hope you find them useful.

Good Luck to everyone.

430 Immunology Team Members

Abdelelah AlFreem AlEnazi

Alaa AlHumaid

Hadeel F. AlSajjan

Hanan AlSalman

Lujain AlYousef

Mohammed AlAnazi

Nora AlMfgaie

Immunology

Immunology of Asthma.

Some main points you can go through and revise from:

Note:

People with asthma have a very sensitive air way can be triggered by any thing

Note:

Asthma → inflammation on the lung by an allergen or any disease

Asthma is a clinical syndrome characterized by :


- Episodes of reversible airway obstruction. #(important in diagnosing asthma)
- Increased bronchial reactivity.
- Airway inflammation

Patients with asthma present with one or more of the following symptoms:

- breathlessness (difficulty in breathing).
- Wheezing .
- persistent cough. (the common symptoms of asthma)
- chest tightness

Note:

Reversible Airway Obstruction → (an important feature of asthma)



Classification of asthma :

- Extrinsic (atopic).#Main type
- Intrinsic (non-atopic).

Note:

Atopy → A genetic tendency to develop allergy

Non-atopic (intrinsic) asthma (10-33% of asthmatics)

- ! Negative skin tests .
- ! No clinical/family history of allergy
- ! Serum [IgE] is normal .
- ! Older patients .
- ! More severe .

Atopic (extrinsic) asthma .

*Allergies trigger asthma attacks in :

- Children (60-90%)
- Adults .(50%)

Note:

Approximately 75-85% of patients with asthma have positive (immediate) skin test reactions to various allergens .

#Some Allergies get treated by the increase of hormones at puberty

Role of Allergens in Asthma

#Allergen sensitization is linked to the risk of developing asthma.

Indoor allergens #Patient would present with asthma all year long

- House dust mites.
- Domestic pets (cat fur & dander).
- Cockroaches (insects).
- Molds (fungal spores)

Outdoor allergens : #Patients present with asthma at certain times of the year


- Fungal spores :(e.g. Alternaria) .
- Grass, tree & weed pollens.

Antigen presenting cells (APC) in the lung :

#Cell mediated immunity

Two subsets of dendritic cells in the lungs.

- Respiratory tract myeloid DCs (mDCs) aid in the development of asthma symptoms → #Causes Allergy
- Plasmacytoid DCs (pDCs) aid in respiratory tolerance to allergens. → # Does not cause allergy



In susceptible individuals :


first encounter with allergens activate B-cells to produce IgE.

Subsequently:

inhaled allergens activate submucosal mast cells in the lower airways .


mediators are released within seconds & lead to :

1. bronchoconstriction.
2. Influx of eosinophils & other inflammatory cells.



Asthma results from complex interactions of among the

Inflammatory cells and :




1. Airway epithelium


2. Nervous system #Mast cells stimulate it

3. Bronchial smooth muscles

Factor contributing to airflow obstruction leading to difficulty in breathing



Response to allergen occurs in two phases.



Early allergic response :

- Occur within minutes.
- manifest clinically as :
 - bronchial constriction .
 - airway edema .
 - mucus plugging .

*respond to bronchodilators .

Late allergic response :

- Appear 4 to 10 hours later.
- Result from infiltration by inflammatory cells.
- Activation of lymphocytes & Eosinophils .

*respond to steroids. (anti-inflammatory drugs).

Th2 cells and role of cytokines in allergic asthma.

Allergens drive T-cells towards TH 2 type :

TH2 secrete the cytokines : **IL-4, IL-5, IL-9 & IL-13** which Promote :

- B-cell IgE production.
- eosinophil attraction.
- airway inflammation .
- increased bronchial reactivity .

Role of IL-4 in allergic asthma.


The main role of IL-4 is carried out during the initial priming of Th2 cells:

- Regulate isotype switching in B cells to IgE. #When B-Cells are stimulated by IL-4 they form IgE
- Induce MHC II on antigen-presenting cells .
- Induce adhesion molecule expression.
- Activate mast cells and eosinophils .

Role of IL-13 in allergic asthma .

- IL-13 induces inflammation.
- Stimulate mucus hypersecretion.
- Induce subepithelial fibrosis. # The fibroses of the tissue is irreversible and will deposit in the bronchi which leads to the narrowing of the air way
- stimulate eotaxin production.

IL-4 & IL-13 act on: Eosinophils. Mast cells, B-cells, fibroblasts and airway smooth muscles.



Role of IL-5 in allergic asthma .

- IL-5 induces an increase in eosinophil production in the bone marrow.
- release of eosinophils from the bone marrow into circulation .

Role of eosinophils in allergic asthma .

- Eosinophils play a prominent role in initiating asthmatic symptoms by causing tissue damage in the airways of the lungs.

Role of IL-10 in allergic asthma.


* IL-10 regulate allergic responses

- 1.IL-10 has direct inhibitory effects on APCs and T cells .
#Inhibits the allergic reaction
- modulates eosinophil accumulation in airways by possibly inhibiting eosinophil production in bone marrow.

#Asthmatic patients have a decreased expression of IL-10 .

Role of NKT cells in asthma .

- NKT cells constitute less than 1% of CD4+ T cells in the blood and constitute 63% of CD4+ T cells in patients with moderate-to-severe asthma.
- NKT cells activate Th2 cells to secrete IL-4 and IL-13.



Role of regulatory T – cells :

- Regulatory T cells suppress the effector mechanisms that induce asthmatic symptoms.
- Asthmatics may lack functional regulatory T cells that can inhibit an asthmatic response.

*Activation of inflammatory cells (mast cells , eosinophils etc,) is a major inducer of airway inflammation.

*Airway inflammation is the hallmark in the asthmatic lung .

Airway inflammation leads to :

- Increased bronchial reactivity.

Products of the inflammatory cells act on :

- Airway smooth muscle cells . #Causes hypertrophy
- Lung fibroblasts .
- Mucous glands .

Products of the inflammatory cells cause :


- Airway remodeling.

Products of these cells acts on:


- 1- Air smooth muscle cells.
- 2- Lung fibroblasts.
- 3- Mucous glands.

Which will cause → **Airway remodeling**

Airway remodeling could cause an irreversible lung damage by:



airway remodelling refer to .



1. Smooth muscle hyperplasia & hypertrophy

2. Mucous gland hyperplasia

3. Collagen deposition

5. Chronic inflammation

4. Fibroblast activation

Inflammatory cells & their mediators



Airway Inflammation



Increased bronchial reactivity



airway remodelling

Outcome of airway remodelling



Can ultimately lead to fibrosis and irreversible airway obstruction in some patients .

Outcome of increased airway reactivity



predispose patients to develop asthma attacks
on exposure to non-specific irritants :

1. chemical irritants
2. smoke & strong perfumes.
3. sulphur dioxide & air pollutants
4. viral and bacterial respiratory infections.



If the patient
with
airway
reactivity is
exposed to any
of this triggers
he/she will
develop asthma

Summary:

- Asthma is characterized by episodic reversible
- airway obstruction
- Classified in 2 types : intrinsic & extrinsic.
- In the extrinsic type, allergens drive T-cells
- into TH2 pattern
- Airway inflammation is a hallmark finding in
- the asthmatic lung
- Inflammatory cells lead to increased bronchial
- reactions & airway remodeling