



Cardiovascular Physiology

Contractile Mechanism in Cardiac Muscle

Dr. Abeer A. Al Masri

MBBS, MSc, PhD
Assistant Professor
Consultant Cardiovascular Physiologist
College of Medicine, KSU



At end of this lecture you should be able to:

- ✓ Define cardiac muscle contractility
- ✓ Describe the mechanism of 'Excitation-Contraction Coupling'
- ✓ Understand the mechanism of isovolumetric & isometric contraction
- ✓ Factors affecting cardiac contractility
- ✓ Understand what's meant by inotropics (+ve & -ve)



PROPERTIES OF THE CARDIAC MUSCLE

- I. Excitability
- II. Conductivity
- III. Contractility
- IV. Rhythmicity

3



Contractility:



Is the force of contraction for a given fiber length

- ☐ Or/ is the ability of cardiac muscle to convert chemical energy into mechanical work
- ☐ Its essential for the pumping action of the heart

4


Excitation-Contraction Coupling:

The mechanism by which the action potential triggers contraction of the myofibrils

- ❑ Is almost similar to that of skeletal ms
- ❑ It depends more on ECF $[Ca^{2+}]$

5

Excitation-Contraction Coupling in Heart Muscle



6



In resting muscle fibers:

- ❑ Ca^{2+} is stored in the endoplasmic 'sarcoplasmic' reticulum (SR)
- ❑ Along the plasma membrane 'sarcolemma' there are membrane invaginations that form transverse tubules 'T tubules' (TT), which are long & tortuous
- ❑ TT terminate near the Ca^{2+} -filled sacs of SR

7




Excitation-Contraction Coupling:

- ❑ Initiated by impulse (AP) on the surface of myocardium
- ❑ Depolarization wave passes through TT
- ❑ Ca^{2+} enters through Ca^{2+} voltage channel in exterior membrane & dihydropyridine receptors (DHPRs) in TT
- ❑ Ca^{2+} entry triggers the release of Ca^{2+} from SR
- ❑ causes \uparrow in intracellular Ca^{2+}

8

Excitation-Contraction Coupling ... (Cont.)

- ❑ Ca^{2+} then binds to troponin C on thin filaments to start cross bridge cycle
- ❑ hydrolysis of ATP occurs to provide the energy for sliding of myosin over actin, causing contraction



9

Excitation-Contraction Coupling ... (Cont.)

- ❑ Relaxation occurs when Ca^{2+} is removed from troponin C:
 - Ca^{2+} will actively re-uptaken by the SR ... (using a Ca^{2+} ATPase)
 - Ca^{2+} will also pumped out of cell via Na^+ - Ca^{2+} exchanger
- ❑ Resulting in cessation of contraction until new AP occurs

10



Factors Affecting Myocardial Contractility: (Inotropic Effectors)

- 1. Cardiac innervation: (ANS)**
- 2. Oxygen supply**
- 3. $[Ca^{2+}]$ & $[K^+]$ ions in ECF**
- 4. Physical factors: (e.g. warming; cooling)**
- 5. Hormonal & Chemical factors**
- 6. Mechanical factors**

11



Factors Affecting Myocardial Contractility: (Inotropic Effectors)

- ☐ Inotropes are agents that \uparrow or \downarrow the force of muscular contraction:

Negatively Inotropic agents weaken the force of muscular contractions

Positively Inotropic agents increase the strength of muscular contraction

- ☐ Inotropes work via the autonomic nervous system (ANS)

12

How do they work?

- ❑ **'Ach'** is the neurotransmitter of the parasympathetic nervous system (PNS)
- ❑ **'NA'** is the primary neurotransmitter released by the sympathetic nervous system (SNS)
- ❑ **When SNS is stimulated, Noradrenaline & adrenaline levels ↑, leading to stimulation of adrenergic receptors that located in the endothelium:**
 - Alpha1 (α_1), Alpha2 (α_2)
 - Beta1 (β_1), Beta2 (β_2), &
 - Dopamine (D)

13

Adrenergic Receptors

Alpha Receptors		Beta Receptors		Dopamine
Alpha1 (α_1)	Alpha2 (α_2)	Beta1 (β_1)	Beta2 (β_2)	Dopamine (D)
Cardiac	In peripheral vessels	Heart	<ul style="list-style-type: none"> • Bronchial smooth muscles • Skeletal muscles 	
↑ Contractility w/out ↑ in rate	Mediate vasoconstriction	↑ Contractility w ↑ in HR	<ul style="list-style-type: none"> • Dilation of BSM • Vasodilation in SM • ? Some cardiac effects 	<ul style="list-style-type: none"> • ↑ Renal & ↑ Coronary blood flow • Arterial vasodilatation

14


Factors Affecting Myocardial Contractility:

1. Cardiac innervation: (ANS)

- Sympathetic NS → ↑ force of contraction
- Parasympathetic NS (vagus) → ↓ atrial force of contraction w no significant effect on ventricular ms

15


Mechanism of action of ANS on cardiac contractility



16

Effect of autonomic innervation on cardiac contractility

Affected by epinephrine (\uparrow) & ACh (\downarrow)



17

Factors Affecting Myocardial Contractility ... (Cont.)



2. Oxygen supply:

- Hypoxia \rightarrow \downarrow contractility

3. $[\text{Ca}^{2+}]$ & $[\text{K}^{+}]$ ions in ECF:

- $\uparrow \text{Ca}^{2+} \rightarrow \uparrow$ contractility
- $\uparrow \text{K}^{+} \rightarrow \downarrow$ contractility

18





Factors Affecting Myocardial Contractility ... (Cont.)

4. Physical factors:

- Warming → ↑ contractility
- Cooling → ↓ contractility

21



Factors Affecting Myocardial Contractility ... (Cont.)

5. Hormonal & Chemical factors:

Positive (↑Cardiac Contractility)	Negative (↓Cardiac Contractility)
Digoxin, digitalis	Beta blockers (β blockers)
Adrenaline & Noradrenaline	Acetylcholine
Dobutamine	Ether
Dopamine	Some bacterial toxins (e.g. diphtheria toxins)
Isoprenaline	K ⁺
Alkalosis	Acidosis
Ca ²⁺	Ca ²⁺ channel blockers
Caffeine	

22





Risk & Benefits of Inotropes

Benefits	Risk
Improves cardiac performance	↑ Heart rate, causing further deterioration of failing heart pump
Improves myocardial contractility	↑ Myocardial oxygen requirements
↑ Blood pressure	Potentially arrhythmogenic
	Can ↑ ischemia

25



Factors Affecting Myocardial Contractility ... (Cont.)



6. Mechanical factors:

- Cardiac ms obeys 'all or none law'
- Cardiac ms can perform both isometric & isotonic types of contractions
- Staircase phenomenon
- Cardiac ms can't be tetanized
- Starling's law of the heart

26



Mechanical Factors ... (Cont.)

a. Cardiac ms obeys 'all or none law'

- ❑ Action potential fails to occur if the stimulus is subthreshold in magnitude
- ❑ It occurs with constant amplitude & form regardless of the strength of the stimulus if the stimulus is at or above the threshold
 - This is made possible through rapid spread of excitation from cell to cell (as cardiac ms behaves as a syncytium)

27



Mechanical Factors ... (Cont.)

b. Cardiac ms Can perform both isometric & isotonic types of contractions

- ❑ **Isometric 'same length' contraction:**
Occurs when the muscle is stimulated after its ends has been fixed 'held'. It contracts & exerts an internal tension but cannot be shortened, so it does no external work
- ❑ **Isotonic 'same tension' contraction:**
Occurs when stimulated muscle is allowed to shorten

28



Mechanical Factors ... (Cont.)

c. Staircase Phenomenon

- ❑ After brief rest, on stimulation at regular frequency, the force of contraction \uparrow progressively to a maximum & then is maintained at a plateau
- ❑ Causes of staircase effect:
 - \uparrow accumulation of Ca^{2+}
 - \uparrow temperature
 - \downarrow viscosity





29



Mechanical Factors ... (Cont.)

d. Cardiac ms can't be tetanized

- ❑ Cardiac ms can't be stimulated while it's contracted
- ❑ Its excitability during contraction is zero, due to long ARP



KEY
▲ = Stimulus for action potential

30

Mechanical Factors ... (Cont.)

e. Starling's law of the heart

❑ Effect of load on contractility:

- **Pre-load:**
the load acting on heart before it starts contracting
- **After-load:**
the load acting on heart after it starts contracting ...
(resistance)


31

Mechanical Factors ... (Cont.)

e. Starling's law of the heart ... (Cont.)

- ❑ Starling's law of the heart states that the force of contraction is proportional to the initial length of the muscle within the physiological limits
 - ❑ Initial length depends on pre-load, i.e. end-diastolic volume (EDV) \approx 130 ml
 - ❑ Cardiac ms accommodates itself (up to certain limit) to the changes in venous return

32



Length - Tension Relationship

- ❑ As the pre-load \uparrow , the tension \uparrow
- ❑ **Passive tension** is given by diastolic intraventricular pressure
- ❑ **Active tension** is given by systolic intraventricular pressure
- ❑ **Total tension** is the sum of the parallel elastic tension, i.e. 'passive' & 'active' tension
- ❑ **Descending limb at high degree of stretch is due to disruption of myocardial fibers**

34

