

# Physiology Team 431



## Team leaders

**Mohammed Asiri**  
**Nour Al-Khawaja**


**Yafa Al-shamlan**  
**Sara Al-anazy**  
**Lama Mokhlis**  
**Tmader alaofi**  
**Hayfa alabdulkareem**  
**Dalal fatani**  
**Jomanah alshammari**  
**Shehanah alomair**

**Abdullah Al-Towim**  
**Khalid Al Mohaimedi**  
**AbdulRahman Al-Bakr**  
**Fahad Al-showishi**  
**Abdullah Al-Turki**  
**Saad Al-Mdemig**  
**Ahmad Al-Zuhair**  
**Mohammed Al Numeir**  
**Majid Al-Oriny**  
**Abdullrahman Alshahrani**  
**Tariq Al-Otaibi**  
**Abdulmalik Almufarrih**  
**Ahmed Almarzuqi**  
**Nasser Al-moosa**



# CARDIOVASCULAR SYSTEM BLOCK

## CARDIAC ELECTRICAL ACTIVITY (PHYSIOLOGY)



**Dr. Mona Soliman, MBBS, MSc, PhD**  
Associate Professor  
Department of Physiology  
Chair of Cardiovascular Block  
College of Medicine  
King Saud University

# Cardiac Electrical Activity

Automaticity of the heart: the heart is capable of □

**Generating** rhythmical electrical impulses . ۱

**Conduct** the impulses rapidly through the heart . ۲


**The atria contract about one sixth of a second ahead of ventricular contraction** □

Why?

To allow filling of the ventricles before they pump the blood into the circulation ■

# The Specialized Excitatory and Conductive System of the Heart

1. The sinoatrial (S-A node)
2. The internodal pathway
3. The atrioventricular (A-V node)
4. The atrioventricular bundle (Bundle of His)
5. Purkinje fibers



# Conduction of Impulses

Notice that the sino-atrial node is the:

- 1-Pacemaker of the heart.
- 2-Generator of the impulse.

## Sinoatrial node (S-A node):

Located in the superior lateral wall of the right atrium near the opening of the superior vena cava

Why?

### Pacemaker of the heart □

Its rate of rhythmic discharge is greater than any other part in the heart □

Highest frequency □

Is capable of originating action potentials □



# Conduction of Impulses


## Atrioventricular (A-V) node:

Located in the posterior wall of the right atrium □

## Delay in the conduction of impulses (0.1 sec) □

Why?

Allows time for the atria to empty the blood into the ventricles before ventricular contraction begin ■



# Conduction of Impulses

## The Purkinje System

- ▣ Purkinje fibers are very large fibers
- ▣ Transmit action potentials at a very high velocity (0.1-4.0 m/sec)



Why?

very high permeability of gap junctions ■  
→ ions are transmitted easily from one cell to the next ■  
→ enhance the velocity of transmission ■

Ventricular muscle contract at almost the same time ▣

# Conduction of Impulses

## The Purkinje System


Penetrate atrioventricular fibrous tissue

→ divides into right and left bundle branches

→ each branch spread toward the apex of the heart


→ divide into small branches

→ penetrate and become continuous with cardiac muscle fibers





# Spread of the cardiac impulse through the heart



Not Important

# Control of Excitation and Conduction in the Heart

The impulse normally arise s in the sinus node □

Why?

The Sinus Node is the Pacemaker of the Heart □

Its rate of rhythmical discharge is faster than □  
that of any other part of the heart

# Abnormal Pacemakers

Ectopic pacemaker: a pacemaker elsewhere than the sinus node □

The cause: □

Any other part of the heart develops a rhythmical discharge rate that is more rapid than that of the sinus node

Example: the A-V node or in the Purkinje fibers

# Abnormal Pacemakers

Blockage of transmission of the cardiac impulse  
from the sinus node to the other parts of the heart

Example: A-V block

→ cardiac impulses fails to pass from atria into the ventricles

the atria continues to beat at the normal rate of  
rhythm of the S-A node →


a new pacemaker develops in the Purkinje system →  
with a new rate

# Control of Heart Rhythmicity and Impulse Conduction by the Cardiac Nerves

The heart is supplied with both sympathetic and parasympathetic nerves

Parasympathetic nerves (vagi): mainly to the S-A and A-V nodes

Sympathetic nerves: all parts of the heart with strong supply to the ventricles



# Parasympathetic stimulation of the heart

↓ rate of rhythm of the S-A node    □

↓ transmission of impulses to the A-V node    □

Strong stimulation of the vagi:    □

Stop completely the rhythmical excitation by the S-A node    ■

Block completely transmission of cardiac impulses from the atria to the ventricle    ■

Some point in the Purkinje fibers develops a rhythm of its own    ■

## “Ventricular Escape”

Sympathetic Stimulation Of  
The Heart..

↑ rate of rhythm of the S-A node

↑ transmission of impulses to the A-V node

↑ force of contraction



# Questions:



1) which statement is correct according to S-A node? 

a) It has the highest rate if the rhythmic discharge .. 

b) Delay in the conduction of impulse.. 


c) Transmit AP at very high velocity.. 



Sympathetic stimulation of the heart leads to: 

a) decrease rate of rhythm of the S-A node.. 

b) increase force of contraction.. 

c) decrease transmission of impulses to the A-v node.. 

Good Luck..

PS: MALE'S NOTES INCLUDED IN THE WORD FILE