

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

Glucose Metabolism: Glycolysis

By

Reem M. Sallam, M.D.; Ph.D.

**Assistant Prof., Clinical Chemistry Unit, Pathology Dept.
College of Medicine, KSU
sallam@ksu.edu.sa**

Glycolysis: Revision

- **Major oxidative pathway of glucose**
- **The main reactions of glycolytic pathway**
- **The rate-limiting enzymes/Regulation**
- **ATP production (aerobic/anaerobic)**
- **Pyruvate kinase deficiency hemolytic anemia**

Substrate-level phosphorylation Vs. Oxidative phosphorylation

- **Phosphorylation** is the metabolic reaction of introducing a phosphate group into an organic molecule.
- **Oxidative phosphorylation:** The formation of high-energy phosphate bonds by phosphorylation of ADP to ATP coupled to the transfer of electrons from reduced coenzymes to molecular oxygen via the electron transport chain (ETC); it occurs in the mitochondria.
- **Substrate-level phosphorylation:** The formation of high-energy phosphate bonds by phosphorylation of ADP to ATP (or GDP to GTP) coupled to cleavage of a high-energy metabolic intermediate (substrate). It may occur in cytosol or mitochondria

Summary: Regulation of Glycolysis

Regulatory Enzymes (Irreversible reactions):

Glucokinase/hexokinase

PFK-1

Pyruvate kinase

Regulatory Mechanisms:

Rapid, short-term:

Allosteric

Covalent modifications

Slow, long-term:


Induction/repression

Apply the above mechanisms for each enzyme where applicable

Long-Term Regulation of Glycolysis

Insulin: Induction

Glucagon: Repression




Pyruvate Kinase Deficiency


Hemolytic Anemia

PK Mutation may lead to:

1. Altered Enz. kinetics
2. Decreased Enz. stability
3. Altered response to activator



Aerobic Glycolysis: Total Vs Net ATP Production



Aerobic Glycolysis: ATP Production

ATP Consumed:

2 ATP

ATP Produced:

Substrate-level 2 X 2 = 4 ATP


Oxidative-level 2 X 3 = 6 ATP

Total 10 ATP

Net:

10 - 2 = 8 ATP

Aerobic Vs Anaerobic Glycolysis




Anaerobic Glycolysis

NADH produced **cannot be** used by ETC for ATP production


(No O₂ and/or No mitochondria)

Less ATP production, as compared to aerobic glycolysis


Lactate is an obligatory end product, **Why?**




Lactate Dehydrogenase



Anaerobic Glycolysis in RBCs: 2,3-BPG Shunt



Anaerobic Glycolysis in RBCs: 2,3-BPG Shunt



Glycolysis in RBCs: ATP Production

ATP Consumed:

2 ATP

ATP Produced:

Substrate-level OR $2 \times 2 = 4$ ATP

$1 \times 2 = 2$ ATP

~~Oxidative-level $2 \times 3 = 6$ ATP~~

Total 4 OR 2 ATP

Net: OR $4 - 2 = 2$ ATP

$2 - 2 = 0$ ATP

Glycolysis in RBCs: Summary

End product:

Lactate

No net production or consumption of NADH

Energy yield:

If no 2,3-BPG is formed: 2 ATP

If 2,3-BPG shunt occurs: 0 ATP

PK Deficiency hemolytic anemia depends on:

Degree of PK Deficiency

Compensation by 2,3-BPG

Take Home Message

- **Glycolysis is the major oxidative pathway for glucose**
- **Glycolysis is employed by all tissues**
- **Glycolysis is a tightly-regulated pathway**
- **PFK-1 is the rate-limiting regulatory enzyme**

Take Home Message

- Glycolysis is mainly a catabolic pathway for ATP production, **But** it has some anabolic features (amphibolic)
- Pyruvate kinase deficiency in RBCs results in hemolytic anemia

Take Home Message

- **Net energy produced in:**
 - **Aerobic glycolysis:** 8 ATP
 - **Anaerobic glycolysis:** 2 ATP
- **Net energy produced in glycolysis in RBCs:**
 - **Without 2,3 BPG synthesis:** 2 ATP
 - **With 2,3 BPG synthesis:** 0 ATP