

## FERTILIZATION & IMPLANTATION AND TWINNING




By Prof. Saeed Abuel Makarem & Dr. Sanaa Alshaarawi

# **OBJECTIVES**

- By the end of the lecture, you should be able to:
- Identify <u>fertilization</u> and its site.
- List the **phases** of fertilization.
- Describe the **results** of fertilization.
- Describe the **formation** of **blastocyst**.
- Identify <u>implantation</u> and <u>its site</u>.
- Describe the mechanism of implantation.
- Describe the formation of primary chorionic villi.
- List the sites of ectopic pregnancy.



# FERTILIZATION




#### Definition:

It is the process
during which a
male gamete
(sperm) <u>unites</u>
with a female
gamete (oocyte)
to form a single
cell (ZYGOTE).


## **Fertilization**

- It is a complex process.
- It begins with a contact between sperm & ovum.
- Ends up with intermingling of the maternal and paternal chromosomes.



# Site of Fertilization


- Usually in the ampulla of uterine tube.
- Ampulla is the longest and widest part of the tube.
- Fertilization may occur in any other part of tube.
- <u>Never occurs in the</u> <u>uterine cavity.</u>
- Chemical signal from oocyte attracts the sperms.




#### **Phases of Fertilization**

- 1&2- Passage of the sperm through the cells of the <u>corona radiata</u> by <u>the effect of</u>:
  - a) <u>Hyaluronidase enzyme</u> secreted from the sperms.
  - b) By movement of its tail.
- 3- Penetration of the zona pellucida by <u>acrosine</u> (a substance secreted from <u>acrosomal cap</u>).
- 4- Fusion of the plasma membranes of the oocyte and the sperm.
- 5- Completion of the second meiotic division of the oocyte & formation of the female pronucleus.

6- Formation of the male pronucleus.






- **Zygote** is genetically unique.
- Half of its chromosomes comes from the <u>father</u> and the <u>other half</u> comes from the <u>mother</u>.
- New combination is formed which is <u>different</u> from either of the parents.
- This mechanism forms <u>biparental inheritance</u> and <u>leads to</u> <u>variation of the</u> <u>human species</u>.

### Sex of the Embryo


- <u>Embryo's chromosomal</u> <u>sex</u> is determined at the time of fertilization.
- <u>Sex</u> is determined by the type of sperm (X or Y) that fertilizes the oocyte.
- So, it is the <u>father</u> whose gamete <u>decides the sex.</u>



Zonal reaction : it is a change in properties of zona pellucida that makes it impermeable to other sperms.


# Results of Fertilization

- 1. <u>Stimulates</u> the penetrated <u>oocyte</u> to complete its 2<sup>nd</sup> meiotic division.
- 2. Restores the normal diploid number of chromosomes.
- 3. Determines the sex of the embryo.
- 4. Initiates cleavage (cell division) of the zygote.




## **Cleavage of Zygote**

- It is the repeated mitotic divisions of the zygote.
- Normally <u>occurs</u> in the uterine tube.
- Rapid increase in the number of the cells.
- These smaller embryonic cells are now called, <u>Blastomeres</u>.




- It begins about <u>30</u> <u>hours</u> after fertilization.
- Zygote divides into <u>2</u>, then <u>4</u>, then <u>8</u>, then <u>16</u> cells.
- Zygote lies within the thick zona pellucida during cleavage.
- Zygote migrates in the uterine tube during cleavage from lateral to medial.
- Under the microscope, the zona pellucida is a translucent membrane

## **Cleavage of Zygote**



# Morula

- When there are 16-32 blastomeres the developing human is called MORULA.
- The Morula reaches the uterine cavity at this stage.
- Spherical <u>Morula</u> is <u>formed</u> about 3 days <u>after</u> <u>fertilization.</u>
- It resembles mulberry or blackberry.




# BLASTOCYST

A cavity appears within the morula dividing its cells into <u>2 groups</u>:

- 1. Outer cell layer called trophoblast.
- Inner cell layer (mass) called Embryoblast attached to one of the poles of the blastocyst.
   The cavity is called


blastocystic cavity or blastocele.



#### • **Definition** :


- It is the process by which the Blastocyst penetrates the superficial (Compact) layer of the endometrium of the uterus.
- <u>Site:</u>
- The normal site of *implantation* is the posterior wall of the body of the uterus near the fundus.
- <u>Time:</u>
- It begins about the <u>6<sup>th</sup> day</u> <u>after fertilization.</u>
- It is completed by the 11th or 12th day.

# IMPLANTATION



#### <u>Mechanism:</u>

- The Morula reaches the uterine cavity by the 4<sup>th</sup> day after fertilization, & remains free for one or two days.
   Fluid passes from uterine cavity to the Morula.
- Now the Morula is called <u>Blastocyst</u>, its cavity is called blastocystic cavity, its cells divided into Embryoblast & Trophoblast.




By the 5<sup>th</sup> day the Zona pellucida degenerates.

Blastocyst begins implantation by the 6<sup>th</sup> day.

Trophoblast cells penetrate the epithelium of the endometrium.

Penetration results from **proteolytic enzymes** (eg.COX-2) produced by the **trophoblast**.




- Zona pellucida degenerates & disappears by the <u>5<sup>th</sup></u> day to allows the blastocyst to increase in size and penetrates the endometrium.
- The embryoblast projects into the blastocystic cavity, while the trophoblast forms the wall of the blastocyst.
- <u>By 6<sup>th</sup> day the blastocyst</u> adheres to the endometrium (A).
- By 7<sup>th</sup> day, <u>Trophoblast differentiated</u> into 2 layers: (B)


<u>Cytotrophblast</u>, inner layer, mitotically active.

Syncytiotrophoblast (outer

multinucleated mass, with *indistinct* cell boundary.

By 8<sup>th</sup> day the blastocyst is superficially embedded in the compact layer of the endometrium.





# Blood-filled Lacunae appear in the Syncytiotrophoblast which communicate forming a <u>lacunar network</u> by the 10<sup>th</sup> or 11<sup>th</sup> day.


 Syncytiotrophoblast erodes the endothelial

lining of the **maternal capillaries** which known as sinusoids.

**Now** blood of maternal capillaries reaches the lacunae so,

#### **Uteroplacental circulation**

is established by **11<sup>th</sup> or 12<sup>th</sup> day.** 



Endometrial cells undergo a process called apoptosis (programmed cell death) to facilitates invasion of endometrium by the Syncytiotrophoblast.


Syncytiotrophoblast engulf these degenerated cells for nutrition of the embryo.

#### **Implantation**

can be **detected** by:

1- Ultrasonography.

2- hCG (human chorionic gonadotrophin) which is secreted by the Syncytiotrophoblast about the <u>end</u> of 2<sup>nd</sup> week.



# **Early Pregnancy Factor**


• Is an immunosuppressant protein.

<u>Secreted</u> by trophoblast cells.


- <u>Appears in maternal serum</u> within 24--48
   hrs., after fertilization.
- It is the basis for EPT (Early pregnancy test) in the first 10 days of development.

# Formation of The Primary Chorionic villi

By the 13<sup>th</sup> day Proliferation of **Cytotrophblast cells** produce extension inside the **Syncytiotrophoblast** to form the **primary** chorionic villi.




## **Ectopic Implantation (Pregnancy)**




3 - 10. Implantation sites of blastocysts. The usual site in the posterior wall of the uterus is indicated by an X. The approximate order of new of ectopic implantations is indicated alphabetically (*A*, most common, *H*, least common). *A* to *F*, Tubal pregnancies. *G*, Abdominal ancy. *H*, Ovarian pregnancy. Tubal pregnancies are the most common type of ectopic pregnancy. Although appropriately included with uterine ancy sites, a cervical pregnancy is often considered to be an ectopic pregnancy.

#### **Ectopic Pregnancy**

- It means implantation <u>outside the uterine</u> <u>cavity.</u>
- 95 to 97% of ectopic pregnancies occurs in the uterine tube.
- Most are in the ampulla & isthmus.
- Placenta previa:
- Implantation occurs in the lower uterine segment.



Ectopic Pregnancy: 1- Placenta Previa. 2- Tubal. 3- Ovarian. 4- Abdominal. 5- Pelvic. 6- Cervical.



# THANK YOU