

FOUNDATION BLOCK

Pathology Practical

Prepared by:

- *Prof. Ammar Al Rikabi*
- *Dr. Sayed Al Esawy*
- *Dr. Marie Mukhashin*
- *Dr. Shaesta Zaidi*



Head of Pathology Department: Dr. Hisham Al Khalidi

PACTICAL 1

CELL INJURY & INFLAMMATION - I

***NORMAL ANATOMY AND
HISTOLOGY OF ORGANS RELATED
TO THIS CHAPTER***

Anatomy of the Respiratory System




Normal Histology of the Lung (Bronchiole, alveolar duct and alveoli)



**Microscopic section of normal lung showing:
terminal bronchiole (T) , respiratory bronchiole (R), alveolar duct (AD),
alveolar sac (AS), and alveoli (A).**


Normal Histology of the Lungs - Bronchiole



This view shows a BRONCHIOLE (right) and Blood Vessel (left) in cross-section as well as numerous ALVEOLI in The bronchiole inner membrane is composed of pseudostratified columnar epithelium.


Portions of hyaline cartilage rings can also be seen outside of the bronchiole.

Anatomy of the Heart





- ***The heart serves as a mechanical pump to supply the entire body with blood, both providing nutrients and removing waste products.***
- ***The great vessels exit the base of the heart.***
- ***Blood flow: Body → venae cavae → right atrium → right ventricle → lungs → left atrium → left ventricle → Aorta → body***

Anatomy of the Heart – inside view




Histology of the Heart

- The heart consists of 3 layers
 - the **endocardium**,
 - the **myocardium**, and
 - the **epicardium**.
- The **epicardium** consists of arteries, veins, nerves, connective tissue, and variable amounts of fat.
- The **myocardium** contains **branching, striated muscle cells with centrally located nuclei**. They are connected by **intercalated disks** (arrowheads).



Anatomy of the Normal Liver

Surfaces and Bed of Liver Anterior View



Rt

Cut surface of a Normal Liver



Lt



The cut surface of a normal liver has a brown color. Near the hilum here, note the portal vein carrying blood to the liver, which branches at center right, with accompanying hepatic artery and bile ducts.

At the lower left is a branch of hepatic vein draining blood from the liver to the inferior vena cava


Histology of Normal Liver



Gall Bladder & Extrahepatic Bile Duct


Gallbladder and Extrahepatic Bile Ducts

Right and left hepatic ducts




Gallbladder and Extrahepatic Bile Ducts

Sectioned




Histology of Gall Bladder



The gallbladder is a distensible sac and, when not distended, its mucosa is thrown into many folds. The lumen of the gallbladder is lined with a high columnar epithelium. The connective tissue wall contains abundant elastic fibers and layers of smooth muscle which predominantly run obliquely

Histology of Extra Hepatic Bile Duct




Normal histologic characteristics of the extra-hepatic bile duct. Photomicrograph (H&E stain) shows the epithelium as a single layer of columnar cells (arrow) with an underlying dense connective tissue wall.

Normal Appendix - Gross



The appendix (arrow) is a coiled 8-12 cm tube attached to the caecum, usually located in the right lower abdominal quadrant.

Histology of the appendix




The mucosa of the appendix, like that of the colon, is characterized by straight crypts with no villi. It is also characterized by an abundance of lymphoid tissue, including numerous well-organized lymph nodules.

Normal Kidney - Gross




In cross section, this normal adult kidney demonstrates the lighter outer cortex and the darker medulla, with the renal pyramids into which the collecting ducts coalesce and drain into the calyces and central pelvis.

Anatomy of the Kidney




Renal Corpuscle – Normal Histology



Normal glomerulus by light microscopy. The glomerular capillary loops are thin and delicate. Endothelial and mesangial cells are normal in number. The surrounding tubules are normal

Histology of the SKIN






Normal Skin. Epidermis has 4 layers :


- 1. Stratum basale**
- 2. Stratum spinosum (cells gain more cytoplasm)**
- 3. Stratum granulosum**
- 4. Stratum corneum (anucleate layer)**

Anatomy of the Brain


Brain Surface Anatomy



Histology of the Brain





Normal white matter: 85% of the cells are oligodendrocytes. Note some tendency for the oligodendrocytes to line up in rows



Cerebral cortex: some degree of satellitosis (i.e., oligodendrocytes surrounding neurons) is a normal finding

Spinal Cord Nerve Branches and Histology



Spinal motor neuron, with its prominent nucleolus in the nucleus and the basophilic tigroid appearance of its cytoplasm, which is due to the presence of abundant ribosomes

CELL INJURY

Gross and Histopathology

1 - FATTY LIVER (STEATOSIS)

Normal Liver & Cut Section of Fatty Liver




Normal Liver : This is the external surface of a normal liver. The color is brown and the surface is smooth



Steatosis : This liver is slightly enlarged and has a pale yellow appearance, seen both on the capsule and cut surface


Steatosis – Fatty Liver



This is the histologic appearance of hepatic fatty change. The lipid accumulates in the hepatocytes as vacuoles. These vacuoles have a clear appearance with H&E staining.

The most common cause of fatty change in developed nations is alcoholism.

Steatosis – Fatty Liver



***Here are seen the lipid vacuoles within hepatocytes.
The lipid accumulates when lipoprotein transport is disrupted and/or when
fatty acids accumulate.***

Alcohol is the most common cause


2- COAGULATIVE NECROSIS

Coagulative Necrosis of the Kidney




A typical pattern with ischemia and infarction of the kidney. Here, there is a wedge-shaped pale area of coagulative necrosis (infarction) in the renal cortex of the kidney.

Coagulative Necrosis of the Kidney - LPF



Coagulative necrosis of glomeruli, tubules and interstitial tissue with loss of cell nuclei. The haemorrhagic zone at the periphery of the infarct shows dilated and congested blood vessels and cellular infiltrate by neutrophils, red blood cells and lymphocytes

Coagulative Necrosis of the Kidney - HPF




The majority of the tubules seen here are proximal convoluted tubules. The PAS stain colors the brush border of these structures a deep pink-lavender. A pale-staining collecting duct stands out in contrast to the abundant proximal tubules

Coagulative Necrosis of the Spleen




Two large infarctions (areas of coagulative necrosis) are seen in this sectioned spleen

Coagulative Necrosis of Infarcted Myocardium



Many nuclei have become pyknotic (shrunken and dark) and have then undergone karyorrhexis (fragmentation) and karyolysis (dissolution). The cytoplasm and cell borders are not recognizable.


Coagulative Necrosis of Infarcted Myocardium



**The nuclei of the myocardial fibers are being lost.
The cytoplasm is losing its structure, because no well-defined cross-striations
are seen.**


3- LIQUEFACTIVE NECROSIS

Liquefactive Necrosis of the Brain




Grossly, the cerebral infarction at the upper right here demonstrates liquefactive necrosis. Eventually, the removal of the dead tissue leaves behind a cavity.

Liquefactive Necrosis of the Brain




Liquefactive necrosis in brain leads to resolution with cystic spaces. The necrotic area is found in the upper right quadrant of the visual field.

Liquefactive Necrosis of the Brain




This cerebral infarction demonstrates the presence of many macrophages at the right which are cleaning up the lipid debris from the liquefactive necrosis.

Liquefactive Necrosis of the Brain



This is the microscopic appearance of a lacunar infarct. Note that it is a cystic space from the resolved liquefactive necrosis. There can be hemosiderin pigment from hemorrhage as well.

Liquefactive Necrosis - Liver Abscess



The liver shows a small abscess here filled with many neutrophils. This abscess is an example of localized liquefactive necrosis


4- CASEOUS NECROSIS

Caseous Necrosis of the Lung “TB. Lung”



Tuberculosis of the lung, with a large area of caseous necrosis containing yellow-white and cheesy debris


T.B. Granuloma with Central Caseous Necrosis



Multiple caseating granulomas with giant cells and caseous necrosis. Note preserved alveolar spaces at the margins of the field.

5 – FIBRINOID NECROSIS


Fibrinoid Necrosis of an Artery - HPF



Fibrinoid necrosis in an artery. The wall of the artery shows a circumferential bright pink area of necrosis with inflammation (neutrophils with dark nuclei).


6 – FAT NECROSIS

Fat Necrosis in the Mesentery




The areas of white chalky deposits represent foci of fat necrosis with calcium soap formation (saponification) at sites of lipid breakdown in the mesentery

Fat Necrosis in the Mesentery




Fat necrosis of the mesentery in a case of acute pancreatitis
Numerous round white fat necroses

Fat Necrosis – Histopathology



The necrotic fat cells have vague cellular outlines, have lost their peripheral nuclei, and their cytoplasm has become a pink amorphous mass of necrotic material

Fat Necrosis – Histopathology



Fat necrosis adjacent to pancreas is seen here. There are some remaining steatocytes at the left which are not necrotic.

The necrotic fat cells at the right have vague cellular outlines, have lost their peripheral nuclei, and their cytoplasm has become a pink amorphous mass of necrotic material

7 - DYSTROPHIC CALCIFICATION


(AORTIC VALVE – STOMACH - SKIN)

Dystrophic calcification of Aortic Valve




View looking down onto the unopened aortic valve in a heart with calcific aortic stenosis. It is markedly narrowed (stenosis). The semilunar cusps are thickened and fibrotic, and behind each cusp are irregular masses of piled-up dystrophic calcification

Dystrophic calcification of Aortic Valve



Aortic valve. Fibrosis with some lymphocytes and dystrophic calcification (A) hematoxylin and eosin; 1.25× objective magnification; and siderosis (B) Berlin blue 40× objective magnification

Dystrophic Calcification of Stomach




This is a dystrophic calcification in the wall of the stomach. At the far right is an artery with calcification in its wall. There are also irregular bluish-purple deposits of calcium in the submucosa

Dystrophic Calcification of the Skin




Multiple erythematous hard papules in linear configuration on the extensor aspect of the arm. Within the lesion there were several 2-5 mm white calcifications

Dystrophic Calcification of the Skin



Calcifying panniculitis with fibrosis of the subcutaneous connective tissue septae, adjacent inflammation containing plasmacytes and lymphocytes, and a deposit of calcification (arrow).

Dystrophic Calcification of the Skin



Irregular blue granular deposits of calcium in the dermis surrounded by fibrous tissue and foreign body giant cell reaction

8- ATROPHY OF THE ORGANS

(BRAIN – TESTIS)

Atrophy of the Brain




This is cerebral atrophy in a patient with Alzheimer disease. The gyri are narrowed and the intervening sulci are widened, particularly pronounced toward the frontal lobe region.

Right

Atrophy of the Testis

Left




The testis at the left has undergone atrophy and is much smaller than the normal testis at the right.

9 - LEFT VENTRICULAR HYPERTROPHY

Normal and Hypertrophied Left Ventricle



Left ventricular hypertrophy: The number of myocardial fibers does not increase but their size increased in response to an increased workload



Normal ventricles

Right

Left Ventricular Hypertrophy

Left



This cross section view of the heart shows the left ventricle in the right of the picture. The heart is from a severe hypertensive patient. The left ventricle is grossly thickened. The myocardial fibers have undergone hypertrophy.

10- PROSTATIC HYPERPLASIA

Prostatic Hyperplasia - Gross




The normal adult male prostate is about 3 to 4 cm in diameter. The number of prostatic glands, as well as the stroma, has increased in this enlarged prostate

Prostatic Hyperplasia



Nodular hyperplasia of glandular and fibromuscular stromal tissue. Each nodule shows large number of glands of variable sizes lined by tall columnar epithelium and some are cystically dilated.

Prostatic Hyperplasia




Here is one of the nodules of hyperplastic prostate, with many glands along with some intervening stroma.

The cells making up the glands are normal in appearance, but there are just too many of them. Eosinophilic hyaline corpora amylacea is present in some glands.

11 - SQUAMOUS METAPLASIA AND DYSPLASIA

Normal Uterine Cervix




Normal cervix with a smooth, glistening mucosal surface. There is a small rim of vaginal cuff from this hysterectomy specimen. The cervical os is small and round, typical for a nulliparous woman. The os will have a fish-mouth shape after one or more pregnancies

Right


Normal and Dysplastic Cervical Squamous Epithelium

Left



The normal cervical squamous epithelium at the right transforms to dysplastic changes on the left with underlying chronic inflammation

Endocervical Squamous Metaplasia




A section of endocervix shows the normal columnar epithelium at both margins and a focus of squamous metaplasia in the center.

Right

Laryngeal Squamous Metaplasia

Left



Metaplasia of laryngeal respiratory epithelium has occurred here in a smoker .The chronic irritation has led to an exchanging of one type of epithelium (the normal respiratory epithelium at the left) for another (the more resilient squamous epithelium at the right)


GOOD *LUCK*

PRACTICAL 2

CELL INJURY & INFLAMMATION - II


I - Acute Inflammation

Pathogenesis of Exudation




The diagram shown here illustrates the process of exudation, aided by endothelial cell contraction and vasodilation, which typically is most pronounced in venules. Collection of fluid in a space is a transudate. If this fluid is protein-rich or has many cells then it becomes an exudate.

Exudation in the Alveolar Space




Here is vasodilation with exudation that has led to an outpouring of fluid with fibrin into the alveolar spaces along with PMN's indicative of an acute bronchopneumonia of the lung,

Exudation of Fibrin in Acute Inflammation




Here is an example of the fibrin mesh in fluid with PMN's that has formed in the area of acute inflammation. It is this fluid collection that produces the "tumor" or swelling aspect of acute inflammation.

Inflammation with Necrosis - LPF



The vasculitis shown here demonstrates the destruction that can accompany the acute inflammatory process and the interplay with the coagulation mechanism. The arterial wall is undergoing necrosis, and there is thrombus formation in the lumen.


Inflammation with Necrosis - HPF



At higher magnification, vasculitis with arterial wall necrosis is seen. Note the fragmented remains of neutrophilic nuclei (karyorrhexis). Acute inflammation is a non-selective process that can lead to tissue destruction


1 - FIBRINOUS PERICARDITIS

Acute Fibrinous Pericarditis - Gross



Here, the pericardial cavity has been opened to reveal a fibrinous pericarditis with strands of stringy pale fibrin between visceral and parietal pericardium

Acute Fibrinous Pericarditis - Gross




Serous fluid at the bottom of the pericardial cavity (arrow) is visible. The epicardial surface appears roughened, compared to its normal glistening appearance; due to the strands of pink-tan fibrin that have formed

Acute Fibrinous Pericarditis - Microscopically


Right

Left




The fibrinous exudate is seen to consist of pink strands of fibrin gutting from the pericardial surface at the upper right .The exudate on the surface is shown enlarged in the inset. Note a considerable number of erythrocytes trapped in the mesh of fibrin threads.

Acute Fibrinous Pericarditis - LPF



The pericardium is distorted by thick irregular layer of pinkish fibrinous exudate with some red cells and inflammatory cells

Acute Fibrinous Pericarditis - HPF



The subpericardial layer is thickened by edema and shows dilated blood vessels, chronic inflammatory cells and areas of calcification.

2- ACUTE APPENDICITIS

Normal Appendix - Gross




This is the normal appearance of the appendix against the background of the caecum.

Acute Appendicitis - Gross




Seen here is acute appendicitis with yellow to tan exudate and hyperemia, including the periappendiceal fat superiorly, rather than a smooth, glistening pale tan serosal surface

Acute Appendicitis – Longitudinal section




A case of acute appendicitis : The organ is enlarged and sausage-like (botuliform). This longitudinal section shows the angry red inflamed mucosa with its irregular luminal surface. This appendix does not show late complications, like transmural necrosis, perforation, and abscess formation


Acute Appendicitis – LPF of the cut section



Acute Appendicitis – LPF




Acute Appendicitis – HPF



Scattered Neutrophils in the crypt epithelium

Acute Appendicitis – Histopathology



This slide shows the muscle layer of the appendix which is permeated with numerous polymorphonuclear leukocytes


3- ACUTE CHOLECYSTITIS

Acute Cholecystitis – Gross



Mucocele, stone obstructed the neck , distended , aspiration done and removed by lap chole


Acute Cholecystitis – Histopathology HPF



The neutrophils are seen infiltrating the mucosa and submucosa of the gallbladder in this patient with acute cholecystitis and right upper quadrant abdominal pain with tenderness on palpation

4- SKIN PILONIDAL SINUS

Foreign Body Reaction (Pilonidal Sinus)




A pilonidal sinus is a sinus tract which commonly contains hairs. It occurs under the skin between the buttocks (the natal cleft) a short distance above the anus. Usually runs vertical between the buttocks and rarely occurring outside the coccygeal region.

Foreign Body Reaction (Pilonidal Sinus)



Surgically excised pilonidal sinus tracts

Pilonidal Sinus – Histopathology LPF



The lumen of the sinus and wall contain large number of hair shafts with foreign body giant cells, lymphocytes , macrophages & neutrophils

II - CHRONIC INFLAMMATION


1- CHRONIC CHOLECYSTITIS WITH STONES

Chronic cholecystitis with Gall Stones




Gross appearance of gallbladder after sectioning longitudinally. Notice thickness of gall bladder wall, abundant polyhedric stones and small papillary tumor in the cystic duct.

Chronic cholecystitis - Histopathology



Irregular mucosal folds and foci of ulceration in mucosa. Wall is penetrated by mucosal glands which are present in muscle coat (Rokitansky- Aschoff sinuses). All layers show chronic inflammatory cells infiltration and fibrosis.

Chronic cholecystitis - Histopathology



The mucosa is atrophic, with a single layer of flattened epithelium. There is proteinaceous fluid adherent to the mucosal surface, with some bile stained orange-brown crystals toward the upper left in the lumen. The lamina propria shows fibrosis and contains a mononuclear cell infiltrate (small dark blue nuclei).

The muscle is hypertrophied compared to normal gallbladder.

2- BRAIN ABSCESS

Brain Abscess - CT




CT of a cerebral abscess. There is a liquefactive center with yellow pus surrounded by a thin wall. Abscesses usually result from hematogenous spread of bacterial infection, but may also occur from direct penetrating trauma or extension from adjacent infection in sinuses

Right

Brain Abscess - MRI


Left



This trichrome stain demonstrates the light blue connective tissue in the wall of an organizing cerebral abscess. Normal brain is at the left and the center of the abscess at the right.


3 - GRANULATION TISSUE

Granulation Tissue - LPF



Section of fragments of edematous, loose connective tissue shows many small newly formed capillaries lined by plump endothelial cells. Proliferation of fibroblasts is seen

Granulation Tissue - HPF



***Inflammatory cells including macrophages, lymphocytes, plasma cells and neutrophils in the oedematous stroma.
Pink homogenous collagen fibers may be identified.***


GOOD *LUCK*

PRACTICAL 3

THROMBO-EMBOLIC DISORDERS

1 - ORGANIZING THROMBUS

Organizing Thrombus




Organizing thrombus in a case of pulmonary embolism

Organizing Thrombus with Lines of Zahn




This is the microscopic appearance of a pulmonary thromboembolus in a large pulmonary artery. There are interdigitating areas of pale pink and red that form the "lines of Zahn" characteristic for a thrombus. These lines represent layers of red cells, platelets, and fibrin which are laid down in the vessel as the thrombus forms.

Lines of Zahn



Lines of Zahn, gross and microscopic, is evidence to prove a clot is Pre-mortem which is different from the clots appearing like current jelly or chicken fat which are said to be Post-mortem. These lines represent layers of red cells, platelets, and fibrin



Thromboembolus in Pulmonary Artery



Pulmonary thromboembolus in a small pulmonary artery. The interdigitating areas of pale pink and red within the organizing embolus form the “lines of Zahn” (arrow) characteristic of a thrombus. These lines represent layers of red cells, platelets, and fibrin that are laid down in the vessel as the thrombus forms

2- PULMONARY EMBOLUS WITH INFARCTION

Pulmonary Embolus with Infarction



This specimen shows an area of dead lung tissue ("infarction") due to blockage of one of the major arteries to the lung by an embolus ("blood clot") originating from the deep veins of the leg.


Pulmonary Embolus with Infarction



A large pulmonary thromboembolus is seen in the pulmonary artery of the left lung. Such thromboemboli typically originate in the leg veins or pelvic veins of persons who are immobilized


3- MYOCARDIAL INFARCTION

Myocardial Infarction




Complications that might occur : arrhythmias , ventricular aneurysm, rupture of myocardium, cardiac tamponade and others .

Myocardial Infarction




Cross section of the left and right ventricles shows a pale and irregular focal fibrosis in the left ventricular wall with increased thickness .

Myocardial Infarction




Cross section of the left and right ventricles shows a pale and irregular focal fibrosis in the left ventricular wall with increased thickness .

Myocardial Infarction




Transmurular myocardial infarct at 2 weeks

Myocardial Infarction



Acute myocardial infarct, histology. This 3-4 day old infarct shows necrosis of myocardial cells and is infiltrated with polymorphnuclear leukocytes.


Myocardial Infarction



- 1- Patchy coagulative necrosis of myocardial fibers. The dead muscle fibers are structureless and hyaline with loss of nuclei & striations.**
- 2- Chronic ischemic fibrous scar replacing dead myocardial fibers .**
- 3- The remaining myocardial fibers show enlarged nuclei due to ventricular hypertrophy .**


4- INFARCTION OF THE SMALL INTESTINE

Infarction of the Small Intestine




The dark red infarcted small intestine contrasts with the light pink viable bowel. The forceps extend through an internal hernia in which a loop of bowel and mesentery has been caught. This is one complication of adhesions from previous surgery. The trapped bowel has lost its blood supply

Infarction of the Small Intestine



Diffuse violaceous red appearance is characteristic of transmural hemorrhagic intestinal infarction

Infarction of the Small Intestine



Intestinal infarction typically begins in the villi, which are end vasculature without anastomoses. There is complete loss of the mucosal epithelium. Broad areas of hemorrhage with moderate inflammatory infiltrate is present

PRACTICAL 4

GRANULOMATOUS DISEASES

1 - TUBERCULOSIS OF THE LUNG

Pulmonary TB – Caseous Necrosis – Gross




The granulomas have areas of caseous necrosis. This pattern of multiple caseating granulomas primarily in the upper lobes is most characteristic of secondary T.B.

Pulmonary TB – Caseous Necrosis – Gross



Initial (primary) infection with T.B. producing a sub-pleural lesion called a Ghon's focus. The early Ghon's focus together with the lymph node lesion constitute the Ghon's complex..


Pulmonary TB - Ghon's Complex – Gross




The Ghon's complex is seen here at closer range. Primary tuberculosis is the pattern seen with initial infection with tuberculosis in children. Reactivation, or secondary tuberculosis, is more typically seen in adults.

Miliary TB of the Lungs

- ***Miliary TB can occur when TB lung lesions erode pulmonary veins or when extrapulmonary TB lesions erode systemic veins.***
- ***This results in hematogenous dissemination of tubercle bacilli producing myriads of 1-2 mm. lesions throughout the body in susceptible hosts.***
- ***Miliary spread limited to the lungs can occur following erosion of pulmonary arteries by TB lung lesions.***




Tuberculous Granulomas




Well-defined granulomas are seen here. They have rounded outlines. The one toward the center of the photograph contains several Langhan's giant cells. Granulomas are composed of transformed macrophages called epithelioid cells along with lymphocytes, occasional PMN's, plasma cells, and fibroblasts

Pulmonary TB - Granuloma with central early necrosis




The pyknotic nuclei of epithelioid cells in the center of the granuloma (apoptotic bodies) are a precursor of necrosis.

Tuberculous Granulomas



The edge of a granuloma is shown here at high magnification. At the upper is amorphous pink caseous material [1] composed of the necrotic elements of the granuloma as well as the infectious organisms. This area is ringed by the inflammatory component [2] with epithelioid cells, lymphocytes, and fibroblasts.

Acid Fast bacilli of Mycobacterium TB in the Lung




**A stain for *Acid Fast Bacilli* is done (*AFB* stain) to find the mycobacteria .
The mycobacteria stain as red rods, as seen here at high magnification.**

2- TUBERCULOUS LYMPHADENITIS

Tuberculous Lymphadenitis - Gross




Tuberculous Lymphadenitis – Cut Section




Section of a lymph node with connective tissue capsule and lymphoid tissue

Tuberculous Lymphadenitis



Many round and oval tubercles/ granulomas with or without central caseation that appears structureless, homogenous and pink in colour.


Tuberculous Lymphadenitis



The granulomas consists of epithelioid cells, few langhan's giant cells (large cell with multiple peripheral nuclei) and peripheral rim of lymphocytes


3- BILHARZIAL GRANULOMAS

Colonic Bilharziasis - HPF




Colon biopsy of bilharziasis. Fibrosing foreign body granuloma against the miracidium-containing ovum of *S. mansoni* is observed in the submucosal layer (H&E).

Bilharziasis of the Urinary Bladder



Schistosoma haematobium. Urinary Bladder biopsy showing bilharziasis eggs

S. japonicum in the Hepatic portal tract



S. japonicum eggs in hepatic portal tract


4- CUTANEOUS LEISHMANIASIS

Cutaneous Leishmaniasis




Leishmaniasis is caused by parasitic infection, mainly by parasites of the *Leishmania* genus which are carried by a blood-sucking insect known as the sandfly.

Cutaneous Leishmaniasis



Histological view shows marked cellular infiltration and parasites (Leishman bodies) within macrophages

Cutaneous Leishmaniasis



The blood film shows macrophages containing Leishmania amastigotes, each with a prominent kinetoplast (seen as a darkened spot next to the larger nucleus) and no flagella (in contrast with the promastigote form).

GOOD *LUCK*

PRACTICAL 5


NEOPLASIA



(BENIGN TUMORS)

1 - ADENOMATOUS POLYP OF RECTUM / COLON

Adenomatous polyp of the colon



This adenomatous polyp has a hemorrhagic surface (which is why they may first be detected with stool occult blood screening) and a long narrow stalk. The size of this polyp--above 2 cm--makes the possibility of malignancy more likely, but this polyp proved to be benign

Adenomatous polyp of the colon




This small adenomatous polyp (tubular adenoma) on a small stalk is seen microscopically to have more crowded, disorganized glands than the normal underlying colonic mucosa. Goblet cells are less numerous and the cells lining the glands of the polyp have hyperchromatic nuclei

Right

Adenomatous polyp of the colon


Left



A microscopic comparison of normal colonic mucosa on the right and that of an adenomatous polyp (tubular adenoma) on the left is seen here. The neoplastic glands are more irregular with darker (hyperchromatic) and more crowded nuclei


2- LIPOMA

Lipoma of the Neck




Benign, slow growing, subcutaneous skin growth. In this case, the lipoma is rather large and located in the neck region. On palpation, these are soft, non tender, and mobile if it is small size.

Lipoma – Cut section



Lipoma is a benign tumor composed of mature adipose tissue. Most of them are superficially located in the upper part of the body, although they can arise anywhere. Grossly, they appear bright yellow and lobulated

Lipoma with fat necrosis



This picture shows an area of fat necrosis within a lipoma. The masses are comprised primarily of mature adipocytes. Histiocytes present within these areas should not be mistaken for lipoblasts


3- INTRADERMAL NEVUS

Intradermal Nevus




***The lesion is small, symmetrical, and uniformly has different colors
(Pink – Tan – Brown etc)***

Intradermal Nevus - LPF



Nests and clusters of small round or spindle shaped nevus cells with few melanophages in the upper dermis.


Intradermal Nevus - HPF



The cells contain varying amount of brown melanin pigment. No junctional activity


4- UTERINE LEIOMYOMATA

Multiple Uterine Leiomyomata - Gross




**Smooth muscle tumors of the uterus are often multiple.
Seen here are submucosal, intramural, and subserosal leiomyomata
of the uterus.**

Multiple Uterine Leiomyomata - Gross




A well demarcated tumour mass in the muscle coat of uterus without a definite capsule.

Uterine Leiomyoma – LPF Microscopy



Normal myometrium is at the left, and the neoplasm is well-differentiated so that the leiomyoma at the right hardly appears different. Bundles of smooth muscle are interlacing in the tumor mass

Uterine Leiomyoma – HPF Microscopy



The muscle cells are spindle shaped with elongated nuclei and eosinophilic cytoplasm

5- CHONDROMA


Enchondroma of the fibula




The picture shows intramedullary bone expansion, chondromyxoid material, thin bone cortex.

Enchondroma of the Fibula - Microscopic

LPF




HPF



Lobules consist of mature cartilage cells irregularly distributed through pale blue homogenous matrix and are contained within the lacunar spaces singly, in pairs or in tetrads.

Few bony trabeculae are included in the tumour.

Enchondroma of the Fibula - HPF



Cartilage shows hypo to moderate cellularity and contains chondrocytes of variable sizes. Chondrocyte nuclei tend to be small, round and hyperchromatic. Scattered binucleated cells may be found. Irregular purple granules within the matrix represent calcifications.


6 - HEMANGIOMA

Hemangioma of the Skin




A tumour mass in the dermis which consists of large number of vascular spaces of varying shapes and sizes separated by connective tissue stroma.

Capillary Hemangioma of the skin – LPF




Histopathology of cutaneous capillary hemangioma. Skin biopsy. H/E stain

Capillary hemangioma of the skin - LPF



Vascular spaces are lined by the flattened endothelial cells and some contain blood. Delicate connective tissue stroma separated the capillary vascular spaces


Cavernous Hemangioma of Skin – HPF



Large cavernous hemangioma, usually on an extremity, complicated by thrombocytopenic purpura. Blue rubber bleb nevus syndrome: cavernous hemangiomas of the skin and gastrointestinal tract.

7- TERATOMA (DERMOID CYST) OF THE OVARY

Ovary: Mature Cystic Teratoma




Opened mature cystic teratoma (dermoid cyst) shows hair (bottom) and a mixture of tissues .

Ovary: Mature Cystic Teratoma




This 4.0 cm dermoid cyst is filled with greasy material (keratin and sebaceous secretions) and shows tufts of hair. The rounded solid area at the bottom is called Rokitansky's protruberance. Microscopically, it also showed foci of neural tissue.

Ovary: Mature Cystic Teratoma




Ovarian teratoma showing neuroepithelial tubules and rosettes (immature component) adjacent to a hair follicle (mature component). They consist of epidermis, hair follicles, sweat and sebaceous glands and neuroectodermal derivatives

Ovary: Mature Cystic Teratoma



This image shows skin and mucinous glands in a mature solid teratoma of the ovary

Ovary: Mature Cystic Teratoma



Stratified Squamous epithelium with underlying sweat glands, sebaceous glands, hair follicles, columnar ciliated epithelium, mucous and serous glands and structures from other germ layers such as bone and cartilage, lymphoid tissue, smooth muscle and brain tissue containing neurons and glial cells

PRACTICAL 6

NEOPLASIA



(MALIGNANT TUMORS)


1 - SQUAMOUS CELL CARCINOMA OF THE SKIN

Squamous Cell Carcinoma - Gross




Squamous cell carcinoma (SCC) is the second most common cancer of the skin. A sore that does not heal or any change in an existing mole, wart, or skin lesion can point to SCC. There may be an ulcer or reddish skin plaque that grows very slowly, may bleed occasionally (especially if located on the lip), may have an ulcerated center with raised, hard edges, may have a pearly quality with tiny blood vessels, is commonly present on sun-exposed areas (back of hands, lip (usually a small ulcer which will not heal and bleeds sporadically, ears and the scalp.

Squamous Cell Carcinoma - Histopathology




The normal squamous epithelium at the right merges into the squamous cell carcinoma at the left, which is infiltrating downward. The neoplastic squamous cells are still similar to the normal squamous cells, but are less orderly

Squamous Cell Carcinoma - Histopathology



Here is a moderately differentiated squamous cell carcinoma in which some, but not all, of the neoplastic cells in nests have pink cytoplasmic keratin

Squamous Cell Carcinoma - HPF




At high magnification, this squamous cell carcinoma demonstrates enough differentiation to tell that the cells are of squamous origin. The cells are pink and polygonal in shape with intercellular bridges.

The neoplastic cells show pleomorphism, with hyperchromatic nuclei.


A mitotic figure is present near the center

Squamous Cell Carcinoma - HPF



A mitotic figure is seen here in the center, surrounded by cells of a poorly differentiated squamous cell carcinoma, with pleomorphic cells that have minimal pink keratinization in their cytoplasm. In general, mitoses are more likely to be seen in malignant neoplasms

Squamous Cell Carcinoma - HPF



The dermis is infiltrated by masses of well differentiated neoplastic squamous cells separated by fibrous tissue stroma with chronic inflammatory cells. Tumour cells show pleomorphism, hyperchromatism and many mitotic figures .

Pinkish laminated keratin pearls (epithelial cell nests) are present in the center of some cell masses

2- ADENOCARCINOMA OF THE LARGE INTESTINE

Adenocarcinoma of the Colon




This cancer is more exophytic in its growth pattern. Thus, one of the complications of a carcinoma is obstruction (usually partial).

Adenocarcinoma of the Colon




This is an adenocarcinoma arising in a villous adenoma. The surface of the neoplasm is polypoid and reddish pink. Hemorrhage from the surface of the tumor creates a guaiac positive stool. This neoplasm was located in the sigmoid colon

Adenocarcinoma of the Colon




***A moderately differentiated colonic adenocarcinoma .
Tumour consists of crowded irregular malignant acini separated by
thin fibrovascular stroma.***

Adenocarcinoma of the Colon - LPF




Here is an adenocarcinoma in which the glands are much larger and filled with necrotic debris.

Adenocarcinoma of the Colon - LPF



The acini are lined by one or several layers of neoplastic cells with papillary projection showing pleomorphism, hyperchromatism and few mitoses.


Adenocarcinoma of the Colon - HPF



At high magnification, the neoplastic glands of adenocarcinoma have crowded nuclei with hyperchromatism and pleomorphism. No normal goblet cells are seen


3- LEIOMYOSARCOMA

Leiomyosarcoma




Cut surface of this leiomyosarcoma showing ill defined pale and soft large fleshy mass with hemorrhage and necrosis.

Leiomyosarcoma of Small Intestine




This is a leiomyosarcoma of the small bowel. As with sarcomas in general, this one is big and bad. Sarcomas are uncommon at this site, but must be distinguished from other types of neoplasms.

Leiomyosarcoma – HPF Microscopy



**Marked atypia and cellularity with multiple mitoses present.
Classic features of leiomyosarcoma including cigar shaped nuclei and arrangement of
cells in fascicles are seen.**

Leiomyosarcoma of the Uterus - HPF



Sarcomas, including leiomyosarcomas, often have very large bizarre giant cells along with the spindle cells. A couple of mitotic figures appear at the right and lower right

THE END