

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ


# **Collagen and Creatine: Protein and nonprotein nitrogenous compounds**

**Dr. Sumbul Fatma**  
**Department of Pathology**

# Objectives

1. To study the importance of creatine in muscle as a storage form of energy
2. To understand the biosynthesis of creatine
3. To study the process of creatine degradation and formation of creatinine as an end product
4. To understand the clinical importance of creatinine as a sensitive indicator of kidney function
5. To study the structure, function, types, and biosynthesis of collagen
6. To understand the different diseases associated with collagen

# Creatine Metabolism



**End product**

**Energy Source**

**Figure 21.16**  
Synthesis of creatine.

# Creatine Biosynthesis

Three amino acids are required:

**Glycine**

**Arginine**


**Methionine (as S-adenosylmethionine)**

Site of biosynthesis:

**Step 1: Kidneys**


**Step 2: Liver**

# Creatine Biosynthesis



# Distribution of body creatine

- From liver, transported to other tissues
- **98% are present in skeletal and heart muscles**
- In Muscle, gets converted to the high energy source **creatine phosphate (phosphocreatine)**



# **Creatine Phosphate**

- **Is a high-energy phosphate compound**
- **Acts as a storage form of energy in the muscle**
- **Provides a small but, ready source of energy during first few minutes of intense muscular contraction**


**The amount of creatine phosphate in the body is proportional to the muscle mass**



# Creatine Degradation

1. Creatine and creatine phosphate spontaneously form **creatinine** as an **end product**
2. Creatinine is excreted in the urine
3. Serum creatinine is a sensitive indicator of kidney disease (Kidney function test)
4. Serum creatinine **increases** with the impairment of kidney function

# Creatine Degradation




# Urinary Creatinine

- A typical male excretes about 15mmol of creatinine per day
- A decrease in muscle mass due to muscular dystrophy or paralysis leads to decreased level of creatinine in urine
- The amount of creatinine in urine is used as an indicator for the proper collection of 24 hours urine sample

# Creatine Kinase (CK)

- CK is responsible for the generation of energy in contractile muscular tissues
- CK levels are changed in disorders of cardiac and skeletal muscle



**Collagen: The most common  
animal protein**

# Collagen: Overview

- Most abundant protein in the human body
- Collagens are highly stable molecules, having half-lives as long as several years
- A fibrous protein that serves structural functions
- Is a part of connective tissues: bone, teeth, cartilage, tendon, skin, blood vessels
- Has a long rigid structure

# Collagen structure: The $\alpha$ -chain

- Collagen  $\alpha$ -chain ( $\sim 1,000$  amino acids long), is rich in proline and glycine
- The glycine residues are part of a repeating sequence,  $-\text{Gly}-\text{X}-\text{Y}-$ , where X is frequently proline and Y is often hydroxyproline or hydroxylysine
- Collage consists of three  $\alpha$ -chains wound around one another in rope like triple helix

*Compare between the 2 examples of secondary structure of proteins: the collagen helix & the  $\alpha$ -helix*

- The three polypeptide chains are held together by hydrogen bonds

# Structure of Collagen




- Rich in proline and glycine amino acids
- Proline prevents collagen chains to form  $\alpha$ -helix because:
  - It does not have back bone amino group (it is a ring structure with secondary amino group)
  - Therefore hydrogen bonding within the helix is not possible





# Non-standard amino acids in collagen


- Proline and lysine is converted to hydroxyproline and hydroxylysine by **hydroxylase** enzymes during post-translational modification
- The enzyme requires vitamin C for its function



Hydroxyproline

Proline

Glycine




# Types of collagen molecules


- Type and organization of collagen depends on its function
- Variations in the amino acid sequence of  $\alpha$ -chains result in different properties e.g.
  - Type I-  $(\alpha 1)_2 \alpha 2$
  - Type II-  $(\alpha 1)_3$

TYPE	TISSUE DISTRIBUTION
<b>Fibril-forming</b>	
I	Skin, bone, tendon, blood vessels, cornea
II	Cartilage, intervertebral disk, vitreous body
III	Blood vessels, fetal skin
<b>Network-forming</b>	
IV	Basement membrane
VII	Beneath stratified squamous epithelia
<b>Fibril-associated</b>	
IX	Cartilage
XII	Tendon, ligaments, some other tissues

# Biosynthesis of Collagen


- Synthesized in fibroblasts, osteoblasts and chondroblasts (pre-pro- then pro- and finally mature -collagen)
- Polypeptide precursors are enzymatically modified and form triple helix which is secreted into the extracellular matrix as **procollagen**
- **Glycosylation** of some hydroxylysine residues with glucose or galactose
- Procollagen molecules are cleaved by N- and C-procollagen peptidases releasing triple helical **tropocollagen** molecules
- Tropocollagen molecules spontaneously associate to form **collagen fibrils**





# Cross-linking of Collagen fibrils

- **Lysyl oxidase** oxidatively deaminates some of the lysine and hydroxylysine residues in collagen
- The produced reactive aldehydes- allysine and hydroxyallysine condense with lysine or hydroxylysine residues in neighbouring collagen molecules to form covalent cross-links
- This produces **mature collagen fibres**



# Collagen Diseases

- **Acquired disease:**

**Scurvy:** due to vitamin C deficiency

- **Genetic, inherited diseases:**

**Ehlers-Danlos syndromes (EDS)**


**Osteogenesis imperfecta (OI)**



# Collagen Diseases

**Scurvy**: due to vitamin C deficiency

Scorbutic gums in vitamin C deficiency. Gums are swollen, ulcerated, and bleeding due to vitamin C-induced defects in oral epithelial basement membranes and periodontal collagen fiber synthesis.




# Collagen Diseases

## Ehlers-Danlos syndrome:


can be caused by

- deficiency of lysyl hydroxylase or N-procollagen peptidase,
- Mutations in the amino acid sequences of collagen I, III and V.
- Characterized by hyperextensibility of joints and skin





# Collagen diseases



## Osteogenesis imperfecta (brittle bone disease):

Characterized by bones that fracture easily, with minor or no trauma.

Mutations **replace glycine with amino acids having bulky side chains** preventing the formation of triple helical conformation

- **Type I-** **most common**, characterized by mild bone fragility, hearing loss and blue sclerae
- **Type II-** most severe form and typically lethal in the perinatal period. Fractures are seen in utero.
- **Type III-** severe form, characterized by multiple fractures at birth, short stature, spinal curvature leading to a humped back (kyphotic) appearance and blue sclerae.

# References

- Lippincott, pages 43-49 and 287-288
- Bishop 6<sup>th</sup> edition, page 223-227