

Autoimmune Diseases

Dr. Hend Alotaibi

**Assistant professor & Consultant
College of Medicine, King Saud University
Dermatology Department /KKUH**

Email: halotaibi1@ksu.edu.sa

Objectives

- To know that the inflammatory processes in auto immune diseases are mediated by hypersensitivity reactions (type II, III and IV)
- To know that autoimmune diseases can be either organ specific or may be generalized involving many organs or tissues
- To understand that the manifestations of autoimmune diseases depend upon the organ and the degree of damage inflicted on the target tissues

**Disease processes and tissue damage are
due to Type II Type III and Type IV
hypersensitivity reactions**

SOME AUTOIMMUNE DISEASES IN HUMANS

Disease	Self-antigen	Immune response
Organ-specific autoimmune diseases		
Addison's disease	Adrenal cells	Auto-antibodies
Autoimmune hemolytic anemia	RBC membrane proteins	Auto-antibodies
Goodpasture's syndrome	Renal and lung basement membranes	Auto-antibodies
Graves' disease	Thyroid-stimulating hormone receptor	Auto-antibody (stimulating)
Hashimoto's thyroiditis	Thyroid proteins and cells	T _{DTH} cells, auto-antibodies
Idiopathic thrombocytopenia purpura	Platelet membrane proteins	Auto-antibodies
Insulin-dependent diabetes mellitus	Pancreatic beta cells	T _{DTH} cells, auto-antibodies
Myasthenia gravis	Acetylcholine receptors	Auto-antibody (blocking)
Myocardial infarction	Heart	Auto-antibodies
Pernicious anemia	Gastric parietal cells; intrinsic factor	Auto-antibody
Poststreptococcal glomerulonephritis	Kidney	Antigen-antibody complexes
Spontaneous infertility	Sperm	Auto-antibodies
Systemic autoimmune disease		
Ankylosing spondylitis	Vertebrae	Immune complexes
Multiple sclerosis	Brain or white matter	T _{DTH} and T _C cells, auto-antibodies
Rheumatoid arthritis	Connective tissue, IgG	Auto-antibodies, immune complexes
Scleroderma	Nuclei, heart, lungs, gastrointestinal tract, kidney	Auto-antibodies
Sjogren's syndrome	Salivary gland, liver, kidney, thyroid	Auto-antibodies
Systemic lupus erythematosus (SLE)	DNA, nuclear protein, RBC and platelet membranes	Auto-antibodies, immune complexes

spectrum of autoimmune disease

organ specific



non-organ specific

Hashimoto's thyroiditis
Primary myxoedema
Thyrotoxicosis
Pernicious anaemia
Autoimmune atrophic gastritis
Addison's disease
Premature menopause (few cases)
Insulin-dependent diabetes mellitus
Goodpasture's syndrome
Myasthenia gravis
Male infertility (few cases)
Pemphigus vulgaris
Pemphigoid
Sympathetic ophthalmia
Phacogenic uveitis
Multiple sclerosis (?)
Autoimmune haemolytic anaemia
Idiopathic thrombocytopenic purpura
Idiopathic leucopenia
Primary biliary cirrhosis
Active chronic hepatitis (HBs Ag negative)
Cryptogenic cirrhosis (some cases)
Ulcerative colitis
Sjögren's syndrome
Rheumatoid arthritis
Dermatomyositis
Scleroderma
Mixed connective tissue disease
Discoid lupus erythematosus
Systemic lupus erythematosus (SLE)

Examples of Autoimmune Diseases Affecting Different Systems:

Nervous System:

Multiple sclerosis
Myasthenia gravis

Autoimmune neuropathies such as:

- Guillain-Barré Syndrome (GBS)

Autoimmune uveitis

Blood:

Autoimmune hemolytic anemia
Pernicious anemia
Autoimmune thrombocytopenia

Blood Vessels:

Temporal arteritis
Anti-phospholipid syndrome
Vasculitides such as
Wegener's granulomatosis
Behcet's disease

Skin:

Psoriasis
Dermatitis herpetiformis
Pemphigus vulgaris
Vitiligo

Gastrointestinal System:

Crohn's Disease
Ulcerative colitis
Primary biliary cirrhosis
Autoimmune hepatitis

Endocrine Glands:

Type 1 or immune-mediated diabetes mellitus
Grave's Disease
Hashimoto's thyroiditis
Autoimmune oophoritis and orchitis
Autoimmune disease of the adrenal gland

Multiple Organs, Musculoskeletal System

Rheumatoid arthritis
Systemic lupus erythematosus
Scleroderma
Polymyositis, dermatomyositis
Ankylosing spondylitis
Sjogren's syndrome

Organ Specific Autoimmune Diseases

Mediated by **stimulating** or **blocking**
auto-antibodies

- 1) Graves' disease (**Stimulating antibodies**)
- 2) Myasthenia gravis (**Blocking Antibodies**)

1. Graves' Disease (Thyrotoxicosis)

- Production of thyroid hormones is regulated by **thyroid-stimulating hormones (TSH)**

- The binding of TSH to a receptor on thyroid cells stimulates the synthesis of two **thyroid hormones: thyroxine and triiodothyronine**






Fig. 1A



Fig. 1B


- A person with Graves' Disease makes **auto-antibodies to the receptor for TSH**.
- Binding of these auto-antibodies to the receptor **mimics** the normal action of TSH leading to over-stimulation of the thyroid gland

2. Myasthenia Gravis


- Clinically characterized by weakness and fatigability on sustained effort
- Antibodies directed against acetylcholine receptor (AChR)
- IgG Ab interact with the postsynaptic AChR at the nicotinic neuromuscular junction (NMJ)
- There is reduction in the number of functional AChR receptors by increasing complement mediated degradation of receptors

Myasthenia gravis

Normal events at the neuromuscular junction



Myasthenia gravis



Motor end-plates of muscles

Fig 13.10 © 2001 Garland Science

Systemic Autoimmune Immune diseases

I. Systemic lupus erythematosus (SLE)

- o Systemic lupus erythematosus is the most common autoimmune disorder
- o The characteristic “**butterfly rash**” is made worse by exposure to sunlight
- o Lupus is a potentially fatal autoimmune disease



Figure 13.11 The Immune System, 3ed. (© Garland Science 2009)

Genetic + Environment factors

Pathogenic Auto antibodies
-DNA/protein, RNA/protein complexes

Immune complexes

Complement activation

Chemotaxins

leukocytes
mononuclear cells

Inflammatory factors
IL-4, IL-6
IL-10

Destruction of cells

Symptom complex

Constitutional

Fatigue:

Myalgia

Fever:

Weight change:

Arthritis:

- migratory and asymmetrical. Only a few joints are usually affected, especially the hands

- Joint deformities including ulnar deviation, MCP subluxation, and **swan-neck deformities** caused by tendon laxity, rather than bony destruction.

Dermatological: CNS:

- malar rash
- discoid lesions
- hair loss
- oral ulcers
- Raynaud's
- Nailfold erythema/crises
- livedo on hands/legs
- Bullous rash on legs
- dermatitis on fingers

- cognitive defects, anxiety, depression, psychosis, seizures, and/or neuropathies, cerebral punctate vasculitis

Cardiovascular

- Pericarditis
- **Verrucous endocarditis** => emboli
- CAD from steroids

Pulmonary:

- Dyspnea and restrictive LFTs
- Pleurisy, pleural effusion, pneumonitis, interstitial lung disease, and pulmonary hypertension

Renal:


- glomerulonephritis

Hematologic

- Anemia of chronic disease
- Asymptomatic leukopenia
- Thrombocytopenia
- lymphadenopathy

- Gastritis/peptic ulcer due to NSAID/corticosteroids


- Pancreatitis, peritonitis, and colitis: due to SLE vasculitis
- Lupoid hepatitis
- hepatosplenomegaly



GI:

Auto antibodies

- The anti-nuclear antibody (ANA) test is the best screening test for SLE and is determined by **immunofluorescence** or **ELISA** tests
- The ANA is positive in significant titer (usually 1:160 or higher) in virtually all patients with SLE



Significance of Autoantibodies in SLE

Antigen	SLE	Clinical Associations
ds DNA	70%	Nephritis (and flare)
Anti RNP	40%	Scleroderma, myositis
Histones	70%	Drug-Induced Lupus
SM Antigen	30%	Severe SLE
Anti ribosomal	20%	Psychosis, Depression
Antiphospholipid	50%	Clotting, fetal loss
SSA/Ro	35%	SCLE, Sjogren's, NLS
SSB/La	15%	SCLE, Sjogren's, NLS
Anti neuronal	60%	Active CNS lupus

Other investigations

- **Anti-double-stranded DNA titers**
- **Complement Levels (CH50, C3, C4)**
- **ESR**
- **CRP**
- **Complement Split products**
- **Decreased complement C1q**

Treatment

NSAIDs

(Non-steroidal anti-inflammatory drugs)

Antimalarials (Hydroxychloroquine)

Immunosuppressive agents

2. Rheumatoid Arthritis

- Is an autoimmune disease in which the normal immune response is directed against an individual's own tissue, including the :
 - Joints
 - Tendons
 - Bones

Resulting in inflammation and destruction of these tissues

Rheumatoid Arthritis (*Contd.*)

- The cause of rheumatoid arthritis is **not known**
 - Investigating possibilities of a foreign antigen, such as a **virus**
- Both prevalence and incidence are 2-3 times greater in **women** than in men

Pathogenesis


(Type III hypersensitivity reaction)

In rheumatoid arthritis, many individuals produce a group of auto-antibodies known as **rheumatoid factor**

These antibodies react with determinants in the **F_C region of IgG**

Rheumatoid Factor

The classic rheumatoid factor is an **IgM** antibody with this kind of reactivity



Pathogenesis

(Type III hypersensitivity reaction)

Such auto-antibodies bind to normal circulating IgG, forming **IgM-IgG complexes** which may be deposited in joints.

This leads to activation of synovial macrophages

The macrophages engulf the immune complexes and then release TNF and other pro-inflammatory cytokines e.g., IL-1


Pathogenesis (Type III hypersensitivity reaction)

TNF induces the secretion of **metalloproteinases**; which are known to cause joint destruction.

T cell activation due to unknown antigens also contributes to the inflammation in RA

Rheumatoid Arthritis

Rheumatoid arthritis (RA) affects peripheral joints and may cause destruction of both cartilage and bone.



Treatment and Prognosis

Medications

- **NSAIDS** (Non-steroidal anti-inflammatory drugs)
- **Disease-modifying drugs** (eg, gold, hydroxychloroquine, sulfasalazine, penicillamine)
- **Immunosuppressive therapy:**
 - Corticosteroids
 - Methotrexate
- **Surgery**
- **Physical therapy**

Take home message

- o The spectrum of autoimmune disorders is wide ranging from single organ involvement to a systemic disease
- o The disease process is usually prolonged and is generally associated with significant morbidity and mortality
- o The mainstay of the treatment is to maintain immunosuppression

Thank you



Autoimmune disorders in a nutshell.
•Beatrice the Biologist•