

THROMBOLYTIC DRUGS

(Fibrinolytic drugs)

By

Prof. Hanan Hagar

Learning objectives

- To know mechanism of action of thrombolytic therapy.
- To differentiate between different types of thrombolytic drugs.
- To describe indications, side effects and contraindications of thrombolytic drugs.
- To recognize the mechanisms, uses and side effects of antiplasmins.

Definition of Thrombolytics

Thrombolytic agents are drugs used to lyse **already** formed blood clots in clinical settings where ischemia may be fatal.

What is the role of thrombolytic therapy in antithrombotic plan ?

- **Thrombolytic therapy** is one part of an overall antithrombotic plan that frequently includes **anticoagulants**, **antiplatelet agents** and mechanical approaches to rapidly restore flow and prevent re-occlusion.
- The goal of **thrombolytic therapy** is **rapid restoration of blood flow** in an occluded blood vessel by **accelerating proteolysis** of the thrombus.

Indications of thrombolytics

used for the treatment of thromboembolic disorders as:

- Acute myocardial infarction (ST elevation, **STEMI**).
- Acute ischemic stroke.
- Peripheral artery occlusion.
- Deep venous thrombosis.
- Pulmonary embolism.

Rational for use of thrombolytic drugs in AMI

- Improvement of ventricular function
- Reduction of the incidence of congestive heart failure
- Reduction of mortality following AMI.
- Thrombolytic drugs need to be given **immediately to the patient after diagnosis of MI**, delay in administration will be of no value.

Mechanism of action of thrombolytic drugs


They have common mechanism of action by stimulating **plasminogen activation** via converting **plasminogen** (proenzyme) to **plasmin** (active enzyme) → lysis of the insoluble fibrin clot into soluble derivatives.

What is plasmin?

Plasmin: is a nonspecific **protease** capable of breaking down:

- **Fibrin**
- **other circulating proteins** including fibrinogen, factor V, and factor VIII.

Plasminogen activators



Types of thrombolytic drugs

Non-fibrin specific

Streptokinase

Anistreplase

Urokinase

Fibrin specific

Tissue plasminogen

Activators (t-PA)


➤ Alteplase

➤ Reteplase

➤ Tenecteplase

Fibrin specific plasminogen activators

Fibrin surface




Fibrin specific plasminogen activators activate **mainly plasminogen bound to clot surface** and have less effect on circulating plasminogen

PAI= plasminogen activator inhibitor

Nonfibrin specific plasminogen activators

Fluid phase



Activate both **plasminogen bound to clot surface** and **circulating plasminogen** in blood leading to extensive systemic plasminogen activation, with degradation of several plasma proteins including fibrinogen, factor V, and factor VIII.

Non fibrin-specific thrombolytic drugs

e.g. Streptokinase – Anistreplase – Urokinase

➤ activates plasminogen both in **blood** and at the **clot surface** thus produces **clot lysis** and **systemic fibrinolysis** leading to **bleeding**.

Fibrin-specific thrombolytic drugs

- are tissue plasminogen activators
- e.g. Alteplase – Reteplase - Tenecteplase
- selective in action (clot or fibrin specific)
- binds preferentially to plasminogen at the **fibrin surface (non-circulating)** rather than **circulating** plasminogen in blood.
- Risk of bleeding is **less than** non specific agents.
- Activity is enhanced upon binding to fibrin.


Non fibrin-specific thrombolytic drugs

- Streptokinase
- Anistreplase
- Urokinase

Streptokinase (SK)

- Is a bacterial protein produced by B-hemolytic streptococci.
- It acts **indirectly** by forming plasminogen-streptokinase complex "**activator complex**" which converts inactive plasminogen into active plasmin.
- Streptokinase is non-fibrin specific
- can degrade **fibrin clots** as well as **fibrinogen** and other plasma proteins.

Mechanism of action of streptokinase



Streptokinase

- T 1/2 = less than 20 minutes.
- given as intravenous infusion (250,000 U then 100,000 U/h for 24-72 h).
- It is the least expensive among others.
- used for venous or arterial thrombosis.

Side effects of streptokinase

- **Antigenicity:** high-titer antibodies develop 1 to 2 weeks after use, precluding retreatment until the titer declines.
- **Allergic reaction:** like rashes, fever, hypotension
- **Bleeding** due to activation of circulating plasminogen (systemic fibrinolysis).
- Not fibrin specific.

Precautions

Not used in patients with:

- Recent streptococcal infections or
- Previous administration of the drug
- These patients may develop **fever, allergic reactions and resistance** upon treatment with streptokinase due to **antistreptococcal antibodies.**

Anistreplase (APSAC)

- Anisoylated Plasminogen Streptokinase Activator Complex (APSAC) acylated plasminogen combined with streptokinase.
- It is a **prodrug**, de-acylated in circulation into the active plasminogen-streptokinase complex.
- $T_{1/2}$ is 70-120 min

Advantages

- Given as a bolus I.V. injection (30 U over 3 - 5 min.).
- Longer duration of action than streptokinase.
- More thrombolytic activity.
- Greater clot selectivity.

Disadvantages

Similar but less than streptokinase alone in:

- ❑ Antigenicity.
- ❑ Allergic reactions.
- ❑ Minimal fibrin specificity
- ❑ Systemic lysis.

But more expensive than streptokinase

Urokinase

- Human enzyme synthesized by the kidney
- obtained from either urine or cultures of human embryonic kidney cells.
- is a **direct** plasminogen activator
- Given by intravenous infusion (300,000U over 10 min then 300,000U/h for 12h).

Urokinase

- Has an elimination half-life of 12-20 minutes.
- Used for the lyses of acute massive pulmonary emboli

Advantages No anaphylaxis (not antigenic).

Disadvantages

- Minimal fibrin specificity
- Systemic lysis (acts upon fibrin-bound and circulating plasminogen).
- Expensive (its use is now limited).

Tissue Plasminogen Activators (t-PAs)

- All are recombinant human tissue plasminogen activators (t-PA).
- Prepared by recombinant DNA technology.
- Include drugs as
 - Alteplase
 - Reteplase
 - Tenecteplase

Mechanism of t-PAs

- They activate **fibrin-bound plasminogen** rather than free plasminogen in blood.
- Their action is **enhanced by the presence of fibrin**.
- They bind to fibrin in a thrombus and convert the entrapped plasminogen to plasmin followed by activated local fibrinolysis with **limited systemic fibrinolysis**.

Advantages of t-PAs

- Fibrin-specific drugs (clot specific).
- Limited systemic fibrinolysis.
- Reduced risk of bleeding
- Not-antigenic (can be used in patients with recent streptococcal infections or antistreptococcal antibodies).

Alteplase

- is a recombinant form of human tPA.
- has very short half life (~5 min)
- is usually administered as an intravenous bolus followed by an infusion.
- (60 mg i.v. bolus + 40 mg infusion over 2 h).

Uses

- In ST-elevation myocardial infarction (STEMI)
- Pulmonary embolism.

Retepase

- A variant of recombinant tPA
- It has longer duration than alteplase (15 min.)
- Has enhanced fibrin specificity
- Given as two I.V. bolus injections of 10 U each

Uses

- In ST-elevation myocardial infarction (STEMI)
- Pulmonary embolism.

Tenecteplase

- Is another modified human t-PA.
- prepared by recombinant technology
- It has half life of more than 30 min.
- It can be administered as a single IV bolus.
- It is more fibrin-specific & longer duration than alteplase.
- It is only approved for use in acute myocardial infarction.

Contraindications to thrombolytics

Absolute contraindications include:

- Active internal bleeding
- Cerebral hemorrhagic stroke
- Recent intracranial trauma or neoplasm
- Major surgery within two weeks

Relative contraindications include:

- Active peptic ulcer
- Severe uncontrolled hypertension

Fibrinolytic Inhibitors

Fibrinolytic inhibitors (**Antiplasmins**) inhibit plasminogen activation and thus inhibit fibrinolysis and promote clot stabilization.

Fibrinolytic Inhibitors

Antiplasmins

Aminocaproic Acid & tranexamic acid

- ✓ acts by competitive inhibition of plasminogen activation
- ✓ Given orally

Aprotinin

- ✓ It inhibits fibrinolysis by blocking the action of plasmin (**plasmin antagonist**)
- ✓ Given orally or i.v.

Uses of Fibrinolytic Inhibitors

- ✓ Adjuvant therapy in hemophilia
- ✓ Fibrinolytic therapy-induced bleeding (antidote).
- ✓ Post-surgical bleeding
- ✓ These drugs work like antidotes for fibrinolytic drugs. Similar to **Protamine** (Antidote of the anticoagulant, heparin) or **Vitamin K** (Antidote of the oral anticoagulant warfarin).