

Glucose Metabolism (Glycolysis)

Objectives

By the end of this lecture, students are expected to:


- Recognize glycolysis as the major oxidative pathway of glucose
- List the main reactions of glycolytic pathway
- Discuss the rate-limiting enzymes/Regulation
- Assess the ATP production (aerobic/anaerobic)
- Define pyruvate kinase deficiency hemolytic anemia
- Discuss the unique nature of glycolysis in RBCs.

Glycolysis: An Overview


- ❖ Glycolysis, the major pathway for glucose oxidation, occurs in the cytosol of all cells.
- ❖ It is unique, in that it can function either aerobically or anaerobically, depending on the availability of oxygen and intact mitochondria.
- ❖ It allows tissues to survive in presence or absence of oxygen, e.g., skeletal muscle.
- ❖ RBCs, which lack mitochondria, are completely reliant on glucose as their metabolic fuel, and metabolizes it by anaerobic glycolysis.

Glycolysis



The product of one reaction is the substrate of the subsequent reaction.



Aerobic Vs Anaerobic Glycolysis




Aerobic Glycolysis (1st and 2nd reactions)




Aerobic Glycolysis

(Reactions: 3rd – 5th)



Aerobic Glycolysis

(Reactions: 6th – 10th)




Regulation: Glucokinase/ Hexokinase


- **Hexokinase** – it is inhibited by the reaction product, glucose-6-P which accumulates when further metabolism of this hexose is reduced
- **Glucokinase** – It is inhibited indirectly by Fructose-6-P and is indirectly stimulated by glucose

Glucokinase (GK) Regulation


- In the presence of high fructose-6-phosphate, GK translocates and binds tightly to **GKRP** (glucokinase regulatory protein) in the nucleus, making it inactive
- When glucose levels are high in blood and hepatocytes (GLUT-2), GK is released from GKRP and enters the cytosol



Regulation: PFK-1




Pyruvate Kinase Covalent Modification



Pyruvate Kinase Deficiency Hemolytic Anemia

PK Mutation may lead to:


1. Altered Enz. Kinetics.
2. Altered response to activator.
3. Decreased the amount of the Enz. or its stability



Long-Term Regulation of Glycolysis

Insulin: Induction

Glucagon: Repression



Summary

(Regulation of Glycolysis)

Regulatory Enzymes (Irreversible reactions):

Glucokinase/hexokinase

PFK-1

Pyruvate kinase

Regulatory Mechanisms:


Rapid, short-term: Allosteric, Covalent modifications

Slow, long-term: Induction/repression

Apply the above mechanisms for each enzyme where applicable

Glycolysis

For each NADH, 3 ATP will be produced by ETC in the mitochondria



Substrate-level phosphorylation vs. Oxidative phosphorylation

- **Phosphorylation** is the metabolic reaction of introducing a phosphate group into an organic molecule.
- **Oxidative phosphorylation:** The formation of high-energy phosphate bonds by phosphorylation of ADP to ATP coupled to the transfer of electrons from reduced coenzymes to molecular oxygen via the electron transport chain (ETC); it occurs in the mitochondria.
- **Substrate-level phosphorylation:** The formation of high-energy phosphate bonds by phosphorylation of ADP to ATP (or GDP to GTP) coupled to cleavage of a high-energy metabolic intermediate (substrate). It may occur in cytosol or mitochondria

Aerobic Glycolysis

(Net ATP produced)

ATP Consumed:

2 ATP

ATP Produced:

Substrate-level $2 \times 2 = 4$ ATP

Oxidative-level $2 \times 3 = 6$ ATP


Total 10 ATP

Net: $10 - 2 = 8$ ATP


Anaerobic Glycolysis

- **NADH produced** cannot be used by ETC for **ATP production**.
(No O₂ and/or No mitochondria)
- **Less ATP production**, as compared to aerobic glycolysis.
- **Lactate is an obligatory end product, Why?**

Because if not formed, All cellular NAD⁺ will be converted to NADH, with no means to replenish the cellular NAD → Glycolysis stops → death of the cell



Lactate Dehydrogenase



Anaerobic Glycolysis

(Net ATP produced)

ATP Consumed:

2 ATP

ATP Produced:


Substrate-level $2 \times 2 = 4$ ATP

Total 4 ATP


Net:

$4 - 2 = 2$ ATP

Anaerobic Glycolysis in RBCs (2,3-BPG Shunt)



Anaerobic Glycolysis in RBCs (2,3-BPG Shunt)



Glycolysis in RBCs

(Net ATP produced)

ATP Consumed:

2 ATP

ATP Produced:

Substrate-level

or $2 \times 2 = 4$ ATP
 $1 \times 2 = 2$

Total

4 or 2 ATP

Net:

$4 - 2 = 2$ ATP

$2 - 2 = 0$ ATP

Glycolysis in RBCs (Summary)

End product:

Lactate

No net production or consumption of NADH

Energy yield:

If no 2,3-BPG is formed: 2 ATP

If 2,3-BPG shunt occurs: 0 ATP

PK Deficiency hemolytic anemia depends on:

Degree of PK Deficiency

Compensation by 2,3-BPG

Take Home Messages

- ❑ Glycolysis is the major oxidative pathway for glucose
- ❑ Glycolysis is employed by all tissues
- ❑ Glycolysis is a tightly-regulated pathway
- ❑ PFK-1 is the rate-limiting regulatory enzyme
- ❑ Glycolysis is mainly a catabolic pathway for ATP production, But it has some anabolic features (amphibolic)
- ❑ Pyruvate kinase deficiency in RBCs results in hemolytic anemia

Take Home Messages

❑ Net energy produced in:

Aerobic glycolysis:	8 ATP
Anaerobic glycolysis:	2 ATP

❑ Net energy produced in glycolysis in RBCs:

Without 2,3 BPG synthesis:	2 ATP
With 2,3 BPG synthesis:	0 ATP

Reference

Lippincott Illustrated Review of Biochemistry, 6th edition, 2014, Unit 2, Chapter 8, Pages 91-108.