Parasitic Helminths and **Arthropod Agents** and Vectors of **Diseases** Dr:MONA BADR

Parasitic Helminths and Arthropod Agents and Vectors of Diseases

Objectives:

By the end of this lecture the student should be able to:

- Name the three main groups of parasitic helminths and their characteristic morphological features.
- •Describe the life cycle of **Ascaris lumbricoides** as an example of parasitic helminths.
- •Discuss the role of arthropods as agents and as vectors of diseases in humans.
- Give examples of the main arthropod vectors of diseases.

Classification of Parasites

Helminthes		
Mulicellular		
Specialized cells		
A- Round worms =		
Nematodes		
cylindrical,		
un-segmented(Ascaris)		
B- <i>Flat worms</i>		
1-Trematodes:		
leaf-like, un-segmented.		
2-Cestodes:		
tape-like, segmented		

Location of helminthes in the body:

- Intestinal helminthes:
- Tissue helminthes:


Nematodes (round worm) intestinal Nematode

General features

- Elongated worm, cylindrical, unsegmented and tapering at both ends.
- 2. Variable in size, measure <1 cm to about 100cm.
- Sex separate and male is smaller than female



Ascaris lumbricoides (roundworm)



Ascaris lumbricoides (roundworm)

The commonest intestinal helminthes can cause infection to human.

Found in jejunum and upper part of ileum.


Female (20-40 cm) which is longer than


male (**10-15 cm)**.

Feed on semi digested food.



Ascaris lumbricoides life cycle





http://www.dpd.cdc.gov/dpdx

Life cycle of Ascais Lumbricoides

It infect the human when man ingest an fertilized egg contaminated with food or water, then this fertilized egg become a Larva that penetrate the wall of the duodenum and enter the blood stream to the heart, liver and enter the pulmonary circulation and stay in the alveoli, where it grow and molts for three weeks then **Larva** passes from respiratory system to be **coughed up**, swallowed returned to the small intestine where it mature to adults male &female, fertilization take place producing eggs which pass in stool.

Pathogenicity


- 1-Migrating LARVA:
- Ascaris pneumonia, some times LARVA reach aberrant sites like brain, heart or spinal cord can cause unusual disturbance.
- 2-Adult WORM:
- The worm consumes proteins and vitamins from host's diet and leads to malnutrition.
- Can cause intussusception, intestinal ulcers and in massive infection can cause intestinal obstruction.





Classification of Parasites

Helminthes		
Mulicellular		
Specialized cells		
A- Round worms =		
Nematodes		
cylindrical,		
un-segmented(Ascaris)		
B- <i>Flat worms</i>		
1-Trematodes:		
leaf-like, un-segmented.		
2-Cestodes:		
tape-like, segmented		




















MEDICAL IMPORTANCE OF ARTHROPODS

- 1)As aetiologic agents (causes) of diseases.
 - Tissue damage Scabies
 - Induction of hypersensitivity reactions.
 - Injection of poisons Scorpions.
 - Entomophobia (acarophobia)
- 2) As vectors of diseases:
 - I: Mechanical transmission simple carriage of pathogens. flies
- II: Biological transmission:
 - cyclical filarial parasite
 - propagative e,g;plaque bacillie in rat fleas
 - cyclopropagative e,g;Malaria in mosquito
- III: Transovarian transmission as ricketsis carried within ticks.


Medical importance of Arthropods as vector of diseases

- I: <u>Mechanical transmission</u> simple carriage of pathogens e,g flies.
 - II: Biological transmission:
- 1- cyclical :cyclical change only but does not multiply in the body of the vector e,g :filarial parasite.
- 2-propagative:when the disease agent undergo no cyclical change but multiplies in the vector e,g;plaque bacillie in rat fleas.
- 3- Cyclo-propagative : the disease agent undergoes cyclical change and multiply in the body of arthropods e,g;Malaria in mosquito.
 - III: Transovarian transmission:
- _transmitted as vector from arthropodes parents to offspring as ricketsis carried within ticks.

Scabies as tissue damage example of Arthropod **As aetiologic agents (causes) of diseases.**













Scabies

الجرب

ARTHROPODS OF MEDICAL IMPORTANCE

Class Insecta الحشرات	Class <i>Arachnida</i> العناكب	القشريات Class Crustacea
• Muscid	• Scorpions العقارب	• Water flea
flies:housefly,Tsetse fly		(Cyclops)
• Myiasis-producing flies .		
• Mosquitoes البعوض:	• Spiders العناكب	
Anopheles, Aedes Culex		
• Sandfly ذباب الرمل	• Ticks: القراد	
(Phlebotomus)	hard, soft	
• Black fly(Simulium)	• Mites السوس	
• Fleas البراغيث	-Sarcoptes	
	scabiei,	
• Lice(Pediculus, Phthirus)	-dust mites	
• Bugs:Cimex,Triatoma البق		
• Bees النحل		

Important arthropod vectors for human diseases

Human discases		
House fly (Musca domestica)	Mechanical transmission of many viruses, bacteria and parasites.	
Mosquitoes البعوض	Anopheles :malaria filariasis Culex: filariasis, viruses Aedes: yellow fever, dengue fever, Rift Valley Fever	
Lice القمل	Body louse: vector for: Relapsing fever, typhus and trench fever.	
البراغيث Fleas	Rat flea is vector for plague due to Yersinia pestis.	
Ticks القراد	Soft ticks, some are vestors for: Borrela duttoni Hard ticks Include vectors for Babesiosis (protozoa), Q fever, and Rocky mountain spotted fever:	
نبابة التسي (Glossina) دبابة	Vector for African Trynanosomiasis (African sleeping sickness)	
الذبابة السوداء (Simulium) الذبابة	Vector for Onchocerca (river blindness)	
Sand fly (Phlebotomus) ذبابة الرمل	Vectors for leishmania and sandfly fever virus.	

Vector for Dracunculus medinensis

Cyclops

LICE Louse(singular), Lice (pleural)

Pediculus humanus



Mosquitoes:

Cosmopolitan , more than 3000 species.




Mouth parts in female adapted to piercing and sucking blood.

Genus and species distinguished by morphology of adult and deveopmetal stages.

cyclo-propagative





Malaria

sand flay transmit Leishmania







