

RESPIRATORY SYSTEM BLOCK

Pathology Practical

Classification of Respiratory Diseases

1. Inflammatory lung diseases:

(Asthma, cystic fibrosis, & COPD)

2. Restrictive lung diseases:

(Allergic Alveolitis)

3. Obstructive lung diseases :

(Bronchial Asthma, Bronchiectasis, & (COPD- Ch. Bronchitis & Emphysema))

4. Respiratory tract infections:

-Upper resp. tract infection (sinusitis, tonsillitis, otitis media, pharyngitis & laryngitis)

-Lower resp. tract infection (Pneumonia & Bronchopneumonia , T.B.)

5. Malignant tumors (Squamous CC, adenocarcinoma, Large CC & Small CC)

6. Benign tumors (Pulmonary hamartoma, pulmonary sequestration)

7. Pleural cavity diseases (eg. Mesothelioma, effusion)

8. Pulmonary vascular diseases (Embolism, edema & hypertension)

9. Neonatal diseases (pulmonary hyperplasia.)

RESTRICTIVE LUNG DISEASES

ALLERGIC ALVEOLITIS

Restrictive Lung Disease (Honeycomb lung):

Major Categories of Chronic Interstitial Lung Disease

Fibrosing

Usual interstitial pneumonia (idiopathic pulmonary fibrosis)
Nonspecific interstitial pneumonia
Cryptogenic organizing pneumonia
Associated with collagen vascular disease
Pneumoconiosis
Associated with therapies (drugs, radiation)

Granulomatous

Sarcoidosis
Hypersensitivity pneumonia

Eosinophilic

Loeffler syndrome
Drug allergy-related
Idiopathic chronic eosinophilic pneumonia

Smoking-Related


Desquamative interstitial pneumonia
Respiratory bronchiolitis

Restrictive Lung Disease (Honeycomb lung) – Gross



- ***Represent 15% of non-infectious diseases of lungs.***
- ***End-stage: diffuse interstitial pulmonary fibrosis (Honeycomb lung).***
- ***Acute: Acute Respiratory Distress Syndrome***
- ***Chronic : Occupational: Asbestosis, silicosis, coal worker pneumoconiosis.***
- ***Interstitial lung disease (interstitial pneumonia), Idiopathic pulmonary fibrosis.***

Restrictive Lung Disease (Honeycomb lung) – Cut section



“Honeycomb” lung. (extensive fibrosis from restrictive lung disease)

The gross appearance, as seen here in a patient with organizing diffuse alveolar damage

Restrictive Lung Disease

– Interstitial pulmonary fibrosis (Usual interstitial pneumonia).

Clinical Features:


Usually manifests insidiously, with the gradual onset of Chronic dry and non-productive cough and progressive dyspnea.

On physical examination (detected by auscultation) have characteristic

Dry (velcro like) crackles during inspiration..

Restrictive Lung Disease

– Interstitial pulmonary fibrosis (Usual interstitial pneumonia).




Most likely to involve:

- Lower lobes.**
- Subpleural areas.**


- Wide alveolar septa.**
- Fibrosis.**
- Chronic inflammation.**
- Dilated bronchioles and alveoli (Honey combed pattern).**

Hypersensitivity Pneumonitis– Histopathology & Radiogram




This case of *extrinsic allergic alveolitis* shows interstitial inflammation along alveolar ducts (bronchiolocentric distribution). The inflammation is diffuse, lacks nodularity, and manifests radiologically as a ground-glass pattern

Hypersensitivity Pneumonitis (HP)



Interstitial bronchiolocentric pneumonitis (Extrinsic allergic alveolitis) with lymphocytes, plasma cells and foamy macrophages in alveolar space and terminal airways . Interstitial fibrosis, obliterative bronchiolitis and intra-alveolar exudate . Nodules of organizing fibroblasts, histiocytes and other inflammatory cells

Hypersensitivity Pneumonitis (HP)



Interstitial fibrosis, lymphocyte infiltration in the alveolar wall, mainly collagen fibers hyperplasia, especially in the bronchioles due to their respective muscle fibers and endothelial cell proliferation and thickening


OBSTRUCTIVE LUNG DISEASES



- 1. Bronchial Asthma**
- 2. Bronchiectasis**
- 3. COPD :**
(Chronic Bronchitis & Emphysema)


1. BRONCHIAL ASTHMA

BRONCHIAL ASTHMA - Anatomy




Bronchial Asthma: Inflammation of the airways causes airflow into and out of the lungs to be restricted. The muscles of the bronchial tree become tight and the lining of the air passages swells, reducing airflow and producing the characteristic wheezing sound

BRONCHIAL ASTHMA - LPF



Between the bronchial cartilage at the right and the bronchial lumen filled with mucus at the left is a submucosa widened by smooth muscle hypertrophy, edema, and inflammation (mainly eosinophils). These are changes of bronchial asthma. The peripheral eosinophil count or the sputum eosinophils can be increased during an asthmatic attack.


BRONCHIAL ASTHMA - HPF



Changes of bronchial asthma: the numerous eosinophils are prominent from their bright red cytoplasmic granules in this case of bronchial asthma


2. BRONCHIECTASIS

Diagram of Normal & Bronchiectatic Bronchus




In Bronchiectasis, mucus production increases, the cilia are destroyed or damaged, and areas of the bronchial wall become chronically inflamed and are destroyed .

Bronchiectasis – Gross pathology




Bronchiectasis occurs when there is obstruction or infection with inflammation and destruction of bronchi so that there is permanent dilation.

Bronchiectasis – Gross pathology





- **Red** arrow: Dilated bronchi and bronchioles in the affected lung.
- **Blue** arrow: Fibrous adhesions between the lobes.

Bronchiectasis – LPF



Section of a dilated bronchial wall with florid acute on chronic inflammation of the bronchial wall and surrounding interstitial fibrosis.

Bronchiectasis – HPF



- **This is Bronchiolar wall.**
- **There is Chronic inflammation.**

Bronchiectasis – may be associated with Congenital or hereditary conditions such as:

- ***Cystic fibrosis.***
- ***Kartagener OR (immotile cilia syndrome).***

Bronchiectasis – complications:

- ***Amyloidosis.***
- ***Metastasizing abscesses or brain abscess.***

Chronic Obstructive Pulmonary Diseases (COPD)

also known as

**Chronic Obstructive Lung Disease (COLD),
Chronic Obstructive Airway Disease (COAD),
Chronic Obstructive Respiratory Disease (CORD)**

Include:

3. Chronic Bronchitis

4. Emphysema,

a pair of commonly co-existing diseases of the lungs in which the airways narrow over time.

3. CHRONIC BRONCHITIS

Normal vs Chronic bronchitis




Normal




Chronic Bronchitis

Chronic Bronchitis - LPF




Inflammatory infiltrate in bronchial walls is composed of lymphocytes and plasma cells. In the lumen desquamated epithelial cells (catarrhal inflammation) .In mucosa often occurs mataplasia of cylindric cilliated epithelium into multilayered squamous epithelium. Goblet cells are hyperplastic, hyperplastic are also the sero-mucous glands in the submucosal layer. Muscularis mucosae is hypertrophic

Chronic Bronchitis - LPF



Early - hypersecretion of mucus in large airways with hypertrophy of submucosal glands in tracheobronchial tree . Later - increase in goblet cells in small airways contributes to excessive mucus production and airway obstruction .

Chronic Bronchitis - HPF



● ***Chronic inflammatory infiltrates range from absent to prominent
Increased percentage of bronchial wall is occupied by submucosal mucous glands, as measured
by Reid index; this directly correlates with sputum production, variable dysplasia, squamous
metaplasia, bronchiolitis obliterans***

4. EMPHYSEMA

Emphysema – Clinical Features




Emphysema patient (so-called -pink puffers) exhibit dyspnea without significant hypoxemia and tend to be thin, to have hyperinflated lung fields at total lung capacity, and to be free of signs of right heart failure

Normal Lung




Emphysema



A


Normal Lung



B

Emphysema

Emphysema – Gross Anatomy




The chest cavity is opened at autopsy to reveal numerous large bullae apparent on the surface of the lungs in a patient dying with emphysema. Bullae are large dilated airspaces that bulge out from beneath the pleura.

Emphysema – Gross pathology



Dilated airspaces in emphysematous lung. Although there tends to be some scarring with time because of superimposed infections, the emphysematous process is one of loss of lung parenchyma, not fibrosis

Bullous Emphysema – Gross pathology




A bulla is defined as an emphysematous space larger than 1 cm.

Centrilobular Emphysema – Gross pathology




Centrilobular emphysema : Fixed, cut surface of a lung shows multiple cavities lined by heavy black carbon deposits characteristic of smoking.

Panacinar Emphysema – LPF




Some of the alveolar septae are ruptured and the ruptured septa project with in air spaces on the form of spurs.

Panacinar Emphysema – LPF



Some of the alveolar septae are ruptured and the ruptured septa project within air spaces on the form of spurs.

Panacinar Emphysema - HPF




Destruction of tissue leaves emphysematous spaces with little surface area, few capillaries, and large air spaces. Large vessel at lower left


LOWER RESPIRATORY TRACT INFECTIONS

1. Lobar Pneumonia

Lobar Pneumonia - Gross pathology




Lobar Pneumonia - Gross pathology



A closer view of the lobar pneumonia demonstrates the distinct difference between the upper lobe and the consolidated lower lobe.


Lobar Pneumonia : X - Ray

- **Consolidation** seen in X-ray film.
- Involving Right middle lobe.




Lobar Pneumonia of the right middle lobe

Lobar Pneumonia - Histopathology



- * **Congestion (first 2 days)**
- * **Red hepatisation (fibrinous alveolitis) (2nd to 4th day)**
- * **Grey hepatisation (leukocytic alveolitis) (4th to 8th day)**
- * **Resolution (after 8th day)**

Lobar Pneumonia - HPF



- **Thickened alveolar wall.**
- **Fibrinous exudate.**
- **Inflammatory cells: macrophages and neutrophils.**

Lobar Pneumonia - complications

- Lung abscesses.**
- Empyema.**
- Meningitis.**
- Septic arthritis.**
- Infective endocarditis.**


2. Bronchopneumonia

Bronchopneumonia – Gross pathology




- **The consolidated areas here very closely match the pattern of lung lobules (hence the term "lobular" pneumonia).**
- **Bronchopneumonia is classically a "hospital acquired" pneumonia seen in persons already ill from another diseases e.g. DM , old age , immune deficiency process.**

Bronchopneumonia – Cut section




This bronchopneumonia is more subtle, but there are areas of lighter tan consolidation. The hilum is seen at the lower left with radiating pulmonary arteries and bronchi

Bronchopneumonia – X-Ray




This radiograph demonstrates patchy infiltrates consistent with a bronchopneumonia from a bacterial infection.

Bronchopneumonia – Histopathology




Bronchopneumonia (Lobular pneumonia) is an acute exudative inflammation of the lungs characterised by foci of consolidation surrounded by normal parenchyma. Usually, bronchopneumonia affects one or more lobes and is bilateral.

Bronchopneumonia – LPF



Section of the lung shows foci of inflammatory consolidation surrounding bronchioles: Bronchioles are filled with an inflammatory purulent exudate and show ulceration of mucosa, focal inflammation and necrosis of walls . Surrounding lung parenchyma shows congestion and edema

Bronchopneumonia – MPF



At high magnification, the alveolar exudate of mainly neutrophils is seen. The surrounding alveolar walls have capillaries that are dilated and filled with RBC's. Such an exudative process is typical for bacterial infection. This exudate gives rise to the productive cough of purulent yellow sputum seen with bacterial pneumonias



PULMONARY EMBOLUS AND INFARCTION

Thromboembolism in the Lung – Gross



A large pulmonary thromboembolus is seen in the pulmonary artery to the left lung. Such thromboemboli typically originate in the leg veins or pelvic veins of persons who are immobilized

Thromboembolism in the Lung – Gross



Large thromboemboli can cause death. Medium sized thrombemboli (blocking a pulmonary artery to a lobule or set of lobules) can produce the lesion seen here -a hemorrhagic pulmonary infarction which is a wedge-shaped and based on the pleura.

Pulmonary embolus and infarction in the Lung

A Longitudinal transection of a lung showing a wedge shaped peripheral hemorrhagic infarction .

A thrombus is seen in a major branch of pulmonary artery (arrow head) .




Pulmonary Embolus with infarction – CT scan




CT scan - white arrows show pulmonary embolus with lung infarction

Pulmonary artery thromboembolus - LPF




Microscopic appearance of a pulmonary thromboembolus in a large pulmonary artery. There are interdigitating areas of pale pink and red that form the "lines of Zahn" characteristic for a thrombus.

Small pulmonary artery thromboembolus - HPF



A small peripheral pulmonary artery thromboembolus. If these small PE are showered into the pulmonary circulation at once or over a period of time will lead to pulmonary hypertension.

Fat Embolism in the Lung - HPF



The rounded holes that appear in the vascular spaces here in the lung are fat emboli. Fat embolization syndrome occurs most often following trauma with fracture of long bones that releases fat globules into the circulation which are trapped in pulmonary capillaries

SECOND PRACTICAL


1. TUBERCULOSIS

2. CANCER OF THE LUNG

TUBERCULOSIS


- **Epithelioid and giant cell Granuloma, Ghon's complex or caseation is present**
- **Complications of TB are:**
 - Amyloidosis ,
 - Tuberculous pneumonia
 - Miliary tuberculosis
 - Tuberculous meningitis
 - Addison disease .

Pulmonary TB – Caseous Necrosis – Gross




On closer inspection, the granulomas have areas of caseous necrosis. This pattern of multiple caseating granulomas primarily in the upper lobes is most characteristic of secondary (reactivation) tuberculosis

Pulmonary TB – Caseous Necrosis – Gross



Extensive caseation and the granulomas involve a larger bronchus causing soft, necrotic center to drain out and leave behind a cavity. Cavitation is typical for large granulomas with TB. Cavitation is more common in the upper lobes.


Pulmonary TB - Ghon's Complex – Gross Pathology




The Ghon's complex is seen here at closer range. Primary tuberculosis is the pattern seen with initial infection with tuberculosis in children. Reactivation, or secondary tuberculosis, is more typically seen in adults.

Miliary TB of the Lungs

- **Miliary TB can occur when TB lung lesions erode pulmonary veins or when extrapulmonary TB lesions erode systemic veins.**
- **This results in hematogenous dissemination of tubercle bacilli producing myriads of 1-2 mm. lesions throughout the body in susceptible hosts.**
- **Miliary spread limited to the lungs can occur following erosion of pulmonary arteries by TB lung lesions.**



Miliary TB of the Lungs – Cut section




This is a "miliary" pattern of granulomas because there are a multitude of small tan granulomas, about 2 to 4 mm in size, scattered throughout the lung parenchyma. The miliary pattern gets its name from the resemblance of the granulomas to millet seeds.

Miliary TB of the Lungs – X-Ray




This chest x-ray shows a patient with miliary TB.

Tuberculous Granulomas - LPF




At low magnification, this micrograph reveals multiple granulomas. Granulomatous disease by chest radiograph appear as reticulonodular densities.

Tuberculous Granulomas - HPF




Well-defined granulomas are seen here. They have rounded outlines. The one toward the center of the photograph contains several Langhan's giant cells. Granulomas are composed of transformed macrophages called epithelioid cells along with lymphocytes, occasional PMN's, plasma cells, and fibroblasts

Pulmonary TB - Granuloma with central early necrosis




The pyknotic nuclei of epithelioid cells in the center of the granuloma (apoptotic bodies) are a precursor of necrosis.

Tuberculous Granulomas - HPF




The edge of a granuloma is shown here at high magnification. At the upper is amorphous pink caseous material [1] composed of the necrotic elements of the granuloma as well as the infectious organisms. This area is ringed by the inflammatory component [2] with epithelioid cells, lymphocytes, and fibroblasts.

Epithelioid & Giant cell Granulomas in *Tuberculosis*



At high magnification, the granuloma demonstrates that the epithelioid macrophages are elongated with long, pale nuclei and pink cytoplasm. The macrophages organize into committees called giant cells. The typical giant cell for infectious granulomas is called a Langhan's giant cell and has the nuclei lined up along one edge of the cell

Acid Fast bacilli of Mycobacterium TB in the Lung



*A stain for **Acid Fast Bacilli** is done (**AFB** stain) to find the mycobacteria .
The mycobacteria stain as red rods, as seen here at high magnification.*

LUNG CARCINOMA

TWO TYPES OF LUNG CARCINOMA

- **NON-SMALL CELL CARCINOMA**
 1. **SQUAMOUS CELL CARCINOMA**
 2. **ADENOCARCINOMA**
 3. **LARGE CELL CARCINOMA**
- **SMALL CELL CARCINOMA**

The NON-small cell cancers behave and are treated similarly, the SMALL cell carcinomas are WORSE than the non-small cell carcinomas, but respond better to chemotherapy, often drastically!

1. Squamous Cell Carcinoma of the lung


- **Most commonly found in men and correlated with smoking.**
- **Associated with ectopic parathyroid hormone secretion.**
- **Pathology: If more differentiated → more cytoplasm, keratin whorls.**
- **Transforms from carcinoma in situ.**
- **Grading is based on the amount of keratin & cytoplasm.**

Squamous Cell Carcinoma of the Lung - Gross




This is a squamous cell carcinoma of the lung that is arising centrally in the lung (as most squamous cell carcinomas do). It is obstructing the right main bronchus. The neoplasm is very firm and has a pale white to tan cut surface.

Squamous Cell Carcinoma of the Lung - Gross




This is a larger squamous cell carcinoma in which a portion of the tumor demonstrates central cavitation, probably because the tumor outgrew its blood supply.

Squamous Cell Carcinoma of the Lung – CT scan




This chest CT scan view demonstrates a large squamous cell carcinoma of the right upper lobe that extends around the right main bronchus and also invades into the mediastinum and involves hilar lymph nodes.

Squamous Cell Carcinoma of the Lung - HPF




Microscopic appearance of squamous cell carcinoma with nests of polygonal cells with pink cytoplasm and distinct cell borders. The nuclei are hyperchromatic and angular.

Squamous Cell Carcinoma of the Lung - HPF




In this squamous cell carcinoma at the upper right is a squamous eddy with a keratin pearl. At the left, the tumor is less differentiated and several dark mitotic figures are seen

Squamous Cell Carcinoma of the Lung - HPF



Neoplastic squamous cells show pleomorphism, hyperchromatism, individual cell keratinization, mitoses and areas of necrosis.

Squamous Cell Carcinoma of the Lung - HPF




The pink cytoplasm with distinct cell borders and intercellular bridges characteristic for a squamous cell carcinoma of the lung

2. Adenocarcinoma of the lung

- *The most common type of lung cancer, making up 30-40% of all cases.*
- *Glandular differentiation by tumor cells and 80% of those cells produce mucin.*
- *Not as strongly associated with a smoking history as compared to Squamous or Small Cell Carcinomas*
- *Adenocarcinoma in situ - called bronchoalveolar carcinoma*
- *Early and distant metastases*

Adenocarcinoma of the Lung – Gross




A peripheral adenocarcinoma of the lung. Adenocarcinomas and large cell anaplastic carcinomas tend to occur more peripherally in lung. Adenocarcinoma is the one cell type of primary lung tumor that occurs more often in non-smokers and in smokers who have quit.


Adenocarcinoma of the Lung

CT scan

Ill-defined, peripheral and midzonal right lung mass.




Adenocarcinoma of the Lung – LPF



Adenocarcinoma in Situ (Previously named Bronchioloalveolar Carcinoma)


Invasive Adenocarcinoma of the Lung – HPF



- **Malignant glands.**
- **Pleomorphic cells with prominent nucleoli.**
- **Desmoplasia.**

Gene which is commonly mutated in adenocarcinoma is **Epithelial Growth Factor Receptor 1 (EGFR1)**.

Adenocarcinoma of the Lung – HPF




Differentiated malignant glands lined by pleomorphic and hyperchromatic malignant cells showing conspicuous nucleoli

3. Large Cell Carcinoma of the lung

- **Can be a neuroendocrine carcinoma. Probably represents undifferentiated SCC and adenocarcinomas.**
- **Large nuclei, prominent nucleoli.**
- **Variation in size and shape.**
- **Nuclei normally do not touch due to more cytoplasm.**
- **Moderate amount of cytoplasm.**
- **Early and distant metastases, sometimes cavitating.**


Large Cell Carcinoma of the Lung – Gross



Undifferentiated Large Cell Carcinoma of the Lung – Gross




Large Cell Carcinoma of the Lung – HPF




Pleomorphic carcinoma of lung (large cell and giant cell subtype). It shows mixed composition of large cell carcinoma and pleomorphic multinucleated giant cells (arrows). (H and E, ×200)

Large Cell Carcinoma of the Lung – HPF



This section from lower respiratory tract shows neoplastic cells with abundant pale eosinophilic cytoplasm and a surrounding infiltrate of inflammatory cells

Large Cell Carcinoma of the Lung – HPF



This section shows neoplastic cells with abundant pale eosinophilic cytoplasm and pleomorphic multinucleated giant cells

Small cell carcinoma of the lung

- **Highly Malignant Tumor.**
- **Cells are small, with scant cytoplasm, ill-defined borders, finely granular chromatin (salt & pepper pattern) and absent or inconspicuous nucleoli.**
- **High mitotic count and often extensive necrosis.**
- **Typically not graded as all SCLC are considered High Grade.**
- **Very strong relationship with smoking.**
- **Often lead to paraneoplastic syndromes.**

Small Cell Carcinoma of the Lung “Oat cell” – Gross




Arising centrally in this lung and spreading extensively is a small cell anaplastic (oat cell) carcinoma. The cut surface of this tumor has a soft, lobulated, white to tan appearance. The tumor seen here has caused obstruction of the main bronchus to left lung so that the distal lung is collapsed

Small Cell Carcinoma of the Lung “Oat cell” – Gross




Oat cell carcinoma which is spreading along the bronchi. The speckled black rounded areas represent hilar lymph nodes with metastatic carcinoma

Small Cell Carcinoma of the Lung “Oat cell” : X-Ray




This chest radiograph demonstrates a mass lesion in the right upper lobe. This was an oat cell carcinoma (yellow arrow). It obstructed the right main bronchus, leading to atelectasis on the right, evidenced by a raised right hemidiaphragm. The patient aspirated gastric contents, producing a diffuse pneumonia (blue arrow) on the left (since aspirated material could not pass the obstruction on the right).

Small Cell Carcinoma of the Lung “Oat cell” – HPF



This is the microscopic pattern of a small cell anaplastic (oat cell) carcinoma in which small dark blue cells with minimal cytoplasm are packed together in sheets.

Small cell carcinoma “Oat cell” of the lung - HPF




Section of the tumor shows clusters of malignant cells which are small , round , ovale , or spindle shaped with prominent nuclear molding , finely granular nuclear chromatin (salt and pepper pattern) , high mitotic count and focal necrosis

Metastatic tumours of the lung

METASTATIC TUMORS

- **LUNG** is the **MOST COMMON** site for all metastatic tumors, regardless of the site of origin.
- It is the site of **FIRST CHOICE** for metastatic sarcomas for purely anatomic reasons !

Metastatic Tumors of the Lung – Gross & X-ray




Multiple variably-sized masses are seen in all lung fields. These tan-white nodules are characteristic for metastatic carcinoma. Metastases to the lungs are more common even than primary lung neoplasms



Chest X-ray showing multiple cannon ball opacities in both lung fields.

Metastatic Tumors of the Lung – Gross & CT scan




Here are larger but still variably-sized nodules of metastatic carcinoma in lung.




CT Lung shows Cannonball Metastases-large, hematogenously spread metastatic lesions in the lungs of varying sizes most often from colon, breast, renal, thyroid primaries

Metastatic Tumors of the Lung – LPM



A nest of metastatic infiltrating ductal carcinoma from breast is seen in a dilated lymphatic channel in the lung. Carcinomas often metastasize via lymphatics.

Metastatic Tumors of the Lung – LPF



A focus of metastatic carcinoma from breast is seen on the pleural surface of the lung. Such pleural metastases may lead to pleural effusions, including hemorrhagic effusions, and pleural fluid cytology can often reveal the malignant cells


Mesothelioma of the lung

Mesothelioma of the Lung – Gross




The dense white encircling tumor mass is arising from the visceral pleura and is a mesothelioma. These are big bulky tumors that can fill the chest cavity. The risk factor for mesothelioma is asbestos exposure.

Mesothelioma of the Lung – Gross




RESPIRATORY: Pleura: Mesothelioma: Gross natural color external view of lung with nodules of tumor on pleura

Mesothelioma of the Lung – MPF



Mesotheliomas have either spindle cells or plump rounded cells forming gland-like configurations, as seen here at high power microscopically. They are very difficult to diagnose cytologically.

Mesothelioma of the Lung – HPF



Mesothelioma: Micro epithelial pattern spindle cells or plump rounded cells forming gland-like configurations