


# Effects of exercise on the respiratory system.

**Dr. Aida Korish**  
**Assoc. Prof Physiology**  
**KSU**





When we exercise, we need more oxygen

Dr. Aida Korish ( [iaidakorish@yahoo.com](mailto:iaidakorish@yahoo.com) )

# Objectives

**By the end of this lecture the students should be able to: -**

- 1-Describe the effects of moderate and severe exercise on oxygen consumption, and ventilation volumes.
- 2- Interpret the effects of exercise on arterial  $PO_2$ ,  $PCO_2$  and  $H^+$  ions.
- 3-Define the diffusing capacity of the respiratory membrane, and its typical values at rest, and explain its changes in exercise.
- 4-Explain causes of hyperventilation in exercise.

# The respiratory system and exercise

When we exercise more oxygen is needed by the working muscles and more carbon dioxide must be removed from the muscles.

As a result:

- our rate of breathing increases;
- we increase the depth of our breathing, up to our vital capacity;
- we increase the blood flow through the lungs;
- we increase the oxygen taken up and used by the body.

Oxygen used during exercise can be up to twenty times a person's normal oxygen uptake.



# Effect of Exercise on the respiratory system

- The blood gases do not always have to become abnormal for respiration to be stimulated in exercise.
- Instead, respiration is stimulated mainly by neurogenic mechanisms during exercise.



# Regulation of respiration during exercise.

In strenuous exercise  $O_2$  consumption and  $CO_2$  formation may increase 20 folds but alveolar ventilation increases almost exactly in step with the increased levels of metabolism.

Therefore the arterial  $PO_2$ ,  $PCO_2$ ,  $pH$  all remain almost exactly normal.




Figure 41-8

Effect of exercise on oxygen consumption and ventilatory rate. (From Gray JS: Pulmonary Ventilation and Its Physiological Regulation. Springfield, Ill: Charles C Thomas, 1950.)




Figure 41-9

Changes in alveolar ventilation (bottom curve) and arterial  $PCO_2$  (top curve) during a 1-minute period of exercise and also after termination of exercise. (Extrapolated to the human being from data in dogs in Bainton CR: Effect of speed vs grade and shivering on ventilation in dogs during active exercise. J Appl Physiol 33:778, 1972.)

# What causes intense ventilation during exercise?


- The brain, on transmitting motor impulses to the exercising muscles, transmits at the same time collateral impulses into the brain stem to excite the respiratory center.
- A large share of the total increase in ventilation begins immediately on initiation of the exercise, before any blood chemicals have had time to change.
- It is likely that most of the increase in respiration results from neurogenic signals transmitted directly into the brain stem respiratory center at the same time that signals go to the body muscles to cause muscle contraction.





# What cause intense ventilation during exercise?

- Neural signals from the motor areas of the brain to the respiratory center.
- The joint proprioceptors
- Body temperature (hypothalamus).
- Possibility that the neurogenic factor for control of ventilation during exercise is a learned response.



## Interrelation Between Chemical Factors and Nervous Factors in the Control of Respiration During Exercise.

- Direct nervous signals stimulate the respiratory center *almost* the proper amount to supply the extra oxygen required for exercise and to blow off extra carbon dioxide.
- Occasionally, the nervous respiratory control signals are either too strong or too weak.
- Then chemical factors play a significant role in bringing about the final adjustment of respiration required to keep the O<sub>2</sub>, CO<sub>2</sub>, and H<sup>+</sup> ion concentrations of the body fluids as nearly normal as possible.

# The Neurogenic Factor for Control of Ventilation During Exercise Is a Learned Response.

- The ventilatory response during exercise, is at least partly a *learned* response.
- That is, with repeated periods of exercise, the brain becomes progressively more able to provide the proper signals required to keep the blood PCO<sub>2</sub> at its normal level.
- The cerebral cortex is involved in this learning, because experiments that block only the cortex also block the learned response.






Figure 41-10

Approximate effect of maximum exercise in an athlete to shift the alveolar PCO<sub>2</sub>-ventilation response curve to a level much higher than normal. The shift, believed to be caused by neurogenic factors, is almost exactly the right amount to maintain arterial PCO<sub>2</sub> at the normal level of 40 mm Hg both in the resting state and during heavy exercise.


# Diffusion capacity of the respiratory membrane

- Is the volume of gas that diffuses through the membrane each minute for a pressure difference of 1mmHg.
- **Diffusing capacity for oxygen at rest**  
***21ml/min/mmHg***
- Even if the oxygen pressure difference across the respiratory membrane is 11mmHg----- $11 \times 21 = 230$ ml oxygen diffusing through the membrane each minute.
- During rest tissues consume 250 ml O<sub>2</sub> /min




# Changes in the oxygen- diffusing capacity during exercise

- **65ml/min/mmHg**
- This is due to increased number of open pulmonary capillaries which was dormant, thereby increasing the surface area for gas exchange.
- In addition to increased alveolar ventilation.




# Diffusing capacity for carbon dioxide

- It diffuses 20 times greater than oxygen due to greater diffusion coefficient which is 20 times that for oxygen.
- Diffusion capacity for carbon dioxide 400ml/min/mmHg.
- During exercise 1200 to 1300ml/min/mmHg.




- During exercise the oxygen requirement increased 20 times, and cardiac output increased and so the time blood remained in the pulmonary capillaries becomes less than half normal despite the fact that additional capillaries open up.
- But the blood is almost completely saturated with oxygen when it leaves the pulmonary capillaries.





## *Reasons for this are as follow:*

- 1- The diffusing capacity for oxygen increases almost three fold during exercise, this results mainly from increasing numbers of capillaries participating in the diffusion, and a more even V/Q ratio all over the lung.
- 2- At rest the blood normally stays in the lung capillaries about three times as long as necessary to cause full oxygenation. Therefore, **even with shortened time of exposure in exercise, the blood is still fully oxygenated or nearly so.**



# Oxygen Consumption and Pulmonary Ventilation in Exercise.

- ⊙ Normal oxygen consumption for a young man at rest is about 250 ml/min.
- ⊙ However, under maximal conditions,
- ⊙ this can be increased to approximately the following average levels:
- ⊙ 

	ml/min
⊙ Untrained average male	-----3600
⊙ Athletically trained average male	---4000
⊙ Male marathon runner	----- 5100



Gasping for air after a hard race, to repay the oxygen debt.



*Runners at the end of a race are often left gasping for air*

Dr.Aida Korish ( [iaidakorish@yahoo.com](mailto:iaidakorish@yahoo.com))