#### RESPIRATORY BLOCK

# PHOSPHOLIPIDS OF CLINICAL SIGNIFICANCE

DR. USMAN GHANI

## **OBJECTIVES**

By the end of this lecture the First Year students will be able to:

- Identify the types and functions of phospholipids
- Discuss the physiological importance of phospholipids
- Understand the role of glycerophospholipids in lung surfactant and their clinical implications in respiratory distress syndrome (RDS)
- Identify the classes and physiological functions of phospholipase enzymes

## **OVERVIEW**

- Types and functions of phospholipids
- Glycerophospholipids: Types, functions and role in lung surfactant, cell signaling and protein anchoring
- Respiratory distress syndrome (RDS)
- Sphingophospholipids
- Phospholipids in lipoprotein particles
- Phospholipases: Types and functions

## **PHOSPHOLIPIDS**

- Phospholipids are polar, ionic compounds that contain an alcohol group attached either to:
  - Diacylglycerol or
  - Sphingosine
- Major lipids of cell membranes


#### Two classes:

- Glycerophospholipids
- Sphingophospholipids

## **PHOSPHOLIPIDS**

 Their hydrophobic (non-polar) portion is attached to the membrane

 Their hydrophilic (polar) portion extends outward interacting with the aqueous environment




## **FUNCTIONS OF PHOSPHOLIPIDS**

- Membrane-bound phospholipids act as:
  - Reservoir for intracellular messengers
  - Anchors to cell membranes

- Nonmembrane-bound phospholipids act as:
  - Lung surfactant
  - Components of bile (as detergents to solubilize cholesterol)

## **GLYCEROPHOSPHOLIPIDS**

- Also called phosphoglycerides
- Contain glycerol
- A major class of phospholipids
- All contain phosphatidic acid
   (PA)
- PA is the simplest phospholipid



## **GLYCEROPHOSPHOLIPIDS**


Phospholipids are derived from PA such as:

Serine + PA	Phosphatidylserine (PS)	Cell signaling Blood clotting
Ethanolamine+PA	Phosphatidylethanolamine (PE) (cephalin)	
Choline + PA	Phosphatidylcholine (PC) (lecithin)	Lung surfactant
Inositol + PA	Phosphatidylinositol (PI)	Cell signaling
Glycerol + PA	Phosphatidylglycerol (PG)	Lung surfactant

## SOME EXAMPLES

## Cardiolipin


- Two molecules of PA joined to an additional molecule of glycerol through PO<sub>4</sub> groups
- In the inner mitochondrial membrane
- Function: maintenance of respiratory complexes of electron transport chain



## SOME EXAMPLES

## Platelet activating factor (PAF)

- Binds to cell surface receptors
- Triggers thrombotic and acute inflammatory reaction



## ROLE OF PC IN LUNG SURFACTANT

- Lung surfactant is a complex mixture of:
  - Lipids (90%) including
     Dipalmitoylphosphatidylcholine (DPPC)
  - Proteins (10%)
- Alveolar cells of the lungs are lined by the extracellular fluid layer
- Alveolar cells secrete DPPC (a major lung surfactant)

## ROLE OF PC IN LUNG SURFACTANT

 Surfactant decreases the surface tension of the fluid layer

• Reduces pressure needed to re-inflate alveoli

Prevents alveolar collapse (atelectasis)

## ROLE OF PC IN LUNG SURFACTANT


## Respiratory distress syndrome (RDS)

- In preterm infants due to deficiency of lung surfactant
- A major cause of neonatal death
- Treatment: Glucocorticoids to mother to promote lung maturation
- In adults due to damaged alveoli by infection or trauma

## ROLE OF PI IN CELL SIGNALING

 Plays important role in intracellular signaling

 PI is part of calciumphosphatidyl inositol system






Figure 17.8


Role of inositol trisphosphate and diacylglycerol in intracellular signaling.

## ROLE OF PI IN MEMBRANE PROTEIN ANCHORING

 Anchoring of proteins to membranes through carbohydrate-PI bridge

## **Examples:**

- Alkaline phosphatase (on the surface of small intestine)
- Acetylcholine esterase (on postsynaptic membrane of neurons)
- Anchoring proteins can be cleaved by phospholipase C enzyme




## **SPHINGOPHOSPHOLIPIDS**

 A long-chain fatty acid attached to sphingosine

• Example: Sphingomyelin

 An important component of myelin that protects and insulates nerve fibers



#### Figure 17.4

Structure of sphingomyelin, showing sphingosine (in green box) and ceramide components (in dashed box).

## PHOSPHOLIPIDS IN LIPOPROTEIN PARTICLES

 The outer core of lipoprotein particles is hydrophilic

 Contains phospholipids and free cholesterol

 Allows transport of core lipids in aqueous plasma





Figure 18.14
Structure of a typical lipoprotein particle.

## **PHOSPHOLIPASES**

- Phospholipids are degraded by phospholipase enzymes
- Present in all tissues including pancreatic juice
- Glycerophospholipids are degraded by:
  - Phospholipase A<sub>1</sub>, A<sub>2</sub>, C, D
- Sphingophospholipids are degraded by:
  - Sphingomyelinase

## FUNCTIONS OF PHOSPHOLIPASES

- Digestion of phospholipids by pancreatic juice
- Important for remodeling of phospholipids
- Production of second messengers
- Pathogenic bacteria produce phospholipases to dissolve cell membranes and spread infection



**Figure 17.11**Degradation of glycerophospholipids by *phospholipases*.

## TAKE HOME MESSAGE

- Phospholipids are complex lipids that perform important physiological functions in the body
- Membrane-bound phospholipids are involved in cell signaling, protein anchoring and myelin protective functions
- Nonmembrane-bound phospholipids function as lung surfactant and as detergent in the bile
- Phospholipases are enzymes that degrade phospholipases
- They are important for remodeling of phospholipids

## REFERENCES

• Lippincott's Illustrated Reviews, Biochemistry, 6<sup>th</sup> Edition, Denise R. Ferrier, Lippincott Williams & Wilkins, USA, pp 201-207.