
## RESPIRATORY SYSTEM (III)

Histology of the Lower Respiratory Tract (Trachea, Bronchi, Bronchioles) & the Lung



## **Objectives:**

By the end of this lecture, the student should be able to describe:

#### 1- The microscopic structures of the wall of:

- Trachea.
- Primary or extrapulmonary bronchi.
- Intrapulmonary (secondary and tertiary) bronchi.
- Bronchioles.

#### 2- The microscopic structures of:


- Interalveolar septum.
- Alveolar phagocytes.
- Pleura.

## TRACHEA

#### The wall of trachea is formed of:

- (1) Mucosa.
- (2) Submucosa.
- (3) Adventitia.






## **MUCOSA OF TRACHEA**


- (1) Epithelium: Respiratory epithelium
- (2) Lamina propria.



It is formed of elastic fibers.

It separates lamina propria from submucosa.








## SUBMUCOSA OF TRACHEA

#### Contents:


- 1- C.T.
- 2- Numerous mucous & seromucous glands.
- 3- Lymphoid elements.



#### ADVENTITIA OF TRACHEA

#### Contents:

- 1- Fibroelastic C.T.
- 2- C-shaped rings (12-16) of hyaline cartilage.
  - Trachealis muscle (bundle of smooth muscle fibers) connects the 2 ends of each C-shaped (incomplete) rings of cartilage.







## EXTRAPULMONARY BRONCHUS' (1ry BRONCHUS')

Generally have the same histological appearance as the trachea.

# INTRAPULMONARY BRONCHI (2ry & 3ry BRONCHI)

- 1- Mucosa.
- 2- Muscle coat.
- 3- Submucosa.
- 4- Adventitia.






## INTRAPULMONARY BRONCHUS

#### (1) Mucosa:

- a- Epithelium: Respiratory epith.
- b- Lamina propria.N.B. No elastic lamina.



#### (2) Muscle coat (complete):


Two distinct layers of smooth muscle fibers spirally arranged in opposite direction.

#### INTRAPULMONARY BRONCHUS

#### (3) Submucosa:

C.T. contains:

- a- Seromucous glands.
- b- Lymphoid elements.




#### (4) Adventitia:

Contents: a- Loose C.T.

b- Irregular plates of hyaline cartilage (complete layer).

c- Solitary lymphoid nodules



#### **BRONCHIOLES**


1- Preterminal (1ry) Bronchioles (Bronchioles):

Are less than 1mm in diameter.



3- Respiratory (3ry) Bronchioles.





Supplemental Slide 1

#### Preterminal Bronchioles

- (1) Mucosa: has longitudinal folds:
  - A- Epithelium:

Simple ciliated columnar epith. with occasional goblet cells.




- B- Lamina propria: C.T. rich in elastic fibers.
- (2) Smooth muscle: 2 helically arranged smooth muscle layers.
- (3) Adventitia: C.T.
- N.B. <u>No cartilage</u>, <u>No seromucous glands</u>, <u>No lymph</u> <u>nodules</u>.


#### Terminal Bronchioles

Similar structure to preterminal bronchioles, but: **Epithelium:** 

Simple cuboidal partially ciliated epithelium With Clara cells (With NO goblet cells).


N.B. Are less than 0.5mm in diameter.






## Respiratory Bronchioles

Are similar in structure to terminal bronchioles **But:** their walls are interrupted by the presence of few pulmonary alveoli.






## CLARA CELLS

#### **Structure:**

columnar cells (non ciliated).


#### **Function:**


- 1- Degrade toxins in inhaled air.
- 2- Divide to regenerate the bronchiolar epith.
- 3- Produce surfactant-like material.



#### **ALVEOLAR DUCTS**

The wall of alveolar ducts consist almost of pulmonary alveoli.





N.B. Alveolar duct  $\rightarrow$  ends by: atrium  $\rightarrow$  communicates with: 2-3 alveolar sacs


## **PULMONARY ALVEOLI**


#### **Definition:**

They are small out-pouching of respiratory bronchioles, alveolar ducts & alveolar sacs.

## **Topics:**

- \*Interalveolar septa.
- \*Alveolar epithelium.
- \* Alveolar phagocytes (Lung macrophages).






## INTERALVEOLAR SEPTA

#### **Definition:**

The region between 2 adjacent alveoli.



## **Components:**

(A) Alveolar Epithelium:

lines both sides of interalveolar septum.


(B) Interstitium.

## ALVEOLAR EPITHELIUM

## (1) Type I Pneumocytes

## (2) Type II Pneumocytes





## ALVEOLAR EPITHELIUM

## (1) Type I Pneumocytes:

- line 95% of the alveolar surface.
- Count: less numerous than type II pneumocytes.
- L/M: simple squamous epith.

#### -Function:


Exchange of gases.





## (2) Type II Pneumocytes:

- Line 5% of the alveolar surfaces.
- Are more numerous than type I pneumocytes.
- Are cuboidal or rounded cells,
   With Foamy cytoplasm.
   Nucleus: central & rounded.
- The cytoplasm contains membrane-bound Lamellar bodies

  (contain pulmonary surfactant).







## **Type II Pneumocytes:**

#### **Function:**

1- Synthesis & secretion of pulmonary surfactant.

2- Renewal of alveolar epithelial cells:

Type II cells can divide to regenerate both type I & type II pneumocytes.

## Interstitium of interalveolar septa

(1) Continuous Pulmonary Capillaries.

#### (2) Interstitial C.T.:

a- C.T. Fibers: elastic fibers &

type III collagen (reticular fibers).

b- C.T. Cells: Fibroblasts, Macrophages,

Mast cells, Lymphocytes.

## BLOOD-GAS BARRIER (BLOOD-AIR BARRIER)

Nucleus of endothelial cell

Surfactant — (surface lining)

Alveolar

epithelium Fused basal laminae

Endothelium

Alveolar

lumen

CO2

0.1-1.5 um

Capillary

lumen

#### **Definition:**

It is the region of the interalveolar septum that is

traversed by O2 & CO2

#### **Components:**

- 1- Thin layer of surfactant.
- 2- Type I pneumocyte.
- 3- Fused basal laminae of type I pneumocytes & endothelial cells of the pulmonary capillary.
- 4- Endothelial cells of the pulmonary capillary.

## Alveolar phagocytes


(Alveolar Macrophages)
(Dust Cells)

#### Sites:

- (1) In the lumen of pulmonary alveoli.
- (2) In the interstitium of interalveolar septa.

#### Function:

Phagocytose particulate matter (e.g. dust) & bacteria in the lumen of pulmonary alveoli and in the interstitium of interalveolar septa.





## <u>Pleura</u>

Is formed of two layers:
Parietal and visceral.
It is formed of simple squamous mesothelium. The two layers are separated by serous fluid. The visceral layer has sub-epithelium loose C.T that extends into the lung tissue

## THANK YOU

