#### **Cholesterol Metabolism**

Dr. Usman Ghani

1 Lecture

Cardiovascular Block

# Objectives

- Understand the structure and functions of cholesterol
- Discuss the regulation of cholesterol homeostasis in the body
- Comprehend the important steps of cholesterol synthesis pathway
- Identify different levels of regulation of cholesterol synthesis
- Discuss the association of hypercholesterolemia with abnormal cholesterol metabolism
- Understand the role of statins in the treatment of hypercholesterolemia


#### Overview

- Introduction
- Cholesterol structure
- Cholesteryl esters
- Cholesterol synthesis
- Rate limiting step
- Regulation of cholesterol synthesis
- Regulation of HMG CoA reductase
- Excretion of cholesterol
- Hypercholesterolemia and treatment


#### Cholesterol

- Most important animal steroid
- Maintains membrane fluidity
- Insulating effect on nerve fibres
- Cholesterol is the parent molecule for
  - Bile acids and bile salts
  - Steroid hormones
  - Vitamin D<sub>3</sub>

Liver plays a central role in the regulation of cholesterol homeostasis



#### **Cholesterol Structure**




#### Cholesteryl esters

- Most plasma cholesterol is esterified with a fatty acid
- CEs are not present in membranes
- Present in small amounts in most cells
- More hydrophobic than cholesterol

# Cholesterol synthesis

- Synthesized in all tissues
- Major sites for synthesis: liver, adrenal cortex, testes, ovaries and intestine
- All carbon atoms are derived from acetyl CoA
- Enzymes involved in biosynthesis are partly located in ER and partly in cytoplasm


### Synthesis of HMG CoA



- HMG CoA is present in both cytosol and mitochondria of liver
- Mitochondrialketogenesis
- Cytosolic –
   cholesterol synthesis


## Synthesis of mevalonic acid

- Rate limiting and key step
- Occurs in cytosol
- HMG CoA reductase is an ER membrane enzyme with catalytic unit hanging in the cytosol



### Further steps in synthesis

- Production of a 5-carbon unit:
  - Isopentinyl pyrophosphate (IPP)
- Condensation to a 30C compound: squalene
- Cyclization of squalene to 30C lanosterol
- Synthesis of 27-Carbon cholesterol (defect in this leads to Smith-Lemli-Opitz Syndrome)



# Regulation of Cholesterol Synthesis


 HMG CoA reductase is the rate-limiting enzyme of cholesterol synthesis

#### **HMG CoA Reductase Regulation**

- Sterol-dependent regulation of gene expression
- Sterol-accelerated enzyme degradation
- Sterol-independent phosphorylation/dephosphorylation
- Hormonal regulation

# Sterol-dependent regulation of gene expression of HMG CoA

- When sufficient cholesterol is present, transcription is suppressed and vice versa
- Sterol Regulatory Element (SRE) is a recognition sequence in the DNA
- SREBP (SRE binding protein) binding to SRE is essential for transcription of this gene
- SREBP cleavage-activating protein (SCAP) is an intracellular cholesterol sensor



Copyright @ 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins

### Sterol-dependent regulation

#### **Cholesterol High**


- SCAP binds to insig protein (insulin-induced protein) in ER membrabe
- SCAP-SREBP is retained in the ER
- Down regulation of cholesterol synthesis

#### **Cholesterol Low**

- SCAP-SREBP moves to Golgi bodies
- SCAP is removed from SREBP
- SREBP binds to SRE in DNA
- HMG CoA gene is activated

# Enzyme phosphorylation and dephosphorylation

- AMP- activated protein kinase (AMPK) for phosphorylation
- Phosphorylated form of enzyme is inactive
- Dephosphorylated form is active
- Low ATP or High AMP → cholesterol synthesis decreases




Copyright @ 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins

## Hormonal Regulation

- Insulin and thyroxine increase upregulation of enzyme expression
- Glucagon and cortisol have opposite effect

# HMG CoA Reductase Regulation



#### Excretion of cholesterol

- By conversion into bile acids and bile saltsexcreted in the feces
  - Secretion of cholesterol in bile
  - Transported to intestine for elimination
- In the intestine, some cholesterol is converted by bacteria into coprostanol and cholestanol before excretion

#### Hypercholesterolemia

- High conc. of cholesterol in blood
- Leads to atherosclerosis
- Statin drugs are used to decrease plasma cholesterol levels
- Statins are structural analogs of HMG CoA reductase
- Statins inhibit enzyme activity by competitive inhibition

# **β-Sitosterols/ Phytosterols**

- Plant sterols are poorly absorbed by humans
- Block the absorption of dietary cholesterol
- Clinically useful in the dietary treatment of hypercholesterolemia

#### Take home message

- Cholesterol is important various body functions
- Liver plays a major role in the cholesterol homeostasis in the body
- HMG CoA reductase is a rate-limiting enzyme for cholesterol synthesis

#### References

Lippincott's Biochemistry 5th Edition, Chapter
 18, pp 219–224