

# Transplantation

Immunology Unit  
College of Medicine  
King Saud University

# Objectives

- To understand the diversity among human leukocyte antigens (HLA) or major histocompatibility complex (MHC)
- To know the role of HLA antigens in transplant rejection
- To be familiar with types of immune responses mediating transplant rejections and importance of tissue matching
- To understand the principles of management after transplantation

# Major Histocompatibility Complex and Transplantation

- Major histocompatibility complex (MHC) proteins were discovered for the first time with the advent of **tissue transplantation**
- The success of tissue and organ transplantation depends upon the donor's and recipient's "**human leukocyte antigens**" (HLA) encoded by HLA genes
- These proteins are **allo-antigens**

# MHC Class I and II Proteins

- MHC Class I are glycoproteins found on surface of virtually all the **nucleated cells**
  - Cytotoxic T cell kills virus infected cells in association with **class I MHC** proteins
- MHC Class II glycoproteins are normally found on the surface of **antigen presenting** cells (macrophages, B cells, dendritic cells and Langerhans cells)
  - Helper T cell recognize antigen in association with **class II MHC** proteins

# Major Histocompatibility Complex and Transplantation

- Genes for HLA proteins are clustered in the MHC complex located on the short arm of chromosome 6
- Three genes HLA-A, HLA-B and HLA-C code for Class I MHC proteins
- HLA-D loci encode for Class II MHC proteins ie, DP, DQ and DR

# Major Histocompatibility Complex and Transplantation

Each individual has two “*haplotypes*” i.e, two sets of these genes one paternal and one maternal

<b>MHC class</b>	<b>I</b>			<b>II</b>			<b>III</b>	
<b>Region</b>	<b>A</b>	<b>B</b>	<b>C</b>	<b>DP</b>	<b>DQ</b>	<b>DR</b>	<b>C4, C2, BF</b>	
<b>Gene products</b>	<b>HLA-A</b>	<b>HLA-B</b>	<b>HLA-C</b>	<b>DP</b>	<b>DQ</b>	<b>DR</b>	<b>C' proteins</b>	<b>TNF- α TNF- β</b>
<b>Polymorphisms</b>	<b>47</b>	<b>88</b>	<b>29</b>	<b>More than 300 HLA-D</b>				


# Minor HLA genes and Transplantation

- Minor HLA genes – unknown
  - They mount a **weak immune response**
  - Play role in **chronic rejection** of a graft
  - There are **no laboratory tests** to detect minor antigens

# Transplantation antigens

Slide 4-2

## MHC alleles control allograft rejection



From Abbas, Lichtman, & Pober: Cellular and Molecular Immunology. W.B. Saunders, 1999, Fig. 4-2a



# Transplantation

- Types of transplants:
  - Autografts, Autologous grafts
    - Donor and recipient are same individual
    - Common in skin grafting; bone marrow
  - Syngeneic grafts or (isograft)
    - Donor and recipient are genetically identical
    - Animal models; identical twins


# Transplantation

- Types of transplants:
  - **Allogeneic grafts**
    - Donor and recipient are same species, but genetically unrelated
    - Common heart, lung, kidney, liver graft
  - **Xenogeneic grafts**
    - Donor and recipient are different species
  - **Artificial grafts**


# Transplantation (Rejection)

- Major Barrier to transplantation is the immune response
  - T cells play primary role
  - B cells can/do play a role
  - Classic adaptive/acquired immune response
    - Memory
    - Specificity

# 1<sup>st</sup> set versus 2<sup>nd</sup> set reactions



# Role of CD4<sup>+</sup> versus CD8 T<sup>+</sup> cells



Injecting recipient mice with monoclonal antibodies to deplete one or both types of T cells


# Transplantation

- T cells play primary role in 1st and 2nd set rejection reactions
  - Nude mice accept allografts (no T cells due to genetic modification resulting in **absent thymus**)
  - B cell deficient mice reject allografts




Nude mouse has a transplant of rabbit skin

# Mechanisms involved in Graft Rejection




# Rejection Response





# Clinical manifestations of graft rejection

- I. Hyperacute rejection: very quick
- II. Acute rejection: about 10 days (cell mediated)
- III. Chronic rejection: months-years (both)



# Chronic Rejection

- This occurs months to years after engraftment
- Main pathologic finding in chronic rejection is atherosclerosis of the vascular endothelium
- Main cause of chronic rejection is not known
  - Minor histo-compatibility antigen miss match

# Graft-versus-Host (GVH) Reaction


- Occurs in about two thirds of bone marrow transplants
- Occurs because grafted immunocompetent T cells proliferate in the irradiated immunocompromised host and reject cells with foreign proteins resulting in severe organ dysfunction
- Donor's Tc cells play a major role in destroying the recipient's cells
- Symptoms are: maculopapular rash, jaundice, hepatosplenomegaly and diarrhea
- GVH reactions usually end in infections and death

# HLA Typing in the Laboratory

- Prior to transplantation laboratory test commonly called as ***HLA typing or tissue typing*** to determine the closest MHC match between the donor and recipient is performed
- **Methods**
  - DNA sequencing by Polymerase Chain Reaction (PCR)
  - Serologic Assays
  - Mixed Lymphocyte Reaction (MLR)
  - Crossmatching – (Donor) lymphocytes +(Recipient) serum + complement.


# Tissue Matching

Effect of HLA class I & II matching on survival of kidney grafts



Curve no.	HLA mismatches (no.)	
	Class I	Class II
1	0	0
2	1 or 2	0
3	3 or 4	0
4	0	1 or 2
5	1 or 2	1 or 2
6	3 or 4	1 or 2


# Tissue Matching



# General Immunosuppression Therapy

- 1) Mitotic inhibitor: azathioprine (pre & post)
- 2) Corticosteroids
- 3) Cyclosporin
- 4) Total lymphoid irradiation


# Immunosuppressive Therapy





# Specific Immuno-suppression therapy

- a) Monoclonal antibodies against T cell components or cytokines
- b) Agents blocking co-stimulatory signal



# Immuno-suppressive Therapy

- Downsides
  - Must be maintained for life
  - Toxicity
  - Susceptibility to infections
  - Susceptibility to tumors

# Take home message

- HLA or MHC molecule miss-match can stimulate humoral and cell mediated immunity which is the main cause of rejection of transplants
- Cell mediated immune responses play a major role in transplant rejection
- Tissue matching particularly for HLA-D antigens is important for successful transplantation
- Immuno-suppressive therapy is usually required after transplantation

**Thank you**