



GLUTEAL REGION & BACK OF THIGH



By :
Prof.Saeed Abulmakarem
&
Dr. Sanaa Al-Shaarawi



OBJECTIVES

- ❖ At the end of this lecture, the student should be able to identify :
- ❖ Contents of gluteal region:
 - Groups of Glutei muscles and small muscles (Lateral Rotators).
 - Nerves & vessels.
- ❖ Foramina and structures passing through them as:
 - 1-Greater Sciatic Foramen.
 - 2-Lesser Sciatic Foramen.
- Back of thigh : Hamstring muscles.

CONTENTS OF GLUTEAL REGION


- **I - Muscles:**

- **A- GLUTEI:**

1. Gluteus maximus
2. Gluteus medius
3. Gluteus minimus


- **B- GROUP OF SMALL MUSCLES (Lateral Rotators) :**

1. Piriformis.
2. Obturator internus
3. Superior gemellus
4. Inferior gemellus
5. Quadratus femoris



CONTENTS

- II – NERVES:**
(all from sacral plexus) :
- 1. Sciatic nerve.**
 - 2. Superior gluteal n.**
 - 3. Inferior gluteal n.**
 - 4. Post. cutaneous n. of thigh.**
 - 5. Nerve to obturator internus.**
 - 6. Nerve to quadratus femoris.**
 - 7. Pudendal nerve.**




CONTENTS

III - VESSELS:


(all from internal iliac vessels):

1. Superior gluteal
2. Inferior gluteal
3. Internal pudendal vessels.



Greater sciatic foramen

A




➤ Greater sciatic notch of hip bone is transformed **into foramen** by **sacrotuberous & sacrospinous ligaments**.

- *Structures passing through Greater sciatic foramen :*
- **Piriformis muscle.**
- **Above piriformis :**
- **Superior gluteal nerves & vessels.**
- **Below piriformis :**
- **Inferior gluteal nerves & vessels.**
- **Sciatic nerve.**
- **Posterior cutaneous nerve of thigh.**
- **Nerve to quadratus femoris.**
- **Nerve to obturator internus.**
- **Pudendal N.**
- **Internal pudendal vessels.**

Lesser sciatic foramen


A



- **Lesser sciatic notch of hip bone is transformed into foramen by Sacrotuberous & sacrospinous ligaments.**
- *Structures passing through Lesser sciatic foramen :*
- **Tendon of obturator internus.**
- **Nerve to obturator internus.**
- **Pudendal nerve.**
- **Internal pudendal vessels.**


Glutei Muscles

- **ORIGINS :**
- **Gluteus minimus:**
- **Anterior part of the gluteal surface of ilium**
- **Gluteus medius:**
- **Middle part of the gluteal surface of ilium,**
- **Gluteus maximus:**
- **Posterior part of the gluteal surface of ilium,**
- **Main origin:**
Back of sacrum & coccyx & back of Sacrotuberous ligament



Glutei Muscles

- **Insertion :**
- **Gluteus minimus:**
anterior surface of
the **greater**
trochanter
- **Gluteus medius:**
lateral surface of
the **greater**
trochanter
- **Gluteus maximus:**
 1. **Main insertion:**
iliotibial tract
 2. **Other insertion:**
gluteal tuberosity of
the femur.



- **Gluteus medius & minimus:**

- **Nerve supply:**

- Superior gluteal nerve.

- **Action:**

- **abduction & medial rotation of hip joint.**

- Normally they prevent lateral tilt of the pelvis by contraction of ABDUCTORS of opposite side, on raising the other limb from ground.

- If the pelvis tilts, this is means +ve Trendelenburg's sign.

- Right pelvic tilt as in picture means paralysis of abductors of the opposite side.

- **Gluteus maximus:**

- **Nerve supply:**

- Inferior gluteal nerve.

- **Action:**

- **Extension & lateral rotation of the hip joint.**

- Through its attachment to iliotibial tract, it stabilizes the femur on tibia during standing.

NERVE SUPPLY & ACTION of Gluti muscles






Figure 6a: Trendelenburg gait – inadequate pelvic stability;



Left Figure 6b: normal gait – adequate pelvic stability


- **Obturator Internus:**
- **Origin:**
- Inner surface of the side wall of the pelvis.
- **Insertion:**
- Into the medial surface of the greater trochanter.
- **Nerve supply:**
- Nerve to obturator internus.
- **Superior & Inferior Gemelli:**
- **Origin:**
- Superior gemellus; upper part of lesser sciatic notch.
- Inferior gemellus; lower part of lesser sciatic notch.
- **Insertion:**
- Upper & lower parts into tendon of obturator internus.
- **Nerve supply:**
- Superior gemellus: nerve to obturator internus
- Inferior gemellus: nerve to quadratus femoris.

Small muscles (Lateral Rotators)



- **Piriformis:**
- **Origin:**
- Pelvic surface of **middle 3 sacral vertebrae.**
- **Insertion:**
- It passes through **GSF** to be inserted into the *upper border* of the **greater trochanter.**
- **Nerve supply:**
- Anterior rami of S1,2
- **Quadratus femoris:**
- **Origin:**
- Lateral border of the ischial tuberosity.
- **Insertion:**
- Quadrate tubercle & intertrochanteric crest.
- **Nerve supply:**
- Nerve to quadratus femoris.

Small muscles (Lateral Rotators)



Action: all have **SIMILAR ACTION:**
Lateral rotation of the hip joint.
Control movement of the hip joint.

SUPERIOR GLUTEAL N. :

- **Course:**
- Passes through GSF, above piriformis, then *between* gluteus medius & minimus
- **Branches:**
 1. Muscular to gluteus medius, minimus & tensor fasciae lata
 2. Articular to hip joint


INFERIOR GLUTERAL N. :

- **Course:**
- passes through GSF, below piriformis, then *deep to* gluteus maximus
- **Branches:** muscular to gluteus maximus

NERVE TO QUADRATUS FEMORIS N. :

- **Course:**
- passes through GSF, below piriformis
- **Branches:**
 1. Muscular to quadratus femoris & inferior gemellus
 2. Articular to hip joint

NERVES



POSTERIOR CUTANEOUS NERVE OF THIGH :

Course:

Passes through GSF, below piriformis, then descends deep to deep fascia.

Branches:

Cutaneous branches to: gluteal region, back of scrotum (labium majus) back of thigh & upper part of back of leg.

SCIATIC :


Course:

passes through GSF, below piriformis, then superficial to: ischial spine, superior gemellus, tendon of obturator internus, inferior gemellus, quadratus femoris & adductor magnus.

Branches:


- No branches in gluteal region,
- Divides into tibial & common peroneal nerves, in the middle of back of thigh

NERVES



POSTERIOR COMPARTMENT OF THE THIGH

CONTENTS



- **Muscles:**
- **Hamstring muscles:**
 - Biceps femoris.
 - Semitendinosus.
 - Semimembranosus.
 - Ischial part of adductor magnus.
- **Blood supply:**
 - Branches of the profunda femoris artery.
- **Nerve supply:**
 - Sciatic nerve.

Biceps Femoris :




- **Origin:**
 - **The long head** from the *ischial tuberosity*.
 - **The short head** from the *linea aspera* .
- **Insertion:**
 - Mainly into the *head of the fibula*.
- **Nerve supply:**
 - **The long head** is supplied by the *tibial part of sciatic*;
 - **The short head** is supplied by the *common peroneal part of the sciatic*.
- **Action :**
 - Flexion of knee.
 - Lateral rotation of flexed leg.
 - **Long head:** extends hip.

SEMITENDINOSUS



- **Origin:**
Ischial tuberosity.
 - **Insertion:**
Upper part of the medial surface of the shaft of the tibia (**SGS**)..
- Nerve supply:**
- **Tibial portion of the sciatic.**
- Action:**
- Flexes and medially rotates the leg at the knee joint;
 - Extends the thigh at the hip joint.

SEMIMEMBRANOSUS



- **Origin:**

- Ischial tuberosity.

- **Insertion:**

- Posterior surface of the **medial condyle** of the **tibia**.

- *It forms the oblique popliteal ligament*, which reinforces the capsule on the back of the knee joint.


Nerve supply:

- **Tibial portion of the sciatic nerve.**

Action:


- Flexes and medially rotates the leg at the knee joint;
- Extends the thigh at the hip.

ADDUCTOR MAGNUS (HAMSTRING PART)




- **Origin:**
 - Ischial ramus and **ischial tuberosity**
- **Insertion:**
 - **Adductor tubercle** of the **medial condyle** of the **femur**.
- **Nerve supply:**
 - The **tibial portion of the sciatic**.
- **Action:**
 - Extends the thigh at the **hip joint**.

BLOOD SUPPLY



- The **four perforating branches** of the **profunda femoris artery** (**deep artery of thigh**) provide a rich blood supply to this compartment.
- The **profunda femoris vein** drains the greater part of the blood from the compartment.

NERVE SUPPLY



- **Sciatic Nerve**
- The **sciatic nerve**, a branch of the sacral plexus (**L4 and 5; S1, 2, and 3**), leaves the gluteal region as it descends in the midline of the thigh.
- It is **overlapped posteriorly by** the adjacent margins of the **biceps femoris** and **semimembranosus** muscles.
- **It lies on** the posterior aspect of the **adductor magnus**.
- **In the lower third of the thigh** it ends by dividing into the **tibial** and **common peroneal nerves**.

THANK YOU