# Hip, Knee & Ankle Joints

### Dr Jamila EL Medany

## **OBJECTIVES**


At the end of the lecture, students should be able to:


- List the type & articular surfaces of the hip, knee and ankle joints.
- Describe the capsule and ligaments of the hip, knee and ankle joints.
- Describe movements of hip, knee and ankle joints and list the muscles involved in these movements.
- List important bursae in relation to knee joint.
- Apply Hilton's law about nerve supply of joints.

# **HIP JOINT**


#### **TYPE:**

- <u>Synovial</u>, <u>Ball & Socket</u> joint.
- ARTICULAR SURFACES:
  - Acetabulum of hip (pelvic) bone
  - Head of femur
- Acetabular labrum:
- C-shaped fibro-cartilaginous collar attached to margins of acetabulum, increases its depth for better retaining of head of femur.






### CAPSULE



- Attached <u>Medially</u> to hip bone :
- 1. Labrum acetabulare.
- 2. Transverse acetabular ligament.
- <u>Posteriorly :</u>
- Halfway along the posterior aspect of the neck. (**Part** of the neck lies **inside** the capsule and the other part is **outside** it).
- <u>Anteriorly</u> : the neck of the femur is completely inside the capsule

## Ligaments: (3) Extracapsular



#### **Iliofemoral ligament:**

Y-shaped

- Located anterior to joint
- limits extension




- Located antero-inferior to joint
- limits abduction & lateral rotation

#### **Ischiofemoral ligament:**

Ischiofemoral ligament

- Located posterior to joint
- limits medial rotation





Transverse acetabular ligament: converts acetabular notch into foramen through which pass acetabular vessels


Ligament of femoral head: carries vessels to head of femur

Acetabular labrum


### **Movements**

- Flexion: Iliopsoas (mainly), sartorius, pectineus, rectus femoris.
- <u>Extension</u>: Hamstrings (mainly), gluteus maximus (powerful extensor).
- Abduction: Gluteus medius & minimus, sartorius.
- Adduction: Adductors, gracilis.
- Medial rotation: Gluteus medius & minimus.
- Lateral rotation: Gluteus maximus, quadratus femoris, piriformis, obturator externus & internus.

## **BLOOD SUPPLY**




- The main arterial supply is from branches of the **circumflex femoral** arteries especially the **medial**).
- The blood passes to the joint through :
- (1) Retinacular fibers of the neck.
- (2) Ligament of the head of the femur.




- Damage of the retinacular fibers as in **fracture neck** of the femur can results in
- A vascular necrosis of the head of the femur.
- Fracture neck of the femur is common after age of (60) years especially in women because of Osteoporosis.

## **STABILITY OF THE JOINT**



- The hip joint is one of the **most stable** joints of the body because of :
- (1) The Head of the femur fits very accurately in the acetabulum due to the following
- A. The acetabulum is very deep and its depth is increased by the labrum acetabulare.
- **B.** The labrum acetabulare forms a firm grip on the head of the femur.
- **C.** The atmospheric pressure resists separation between the head of the femur and the acetabulum.
- (2) The three strong Extrinsic ligaments.
- (3) The surrounding strong Muscles.


## **DISLOCATION OF HIP JOINT**



**Figure 15.16** Congenital dislocation of left hip – note the extra skin creases in the upper thigh

### <u>CONGENITAL</u>

- More common in girls and associated with inability to adduct the thigh.
- The upper lip of the acetabulum fails to develop adequately.
- The head of the femur rides up out of the acetabulum onto the gluteal surface of the ileum.



#### • TRAUMATIC:

- It is common in motor vehicle accidents when the thigh is flexed and adducted.
- The dislocated head is displaced **posteriorly** to lie on the posterior surface of the ileum.
- In posterior dislocation the sciatic nerve is liable to be injured.

## **KNEE JOINT**




### • FUNCTIONS:

- 1. Weight bearing.
- 2.Essential for daily activities: standing walking & climbing stairs.
- 3.The main joint responsible for sports: running, jumping, kicking etc.


### **Type & Articular Surfaces**

Knee joint is formed of:

- Three bones.
- Three articulations.
- Femoro-Tibial articulation:
   between the 2 femoral condyles & upper surfaces of the 2 tibial condyles
- <u>Type</u>: synovial, modified hinge
- Femoro-Patellar articulation: between posterior surface of patella & patellar surface of femur
- <u>Type:</u> synovial, plane.








 Is <u>Deficient Anteriorly</u> & replaced by: *quadriceps femoris tendon, patella & ligamentum patellae.* <u>Possesses 2 openings</u>: one for popliteus tendon & one for communication with suprapatellar bursa.

### Ligaments: (4) Extracapsular




- 1. Ligamentum patellae (patellar ligament): from patella to tibial tuberosity.
- 2. Medial (tibial) collateral ligament: from medial epicondyle of femur to upper part of medial surface of tibia (firmly attached to medial meniscus).
- 3. <u>Lateral (fibular) collateral ligament</u>: from lateral epicondyle of femur to head of fibula
- 4. <u>Oblique popliteal ligament:</u> extension of semimembranosus tendon.

## **INTRA Capsular Structures : Menisci**

 They are 2 C-shaped plates of fibro-cartilage attached by anterior & posterior horns, to the articular surface of tibia.

#### **FUNCTION:**

- They deepen articular surfaces of tibial condyles.
- They serve as cushions between tibia & femur.
  - Lateral Meniscus:
  - Small & Circular.
  - Its outer border is separated from lateral collateral ligament by popliteal tendon.




#### Medial Meniscus:

- Large & Oval.
- Its outer border is (firmly attached to the capsule & medial collateral ligament.

medial meniscus is less mobile
 & more liable to be injured.

### **CRUCIATE Ligaments**

- Two in number, situated in the middle of the joint.
- They are called *Cruciate* because they cross each other
- Have received the names
   Anterior and Posterior, from the
   position of their attachments to
   the tibia.




#### **Anterior cruciate ligament:**

• Extends from anterior part of intercondylar area of tibia to posterior part of lateral condyle of femur.

#### **Posterior cruciate ligament:**

• Extends from posterior part of intercondylar area of tibia to anterior part of medial condyle of femur.

## **Functions of Cruciate Ligaments**




#### Anterior ligament:

- Prevents posterior displacement of the femur on the tibia and the tibia from being pulled anteriorly when the knee joint is extended.
- It is taught in <u>Hyper extension</u>.
- Posterior ligament:
- prevents anterior displacement of the femur on the tibia and the tibia from being pulled posteriorly when the knee joint is flexed.
- It is taught in **<u>Hyper flexion</u>**.

### **Bursae Related to Knee joint**

- Suprapatellar bursa: between femur & quadriceps tendon, <u>communicates</u> with synovial cavity of knee joint (Clinical importance?)
- Prepatellar bursa: between patella & skin.
- <u>Deep infrapatellar bursa</u>: between tibia & ligamentum patella.
- <u>Subcutaneous infrapatellar bursa</u>: between tibial tuberosity & skin.
- <u>Popliteal bursa</u> between popliteus tendon & capsule, <u>communicates</u> with synovial cavity of knee joint.




## **MOVEMENTS**

#### FLEXION:

- 1. Mainly by hamstring muscles: biceps femoris, semitendinosus & semimembranosus.
- 2. Assisted by sartorius , gracilis & popliteus.
- EXTENSION: Quadriceps femoris.
- ACTIVE ROTATION (PERFORMED WHEN KNEE IS FLEXED):
- A) MEDIAL ROTATION:
- 1. Mainly by semitendinosus & semimembranosus.
- 2. Assisted by sartorius & gracilis.
- **B)** LATERAL ROTATION:


**Biceps femoris.** 

## **INACTIVE (DEPENDANT) ROTATION**



#### • A) LOCKING OF KNEE:


- The joint assumes the position of **full extension**.
- It becomes a rigid structure.
- The menisci are compressed between the tibial and femoral condyles.
- Results mainly by tension of anterior cruciate ligament.
- Tightening of all the major ligaments.
- The femur is **medially** rotated on the tibia (Lateral rotation of tibia).



### • <u>B) UNLOCKING OF KNEE:</u>

- Medial rotation of tibia (Lateral rotation of femur), at the beginning of flexion
- Performed by <u>Popliteus</u> to relax ligaments & allow easy flexion.


## **STABILITY OF THE JOINT**




#### **1. <u>Muscles :</u>**

- QUADRICEPS particularly the inferior fibers of the Vasti lateralis and medialis.
- Many sport injuries can be preventable through appropriate training and conditioning of the muscle.
- 2. Ligaments :
- The knee joint can function well following a ligamentous strain if the quadriceps is intact.

### **Skeleton of Foot**



### **ANKLE JOINT**



#### **ARTICULAR SURFACES:**

UPPER: A socket formed by: Lateral malleolus, the lower end of tibia & medial malleolus. LOWER: Body of talus.


### Ligaments


### MEDIAL (DELTOID) LIGAMENT:

- A strong triangular ligament.
- Apex: attached to medial malleolus.
- Base: subdivided into 4 parts:
- 1. Anterior tibiotalar part.
- 2. Tibionavicular part.
- 3. Tibiocalcaneal part.
- 4. Posterior tibiotalar part.

#### LATERAL LIGAMENT:

- Composed of 3 separate ligaments
- Anterior talofibular ligament.
- Calcaneofibular ligament.
- Posterior talofibular ligament.





### **Movements**

#### **DORSIFLEXION:**

Performed by muscles of <u>anterior</u> compartment of leg *(tibialis anterior, extensor hallucis longus, extensor digitorum longus & peroneus tertius)*.

#### **PLANTERFLEXION:**

- Initiated by soleus.
- Maintained by gastrocnemius.
- <u>Assisted</u> by other muscles in <u>posterior</u> compartment of leg *(tibialis posterior, flexor digitorum longus & flexor hallucis longus)* + muscles of <u>lateral</u> compartment of leg *(peroneus longus & peroneus brevis)*.

INVERSION & EVERSION MOVEMENTS occur on the talocalcaneo-navicular (subtalar ) joint (Not on ankle joint).

### **Nerve Supply of all JOINTS**

### HILTON'S LAW:

*"The joint is supplied by branches from nerves supplying muscles acting on it".* 

