


PECTORAL REGION AND AXILLA


- *By:*
- *Prof.Saeed Abulmakarem.*
- *DR.Snaa AL-Shaarawy.*



OBJECTIVES


- By the end of the lecture the students should be able to :
- Identify and describe the muscles of the pectoral region.
 - Pectoralis major.
 - Pectoralis minor.
 - Subclavius.
 - Serratus anterior.
- Describe and demonstrate the boundaries and contents of the **axilla**.
- Describe the formation of the **brachial plexus** and its branches.

Pectoralis Major



- **Origin** : 2 heads;
- **Clavicular head**: From;
 - Medial ½ of the front of the clavicle.
- **Sternocostal head**: From;
 - **Sternum.**
 - **Upper 6** costal cartilages.
 - Aponeurosis of the **external oblique muscle**.
- **Insertion** :
- Lateral lip of bicipital groove (**Humerus**).
- **Nerve supply** :
- **Medial & lateral pectoral nerves.**
- **Action** :
- **Adduction and medial rotation of the arm.**
- **Clavicular head helps in flexion of arm (shoulder).**

Pectoralis Minor



Origin:

- From 3rd, 4th, & 5th ribs close to their costal cartilages.

Insertion:

- Coracoid process (scapula).


Nerve supply:

- Medial pectoral nerve.

Action:


- Depression of the shoulder.
- Draw the ribs upward and outwards during deep inspiration.

Subclavius




- **Origin:**
 - From 1st rib at its costal cartilage.
- **Insertion:**
 - Subclavian groove in the **middle 1/3** of the inferior surface of **clavicle**.
- **Nerve supply:**
 - Nerve to subclavius from upper trunk of brachial plexus.
- **Action:**
 - **Fixes the clavicle** during movement of shoulder joint.

Clavipectoral Fascia




- It is a **thickened** membrane of **deep fascia** **between** the subclavius and pectoralis minor.
- **It is pierced by :**
 - **Lateral pectoral nerve.**
 - **Thoraco- acromial artery**
 - **Cephalic vein.**
 - **Few lymph vessels.**



Origin:


- Upper eight ribs.
- Insertion:
- anterior aspect of the medial border and inferior angle of scapula.
- Nerve supply:
- **Long thoracic nerve** (from roots of brachial plexus,C5,6,7).
- Action:
- **Draws the scapula forward in boxing, (protrusion or protraction). "boxer's muscle"**
- **Rotates scapula outwards in raising the arm above 90 degree (Abduction above 90)**

Serratus anterior



AXILLA


- A **pyramid-shaped space** between the upper part of the arm and the **side of the chest** through which major **neurovascular structures** pass between neck & thorax and upper limbs.
- Axilla has an **apex**, a **base** and **four walls**.



Boundaries of the Axilla


■ Apex:

- Is directed **upwards & medially** to the root of the neck.
- It is called
 - **Cervicoaxillary canal.**
- **It is bounded**, by **3 bones:**
 - **Clavicle anteriorly.**
 - Upper border of the **scapula posteriorly.**
 - Outer border of the **first rib medially.**



- **Base:**


- Formed by **skin** stretching between the anterior and posterior walls.
- **is bounded:**
 - **In front** by the **anterior axillary fold** (formed by the **lower border of Pectoralis major**).
 - **Behind** by the **posterior axillary fold** (formed by tendons of **latissimus dorsi** and **teres major**).
 - **Medially** by **upper 4 to 5 ribs & the chest wall.**




- **Anterior wall:**

- Is formed by :

- Pectoralis major
- Pectoralis minor
- Subclavius
- Clavipectoral fascia.



- **Posterior wall:**
- Is formed by:
 - **Subscapularis.**
 - **Latissimus dorsi.**
 - **Teres major**
muscles.




The medial wall:

- It is wide and formed by:
 - Serratus anterior.**
 - Upper 4-5 ribs & Intercostal muscles .**


The lateral wall:

- It is narrow and formed by:
 - Coracobrachialis.**
 - Biceps brachii.**
 - Bicepital groove of the humerus.**



Contents of The Axilla

- **Cords and branches of the brachial plexus**
- **Axillary artery and its branches.**
- **Axillary vein and its tributaries.**
- **Axillary lymph nodes.**
- **Axillary fat.**
- **Loose connective tissue.**



The neurovascular bundle is enclosed in connective tissue sheath, called '**axillary sheath**'


What is a Brachial Plexus ?

Brachial Plexus is a network of nerves that is formed at the root of the neck to enter the upper limb.

Location & Formation

- **Location** : Brachial Plexus is present in the **posterior triangle** of the neck & **axilla**.
- It is **formed by** the union of the anterior Rami of the C 5th, 6th, 7th & 8th and the 1st thoracic spinal nerve.


Roots of C5 & C6 unite to form---- Upper trunk
Root of C7 continuous as the----- Middle trunk
Roots of C8 & T1 unite to form---- Lower trunk



The Plexus can be divided into 5 stages:

- ❖ **Roots:** in the posteriorΔ
- ❖ **Trunks:** in the posteriorΔ
- ❖ **Divisions:** behind the clavicle (in cervico-axillary canal)
- ❖ **Cords:** in the axilla
- ❖ **Branches:** in the axilla
- The first 2 stages lie in the posterior triangle, while the last 2 stages lie in the axilla. 16

Note: Usual composition shown.
Prefixed plexus has large C4 contribution but lacks T1.
Postfixed plexus lacks C5 but has T2 contribution




- ❖ The **anterior divisions** of the **upper and middle trunks** unite to form the **Lateral cord**.
- ❖ The **anterior division** of the **lower trunk** continues as the **Medial cord**.
- ❖ All the posterior divisions of **three trunks** join to form the **Posterior cord**.

B R N C H E S

Lateral cord-3	Medial cord-5	Posterior cord-5
Lateral pectoral nerve.	Medial pectoral nerve.	Axillary nerve.
Musculocutaneous nerve.	Ulnar nerve.	Radial nerve.
Median nerve (lateral root).	Median nerve (medial root).	Upper & lower subscapular nerves.
Medial cutaneous nerve of arm & forearm.		Thoracodorsal (N. to latissimus dorsi).


Note: Usual composition shown.
Prefixed plexus has large C4 contribution but lacks T1.
Postfixed plexus lacks C5 but has T2 contribution



Applied Anatomy

- Lesion of long thoracic nerve (C5,6,7) causes backward projection of medial border of scapula.
- This deformity is called Winging of Scapula.

Paralysis of Serratus anterior muscle...
Winging of scapula



Right winged scapula

THANK YOU

SUMMARY

- Muscles of the pectoral region are connecting the upper limb with anterior and lateral thoracic wall:
 - Pectoralis major.
 - Pectoralis minor.
 - Subclavius.
 - Serratus anterior.
- **The axilla** is a pyramidal space situated between the upper part of arm and the side of the chest, it has 4 walls (anterior, posterior, medial and lateral), base, and apex.
- **The axilla** is an important space as it transmits the neurovascular bundle from the neck and thorax to the upper limb.
It contains:
 - Axillary vessels.
 - Cords and branches of the brachial plexus.
 - Axillary lymph nodes.