

Skin and Soft-Tissue Infections

IMPETIGO, ABSCESSSES, CELLULITIS, AND ERYSIPELA

Objectives

1. Describe the anatomical structure of skin and soft tissues.
2. Differentiate the various types of skin and soft tissue infections and their clinical presentation.
3. Name bacteria commonly involved in skin and soft tissue infections
4. Describe the pathogenesis of various types of skin and soft tissue infections
5. Recognize specimens that are acceptable and unacceptable for different types of skin and soft tissue infections
6. Describe the microscopic and colony morphology and the results of differentiating bacteria isolates in addition to other non-microbiological investigation
7. Discuss antimicrobial susceptibility testing of anaerobes including methods and antimicrobial agents to be tested.
8. Describe the major approaches to treat of skin and soft tissue infections
 - ▶ either medical or surgical.
 - ▶
 - ▶




FIGURE 1. Depth of involvement in skin and soft-tissue infections.

Introduction

- ▶ Common
- ▶ Can be mild to moderate or severe muscle or bone and lungs or heart valves .
- ▶ *Staphylococcus aureus* is the most cause
- ▶ **Emerging antibiotic resistance among**
 - ▶ *Staphylococcus aureus (methicillin resistance)*
 - ▶ *Streptococcus pyogenes (erythromycin resistance)*

key to developing an adequate differential diagnosis requires

- ▶ **History**
 - ▶ patient's immune status, the geographical locale, travel history, recent trauma or surgery, previous antimicrobial therapy, lifestyle, and animal exposure or bites
- ▶ **Physical examination**
 - ▶ severity of infection
- ▶ **Investigation**
 - ▶ CBCs, Chemistry
 - ▶ Swab, biopsy or aspiration
 - ▶ Radiographic procedures
 - ▶ Level of infection and the presence of gas or abscess.
- ▶ **Surgical exploration or debridement**
 - ▶ Diagnostic and therapeutic

IMPETIGO-(Pyoderma)

- ▶ A common skin infection
- ▶ Children **2–5 Yr** in tropical or subtropical regions
- ▶ Nearly always caused by **β -hemolytic streptococci** and/or ***S.aureus***.
- ▶ Nonbullous (Streptococcus) or Bullous (***S. aureus***)
- ▶ Consists of discrete purulent lesions
- ▶ Exposed areas of the body(face and extremities)
- ▶ Skin colonization- Inoculation by abrasions, minor trauma, or insect bites
- ▶ Systemic symptoms are usually absent.
- ▶ **Poststreptococcal glomerulonephritis.**
- ▶ **(anti-DNAse B)**
- ▶ **Cefazolin, Cloxacillin , or erythromycin**
- ▶ Mupirocin



▶ ABSCESSSES, CELLULITIS, AND ERYSIPELA

▶ Cutaneous abscesses.

- ▶ Collections of pus within the dermis and deeper skin tissues.
- ▶ Painful, tender, and fluctuant
- ▶ Typically polymicrobial,, *S. aureus* alone in ~ 25 %
- ▶ Do Gram stain,culture, and systemic antibiotics
 - Multiple lesions, cutaneous gangrene, severely impaired host defenses, extensive surrounding cellulitis or high fever.
 - Incision and evacuation of the pus



▶ **Furuncles and carbuncles.**

- ▶ **Furuncles** (or “boils”) are infections of the hair follicle (**folliculitis**), usually caused by *S. aureus*, in which suppuration extends through the dermis into the subcutaneous tissue
- ▶ **Carbuncle**- extension to involve several adjacent follicles with coalescent inflammatory mass - back of the neck especially in diabetics
- ▶ Larger furuncles and all carbuncles require incision and drainage.
- ▶ Systemic antibiotics are usually unnecessary

Outbreaks of furunculosis caused by MSSA, and MRSA,

- ▶ Families–prisons–sports teams
- ▶ Inadequate personal hygiene
- ▶ Repeated attacks of furunculosis
- ▶ Presence of *S. aureus* in the anterior nares– 20–40%
- ▶ Mupirocin ointment– eradicate staphylococcal carriage nasal colonization



▶ Erysipelas and Cellulitis.

- ▶ Diffuse spreading skin infections, excluding infections associated with underlying suppurative foci
- ▶ Most of the infections arise from streptococci, often group A, but also from other groups, such as B, C, or G.

▶ Erysipelas

- ▶ Affects the upper dermis (raised-clear line of demarcation)
- ▶ Red, tender, painful plaque
- ▶ Infants, young children-
- ▶ β -hemolytic streptococci (group A or *S. pyogenes*)
- ▶ Penicillin-IV or oral.



▶ Cellulitis

- ▶ Acute spreading infection involves the deeper dermis and subcutaneous tissues.
 - ▶ **β -hemolytic streptococci**, Group A streptococci, and group B streptococci-**diabetics**
 - ▶ ***S. aureus*** : commonly causes cellulitis- penetrating trauma.
 - ▶ ***Haemophilus influenzae*** periorbital cellulitis in children
 - ▶ Risk factors ; Obesity, venous insufficiency, lymphatic obstruction (operations), preexisting skin infections- ulceration, or eczema,
 - ▶ CA-MRSA
 - Carry Panton-Valentine leukocidin gene
 - More sensitive to antibiotics
 - Can lead to sever skin and soft tissue infection or septic shock



Diagnosis and Treatment

- ▶ Clinical diagnosis Symptoms and Signs
- ▶ High WBCs, blood culture rarely needed
- ▶ Aspiration and biopsy might be needed in diabetes mellitus, malignancy, animal bites, neutropenia (*Pseudomonas aeruginosa*),immunodeficiency, obesity and renal failure
- ▶ Observe for progression to sever infection(increased in size with systemic manifestation ie . fever, leukocytosis)
- ▶ Treatment: cover streptococcus and staphylococcus
- ▶ Penicillin, cloxacillin, cefazolin(cephalexin),clindamycin
- ▶ Vancomycin or linazolid in case of MRSA
- ▶ Clindamycin, TMP-SMZ for **CaMRSA**




Necrotizing fasciitis

flesh-eating disease

Introduction

- ▶ It is a rare deep skin and subcutaneous tissues infection
- ▶ It can be monomicrobial (Type II) or (polymicrobial Type I) infection
- ▶ Most common in the arms, legs, and abdominal wall and is fatal in 30%-40% of cases.
- ▶ Fournier's gangrene (testicular), Necrotizing cellulitis
- ▶ Group A streptococcus (*Streptococcus pyogenes*)
- ▶ *Staphylococcus aureus* or CA-MRSA
- ▶ *Clostridium perfringens* (gas in tissues)
- ▶ *Bacteroides fragilis*
- ▶ *Vibrio vulnificus* (liver function)
- ▶ *Gram-negative bacteria* (synergy).
 - ▶ *E. coli*, *Klebsiella*, *Pseudomonas*
- ▶ Fungi



Risk factors


- ▶ Immune-suppression
- ▶ Chronic diseases: (diabetes, liver and kidney diseases, malignancy
- ▶ Trauma:(laceration, cut, abrasion, contusion, burn, bite, subcutaneous injection, operative incision)
- ▶ Recent viral infection rash (chickenpox)
- ▶ Steroids
- ▶ Alcoholism
- ▶ Malnutrition
- ▶ Idiopathic

Pathophysiology

- ▶ destruction of skin and muscle by releasing toxins
 - ▶ Streptococcal pyogenic exotoxins
 - ▶ Superantigen
 - ▶ Non-specific activation of T-cells
 - ▶ Overproduction of cytokines
 - ▶ Severe systemic illness (Toxic shock syndrome)



Signs and symptoms

- ▶ Rapid progression of severe pain with fever , chills (typical)
- ▶ Swelling , redness, hotness, blister, gas formation, gangrene and necrosis
- ▶ Blisters with subsequent necrosis , necrotic eschars
- ▶ Diarrhea and vomiting (very ill)
- ▶ Shock organ failure
- ▶ Mortality as high as 73 % if untreated



Diagnosis

- ▶ A delay in diagnosis is associated with a grave prognosis and increased mortality
- ▶ Clinical-high index of suspicion
- ▶ Blood tests
 - ▶ CBC-WBC , differential , ESR
 - ▶ BUN (blood urea nitrogen)
- ▶ Surgery debridement- amputation
- ▶ Radiographic studies
 - ▶ X-rays : subcutaneous gases
 - ▶ Doppler CT or MRI
- ▶ Microbiology
 - ▶ Culture & Gram's stain
 - ▶ (blood, tissue, pus aspirate)
 - ▶ Susceptibility tests



ASM MicrobeLibrary.org © Delišić
Streptococci © Gloria Delišić, author. Licensed for use,
ASM MicrobeLibrary (linked to
<http://www.microbelibrary.org>)

Treatment

- ▶ If clinically suspected patient needs to be hospitalized OR require admission to ICU
- ▶ Start intravenous antibiotics immediately
- ▶ Antibiotic selection based on bacteria suspected
- ▶ broad spectrum antibiotic combinations against
 - ▶ methicillin-resistant *Staphylococcus aureus* (MRSA)
 - ▶ anaerobic bacteria
 - ▶ Gram-negative and gram-positive bacilli
- ▶ **Surgeon consultation**
 - ▶ Extensive Debridement of necrotic tissue and collection of tissue samples
 - ▶ Can reduce morbidity and mortality

Treatment

- ▶ Antibiotics combinations
 - ▶ Penicillin-clindamycin-gentamicin
 - ▶ Ampicillin/sulbactam
 - ▶ Cefazolin plus metronidazol
 - ▶ Piperacillin/tazobactam
 - ▶ *Clostridium perfringens* - penicillin G
- ▶ Hyperbaric oxygen therapy (HBO) treatment

Pyomyositis

- Acute bacterial infection of skeletal muscle, usually caused by *Staph. aureus*
- No predisposing penetrating wound, vascular insufficiency, or contiguous infection
- Most cases occur in the tropics
- 60% of cases outside of tropics have predisposing RF: DM, EtOH liver disease, steroid rx, HIV, hematologic malignancy

Pyomyositis

- Hx of blunt trauma or vigorous exercise (50%), then period of swelling without pain. 10-21 days later, pain, tenderness, swelling and fever, Pus can be aspirated from muscle. 3rd stage: sepsis, later metastatic abscesses if untreated
- Dx: X-ray, US, MRI or CT
- Rx: surgical drainage +abx

Other Specific Skin Infections

Epidemiology	Common Pathgen(s)	Therapy
Cat/Dog Bites	Pasturella multocida; Capnocytophaga	Amox/clav (Doxy; FQ or SXT + Clinda)
Human bites	Mixed flora eikenella corrodens	Hand Surgeon; ATB as above
Fresh water injury	Aeromonas	FQ; Broad Spectrum Beta-lactam
Salt water injury (warm)	Vibrio vulnificus	FQ; Ceftazidime
Thorn , Moss	sporothrix schenckii	Potassium iodine
Meat-packing	Erysipelothrix	Penicillin
Cotton sorters	Anthrax	Penicillin
Cat scratch	Bartonella	Azithromycin

TAKE HOME POINTS

- ▶ Most commonly caused by *Staphylococcus aureus* and *Streptococcus pyogenes*
- ▶ Risk factors for developing SSTIs include breakdown of the epidermis, surgical procedures, crowding, comorbidities, venous stasis, lymphedema
- ▶ Most of the infections are mild and can be managed on an outpatient basis
- ▶ In case

TAKE HOME POINTS

- ▶ • Most SSTIs can be managed on an outpatient basis, although patients with evidence of rapidly progressive infection, high fevers, or other signs of systemic inflammatory response should be monitored in the hospital setting.
- ▶ • Superficial SSTIs typically do not require systemic antibiotic treatment and can be managed with topical antibiotic agents, heat packs, or incision and drainage.
- ▶ • Systemic antibiotic agents that provide coverage for both *Staphylococcus aureus* and *Streptococcus pyogenes* are most commonly used as empiric therapy for both uncomplicated and complicated deeper infections.