



The excitable tissues (Nerve+ Muscle)

TEXTBOOK OF MEDICAL
PHYSIOLOGY

GUYTON & HALL 13TH
EDITION

UNIT II CHAPTER 5




Objectives

At the end of this lecture the student should be able to:

- Discuss the resting membrane potential and its genesis.
- Know the ionic channels involved in resting membrane potential.
- Describe the function $\text{Na}^+\text{-K}^+$ pump and the stages of action potential.
- Explain the threshold Potential, local Response and action Potentials.
- Describe the electrical changes in membrane potential during the action potential, their chemical bases and excitability changes.
- Describe conduction along nerve fibers, role of myelination and how nerve fibers are classified.



The nerve



Neuron:-

DIF: __ unit of function of the central nervous system


Parts of motor neuron & function of each part:



1- Soma (cell body)

2-Dendrites carry nerve impulses from surroundings to the soma

3 Axon hillock

4-Axon & axon terminal






-Histological classification of axons:-

1- myelinated : have myelin sheath (diameter more than 1um)

2- unmyelinated (diameter less than 1um)

-type C : postganglionic autonomic & pain fibers




-Myelin sheath is formed by schwann cell which deposit sphingomyelin -

Functions of myelin sheath

1-insulator

3- increase conduction velocity





The resting membrane potential of nerves




RESTING MEMBRANE POTENTIAL

DIF: it is potential difference across membrane during rest (without stimulation)


Value:- -70 to -90 mv in large nerve fibers (-ve inside)



-The membrane is polarized



Causes of RMP:



- 1. RMP is 100 times more permeable to K^+ than Na^+ . K^+ tends to leak out of the cell down its conc gradient, carrying +ve charge with it. (through K leak channels).
- 2. non-diffusible anions (proteins, sulphate and phosphate ions) cannot leave the cell.
- 3. very small amount of Na^+ diffuses into the cell down its conc gradient. The mb only slightly permeable to Na^+ . (through Na^+ leak channels).
- 4. Na^+-K^+ pump maintain conc gradients of K^+ , and Na^+ between the two sides of the mb.





What does it mean when a neuron “fires”?

- Firing = excitability = action potential = nerve impulse
- Recall resting potential of all cells
 - High K^+ in; high Na^+ out
 - Cell is **polarized**
 - Cell overall neg. charge inside due to molecules like proteins, RNA, DNA
 - Charge measured in millivolts
 - **Potential** = difference in charge across PM
 - **Current** = flow of charge (ions) from one point to another




Changes that occur through the nerve after stimulation by threshold (effective) stimulus:-

1- Electrical changes (nerve action potential)



2- Excitability changes

3- Thermal changes

4- Chemical changes




Nerve physiology: Action potentials



The action potential

- It is sudden reversal of membrane polarity produced by a stimulus to produce a physiological effect such as:
- Transmission of impulse along nerve fibres
- Release of neurotransmitters
- Muscle contraction
- Activation or inhibition of glandular secretion



1- Electrical changes

The nerve action potential

-It is potential difference along nerve membrane after stimulation

by threshold (effective)stimulus

- oscilloscope to measure rapid changes in membrane potential

-Nerve signals (impulses) are transmitted as nerve action potentials conducted along the nerve fiber as a wave of depolarization to its end

-The factors necessary for nerve action potential are voltage gated Na &

Voltage gated k channels

Threshold stimulus








Figure 7.13 Depolarization of an axon affects Na⁺ and K⁺ diffusion in sequence. (1) Na⁺ gates open and Na⁺ diffuses into the cell. (2) After a brief



Reversal Potential
= + 35 mV



Threshold Potential (Firing Level)
= -50 to -65 mV



RMP = -90 mV

Q : What opens the voltage-gated channels ? Opened by a stimulus strong enough to depolarize them to threshold


Increasing Stimulation





Depolarization




c-Repolarization :- due to high K conductance(flow) to outside (K outflux) by opening of all voltage gated K channels (causes negativity inside)



Repolarization




- **Hyperpolarization: Why?**
- Na-K pump now start to move Na out & K in against their concentration gradient, so the RMP is resumed and the membrane is ready for another stimulus



The action potential (cont.)***


Threshold stimulus:

If a stimulus is strong enough to move **RMP** from its resting value (**-70mV**) to the level of (**-55mV**) which leads to production of an **AP**




Subthreshold stimulus:

Stimulus that result only in local depolarisation




All or nothing principle:-

- Once threshold value for excitation is reached a full AP produced ,its intensity can not increased by increasing stimulus intensity (suprathreshold)





What happens after an action potential?


- Refractory period: few milliseconds
 - Time during which can't stimulate neuron a second time
 - Happens until recovery of resting potential
- Two stages
 - **Absolute refractory period**
 - No new action potential possible
 - **Relative refractory period**
 - Can trigger new action potential if stimulus is very strong



Activation-Inactivation-Deactivation



The Na⁺ Voltage-Gated Channel (1)





- Has 2 gates : one on the outer side of the membrane and is called the activation gate ,
- and another one on the inner side of membrane called the inactivation gate .
- And this channel has 3 states :

- (1) Resting state : in the resting cell , when the MP = RMP = -70 to -90 mV , →
- the activation gate is **closed**
- this prevents entry of Na⁺ to the interior of the cell through this gate.




Activated State of Sodium Channel

- (2) Activated state : when a Threshold Depolarizing Stimulus moves the MP from its resting value (-90 mV) to its Threshold value (-65 to -55mV)
- → this opens the activation gate , and now the Na⁺ channel is said to be in the **Activated State**
- (NB in this case BOTH the activation gate & inactivation gate are open) →
- permeability to Na⁺ becomes increased 500 to 5000 times → Na⁺ influx
- Na⁺ flows into the cell in large amounts ,



Inactivated State of Sodium Channel

- (3) Inactivated state : A few milliseconds after the activation gate opens , the channel becomes inactivated :




At the peak of AP the inactivation gate will close

•the inactivation gate will not open by a second stimulus → & the cell becomes Refractory (ممانعة) to another stimulation .



•This goes on until the MP has gone back to its resting (RMP) level (-70 to -90mV).

- → in this case , while the activation gate is still open , the inactivation gate is closed .





The Potassium Voltage-Gated Channel

- Has one gate only .
- During the resting state , the gate of the potassium channel is closed , and K^+ can not enter through it .
- Shortly after depolarization , when the sodium channel begins to be inactivated , the potassium channel opens .
- → K^+ exits (called K^+ Efflux) خروج البوتاسيوم
- → Repolarization



Direction of AP Propagation (Conduction)



Artificial Electrical



- ✓ Under Artificial condition of electrical stimulation in the laboratory , the AP propagates in both directions .
- ✓ But normally AP starts in axon hillock & propagates distally in one directions



Nerve physiology: Action potentials




Propagation of action potential

1- in myelinated nerve fibers:- Saltatory conduction (jumping)



Value:-



- 1-↑ velocity of conduction of nerve impulses
- 2- Conserve energy for axon because only nodes depolarize




How do action potentials travel down the axon?

- Myelinated sheaths
 - Many times faster transmission
 - Action potential skips from one node of Ranvier to the next
 - Called **saltatory conduction**
 - <http://www.blackwellpublishing.com/matthews/actionp.html>






The structure of two kinds of neurons. (a) A motor neuron and (b) a sensory neuron are depicted here.



2- Non- myelinated nerves:-

(local circuits)=point to point

-depolarization pass by local circuits. -




What else influences speed of action potential?



.Axon diameter

-The larger the diameter, the faster the speed of transmission

-Less resistance to current flow with larger diameter



Faster
transduction





Slower transduction



What happens if myelination is lost?

- Multiple sclerosis
 - Autoimmune disease
 - Usually young adults
 - Blindness, problems controlling muscles
 - Ultimately paralysis
 - Immune system attacks myelin sheaths and nerve fibers
 - Scar tissue (scleroses) replaces some damaged cells



• Thank You

