

Hypoxia and cyanosis

Dr. Laila Al-Dokhi

Assistant Professor

Physiology Department

Objectives

- By the end of this lecture you should be able to:
- Define hypoxia and list its various physiological and pathological causes.
- Define hypo and hyper-ventilation in terms of arterial PCO₂ and PO₂.
- Define cyanosis and its clinical presentation
- Define ventilation/perfusion (V_f/Q) ratio and its normal values.


Hypoxia

Is defined as deficiency of oxygen in the tissue cells.

It can be classified into the following groups:-

- Hypoxic or arterial hypoxia
- Anemic hypoxia
- Stagnant hypoxia
- Histiotoxic hypoxia

Hypoxic Conditions



I-Hypoxic or arterial hypoxia

Reduced arterial PO₂ it can be due to

- Alveolar hypoventilation
- Diffusion abnormalities
- Right to left shunt
- Ventilation-perfusion imbalance
(including increased physiological dead space and physiological shunt).

II-Anemic hypoxia

- It is caused by reduction in the oxygen carrying capacity of the blood, due to decreased amount of Hb or abnormal type of Hb which is unable to carry oxygen.
- The PO₂ and % Hb-O₂ is normal.
- Causes:
 - 1- Anemia
 - 2- Abnormal Hb e.g met hemoglobin, carboxyhemoglobin

III-Stagnant hypoxia:

- Caused by reduced blood flow through the tissues,
so more and more oxygen is extracted from the blood, and due to slow circulation less oxygen is carried by the blood at the lung , leading to hypoxia.
- Causes:
- 1-General slowing of the circulation, as in heart failure and shock
- 2-Local slowing e.g vasoconstriction, cold, arterial wall spasm.

IV- Histiotoxic hypoxia

- This is inability of the tissues to use oxygen due to inhibition of the oxidative enzyme activity
- e.g cyanide poisoning causing blockade of the cytochrome oxidase activity

Effects of hypoxia

- According to the degree of hypoxia it could lead to:
 - Impairment of judgment,
 - Inability to perform complex calculations,
 - Headache, nausea, irritability, dyspnea, increased heart rate,
 - Reduction in muscle working capacity,
 - Coma and death may result.

Treatment of hypoxia

- Is by giving oxygen therapy in a tent or high oxygen tension mask.
- This is useful in hypoxic hypoxia, but of less value in other types of hypoxia .
- Histiotoxic hypoxia will not benefit from O₂ therapy.



Hypercapnea

- Excess of CO₂ in body fluids, it usually occurs with hypoxia, PCO₂ increases above 52 mmHg, it decreases the PH
- **Features of hypercapnea**
 - Peripheral vasodilatation
 - Sweating
 - Warm extremities and bounding pulse
 - Muscle twitching
 - Headache, drowsiness and com
 - Papilledema (swelling of optic disc)

Cyanosis



Cyanosis

- Blue discoloration of the skin and mucus membrane due to more than 5 g/dl of reduced (deoxygenated) hemoglobin in blood.
- A person with anemia almost never develop cyanosis due to low amount of Hb for 5 grams to be deoxygenated /100ml blood.

Ventilation –perfusion ratio (V/Q)

- It is the ratio of alveolar ventilation to pulmonary blood flow per minute.

The alveolar ventilation at rest (4.2 L/min)


- The pulmonary blood flow is equal to right ventricular output per minute (5L/min)

$$\text{SO} \quad V/Q \text{ ratio} = \frac{4.2}{5} = 0.84$$

Cont... V/Q ratio

- Average V/Q ratio across the lung is 0.8.
 - At the apex V/Q ratio = 3
 - At the base V/Q ratio=0.6
- So the apex is more ventilated than perfused and the base is more perfused than ventilated.
- During exercise the V/Q ratio becomes more homogenous among different parts of the lung


Ventilation/perfusion abnormalities



Cont... V/Q ratio

- The main function of this ratio is to determine the state of oxygenation in the body.
- Any mismatch in the ratio can result in hypoxia.
- When the V/Q ratio is less than normal this is called physiologic shunt (a certain fraction of the venous blood is passing through the pulmonary capillaries without being oxygenated i.e shunted blood).
- When V/Q is more than normal this is called Physiologic dead space (when the ventilation of some of the alveoli is great but the alveolar blood flow is low, ventilation of these alveoli is wasted).

Ventilation-Perfusion Lung Scan



Abnormalities of the V/Q ratio

- In the Upper and Lower regions of the normal lung
Apex V/Q ratio = 3 (moderate degree of physiologic dead space)
Base V/Q ratio = 0.6 (represent a physiologic shunt).
- In Chronic Obstructive Lung disease COPD.
because of bronchial obstruction in some areas and destruction of the alveolar septa in other areas with patent alveoli those people has some areas of the lung exhibit serious physiologic shunt and other areas serious physiologic dead space.
COPD is the most prevalent cause of pulmonary disability today, lung effectiveness as a gas exchange organ may decrease to 10%