

HYPERTENSION

Sufia Husain
Pathology Department
KSU, Riyadh
March 2018
Reference: Robbins & Cotran Pathology and
Rubin's Pathology



Objectives:

At the end of the lecture, the student should:


- Know the etiology, risk factors and complications of hypertension, so as to be able to identify patient risk factors amenable to treatment by lifestyle modification, and to investigate patients appropriately for causes of secondary hypertension.

Key principles to be discussed:

- Raised systemic blood pressure is a major cause of morbidity and mortality.
 - Hypertension can cause or contribute to: atherosclerosis, left ventricular hypertrophy, chronic renal failure, cerebrovascular disease and retinopathy.
 - Normal values for blood pressure.
 - Causes of secondary hypertension.
 - Genetic and environmental factors contributing to the aetiology of essential hypertension.
 - Pathology of blood vessels (blood vessels changes) in both primary and secondary hypertension.
-

LECTURE OUTLINE

- Definition and risk factors
 - Classification
 - Primary & secondary HTN
 - Causes of secondary HTN
 - Benign vs malignant HTN
 - Pathogenesis
 - Regulation of blood pressure
 - Vascular morphology in HTN
 - Heart in HTN
 - Complications of HTN
-



Hypertension & Hypertensive Vascular Disease

- Common problem
 - Asymptomatic until late - Silent Killer – painless.
 - Complications alert to diagnosis but late.
 - **Hypertension:** *Definition:* a sustained systolic pressure in excess of 140 mm Hg or a sustained diastolic pressure more than 90 mm Hg (>140/90)
 - In the early stages of HTN there are few or no symptoms.
 - Hypertension is an important factor which contributes in development of:
 - Coronary heart disease.
 - Cerebrovascular accidents (stroke)
 - Cardiac hypertrophy
 - Congestive heart failure.
 - Aortic dissection.
 - Renal failure.
 - Retinopathy
-

Risk factors for Hypertension

- **Hereditary, Genetics-** family history
 - **Race.** African-Americans
 - **Gender.** Men & postmenopausal women
 - **Age**
 - **Obesity**
 - **Diet, particularly sodium intake**
 - **Lifestyle-stressful**
 - **Heavy alcohol consumption**
 - **Diabetes**
 - **Use of oral contraceptives**
 - **Sedentary or inactive lifestyle**
-

HTN can be classified in 2 ways:

1. Based on etiology or cause
 2. Based on clinical features
-

Classification: based on etiology/cause

- I. **Primary/Essential Hypertension (95%)** : Mechanisms largely unknown. It is idiopathic.
- II. **Secondary Hypertension (5-10%)**: it can be due to pathology in the renal, endocrine, vascular or neurogenic systems



Causes of Secondary Hypertension

| | |
|-------------------|---|
| Renal | Glomerulonephritis, Renal artery stenosis, Renal vasculitis Adult polycystic disease Chronic renal disease, Renin producing tumors |
| Endocrine | Adrenocortical hyperfunction (Cushing syndrome, primary aldosteronism, congenital adrenal hyperplasia) Hyperthyroidism/Thyrotoxicosis Hypothyroidism/Myxedema, Pheochromocytoma Acromegaly Exogenous hormones (glucocorticoids, estrogen e.g. oral contraceptives) Pregnancy-induced |
| Vascular | Coarctation of aorta Vasculitis e.g. Polyarteritis nodosa Increased intravascular volume Increased cardiac output Rigidity of the aorta |
| Neurogenic | Psychogenic Increased intracranial pressure Sleep apnea Acute stress, including surgery |

Classification based on clinical features.

○ Benign:

- The BP is at modest level (not very high).
- It can be idiopathic HTN or secondary HTN
- Fairly stable over years to decades.
- Compatible with long life.

○ Malignant(5%):

- there is rapidly rising BP which often leads to end organ damage
 - It can be a complication of any type of HTN (i.e. essential or secondary)
 - It is seen in 5% of HTNsive patients.
 - The diastolic pressure is usually over 120mmHg
 - It is associated with:
 - Widespread arterial necrosis and thrombosis
 - Rapid development of renal failure
 - Retinal hemorrhage and exudate, with/without papilledema
 - Hypertensive encephalopathy
 - Left ventricular failure
 - Leads to death in 1 or 2 years if untreated.
-

Regulation of Blood Pressure (BP)

There are 2 hemodynamic variables that are involved in the regulation of BP. They are cardiac output and peripheral vascular resistance

$$\text{BP} = \text{Cardiac Output} \times \text{Peripheral Resistance}$$

- **Cardiac output** is affected by blood volume and is dependent on sodium concentrations.
-

Regulation of Blood Pressure (BP)

- **Peripheral resistance:** it is the resistance of the arteries to blood flow. As the arteries constrict, the resistance increases and as they dilate, resistance decreases. Peripheral resistance is regulated at the level of the arterioles (also known as resistance vessels) and is determined by three factors:
 1. Autonomic activity: sympathetic activity constricts peripheral arteries.
 2. Pharmacologic agents: vasoconstrictor drugs increase resistance while vasodilator drugs decrease it.
 3. Blood viscosity: increased viscosity increases resistance.

Normal BP is maintained by a balance between factors that induce vasoconstriction (e.g. angiotensin II and catecholamines) and factors that induce vasodilation (e.g. kinins, prostaglandins, and nitric oxide).

Note: An increased blood flow in the arterioles induces vasoconstriction to protect tissues against hyperperfusion

PATHOGENESIS of Essential Hypertension

Essential HTN occurs when the relationship between cardiac output and peripheral resistance is altered. Multiple **genetic and environmental factors** ultimately **increase the cardiac output and/or peripheral resistance**

(BP = Cardiac Output x Peripheral Resistance)

1. **Genetic factors:** There is a strong genetic component (family history) e.g. a genetic effect is involved in making people more susceptible or less susceptible to high salt diet etc.
 - a) **Defect in renal sodium homeostasis:** *reduced renal sodium excretion* is a key initiating event in most forms of essential hypertension. This decreased sodium excretion will → lead to increase in fluid volume and therefore → increase in cardiac output, thereby → elevated BP. This is usually due to defect in cell membrane function: affecting Na/Ca transport.
 - b) **Functional vasoconstriction:** abnormality in vascular tone such as increased sympathetic stimulation will cause vasoconstriction leading to increased peripheral resistance.
 - c) **Structural abnormality** in vascular smooth muscle also leads to increased peripheral resistance.
-


PATHOGENESIS of Essential Hypertension

- d) **Also rare gene disorders** can cause HTN by increasing renal sodium reabsorption e.g. Liddle syndrome. Liddle syndrome is an inherited autosomal dominant type of HTN, that begins in childhood. It is caused by mutations of the epithelial sodium channel protein (ENaC) which leads to increased sodium reabsorption in the renal tubules (followed by water), which leads to hypertension. Reabsorption of sodium is also correlates with potassium loss (hypokalemia).


2. *Environmental factors:* stress, obesity, smoking, physical inactivity and heavy consumption of salt also play a role.

NOTE: In hypertension, both increased blood volume and increased peripheral resistance contribute to the increased pressure. However reduced renal sodium excretion in the presence of normal arterial pressure (initially) is probably a key initiating event.

PATHOGENESIS of Essential Hypertension




Regulation of Blood Pressure (BP)



© Elsevier. Kumar et al: Robbins Basic Pathology 8e - www.studentconsult.com

ENDOCRINE FACTORS: Renin, Angiotensin, ADH, Aldosterone

ENDOCRINE FACTORS: role of renin- angiotensin-aldosterone in regulating BP



Atrial natriuretic peptide / factor / hormone (Cardionatine / Cardiodilatine / atriopeptin)

- It is a protein (polypeptide) hormone secreted by the heart muscle cells in the atria of heart (atrial myocytes).
 - It is a powerful vasodilator and is involved in the homeostatic balance of body water, sodium, potassium and fat.
 - It is released in response to high blood volume. It acts to reduce the water, sodium and adipose loads on the circulatory system, thereby reducing blood pressure.
 - It has exactly the opposite function of the aldosterone secreted by the zona glomerulosa
 - **In the kidney:**
 - decreases sodium reabsorption and increases water loss.
 - Inhibits renin secretion, thereby inhibiting the renin–angiotensin–aldosterone system
 - **In adrenal gland:**
 - Reduces aldosterone secretion by the zona glomerulosa of the adrenal cortex.
 - **In arterioles:**
 - Promotes vasodilatation
 - **In adipose tissue**
 - Increases the release of free fatty acids from adipose tissue.
-

Morphology of blood vessels in HTN:

In large Blood Vessels (Macroangiopathy)

- Atherosclerosis. HTN is a major risk factor in AS.

In small Blood Vessels (Microangiopathy)

- Arteriolosclerosis


1. Hyaline arteriolosclerosis:

- Seen in benign hypertension
- Can also be seen in elderly and diabetic patients even without hypertension.
- Can cause diffuse renal ischemia which ultimately leads to benign nephrosclerosis


2. Hyperplastic arteriolosclerosis:

- Characteristic of malignant hypertension.
 - Can show onion-skinning on histology causing luminal obliteration of vascular lumen
 - May be associated with necrotizing arteriolitis and fibrinoid necrosis of the blood vessel.
-

Vascular pathology in hypertension




A. Hyaline arteriosclerosis: hyalinosis of arteriolar wall with narrowing of lumen.




B. Hyperplastic arteriosclerosis (onionskinning) causing luminal obliteration of vascular lumen


Vascular Pathology in Hypertension



Hyaline/ Benign hypertension



Hyperplastic/ Malignant hypertension showing onion skinning




Hyperplastic/ Malignant hypertension showing fibrinoid necrosis.


Left ventricular cardiac hypertrophy (also known as left sided hypertensive cardiomyopathy/ hypertensive heart disease)

- Longstanding poorly treated HTN leads to left sided hypertensive heart disease.
 - Hypertrophy of the heart is an adaptive response to pressure overload due to HTN. HTN induces left ventricular pressure overload which leads to hypertrophy of the left ventricle with increase in the weight of the heart and the thickness of the LV wall.
-

Left ventricular cardiac hypertrophy



Left Ventricular Hypertrophy



Complications in HTN:

The organs damaged in HTN are:

- **Cardiovascular**

- Left ventricular cardiac hypertrophy(left sided hypertensive cardiomyopathy/ hypertensive heart disease)
- Coronary heart disease
- Aortic dissection

- **Kidney**


- Benign nephrosclerosis (photo A)
- Renal failure in untreated or in malignant hypertension

- **Eyes**


- Hypertensive retinopathy (photo B) is especially seen in malignant hypertension.

- **Brain**


- Hemorrhage, infarction leading to Cerebrovascular accidents
-



Subarachnoid Haemorrhage




CEREBRAL HEMORRHAGE



CEREBRAL INFARCTION

Lacunar Infarct





© 1997 Jeff Buchino