


TUBULAR PROCESSING OF FILTRATE

Dr. Maha Saja

MBBS, MSc Physiology, PhD

Msaja@ksu.edu.sa

Contents



- The mechanisms of tubular transport through the different parts of the nephron.
- Tubular reabsorption and tubular secretion.
- Regulation of tubular processing.

Objectives


- Define tubular reabsorption and secretion.
- Identify the role of each tubular segment in glomerular filtrate modification and the types of substances being transported through each.
- Describe the hormonal/physiological factors regulating tubular function at each segment.
- Describe tubular reabsorption of sodium and water.
- Identify and describe mechanism involved in glucose reabsorption.
- Identify the tubular site and describe how amino acids and urea are reabsorbed.
- Identify and describe the characteristics of the loop of Henle, distal convoluted tubule and collecting ducts for reabsorption and secretion
- Describe the role of ADH in the reabsorption of water.
- Identify the site and describe the influence of aldosterone on reabsorption of Na⁺.
- List and explain the factors that control aldosterone and ADH release
- Identify and describe the juxtamedullary apparatus and its role in checking the filtrate.

Tubular Processing of Ultrafiltrate


- After glomerular filtration the ultrafiltrate gets modified as it passes through the tubules before it is finally excreted.
- **Tubular processing includes:**
 - ***Tubular reabsorption*** = reabsorption of substances from the glomerular filtrate into peritubular capillary blood.
 - ***Tubular secretion*** = secretion of substances from peritubular capillary blood into tubular fluid
- ***What is the importance of tubular processing?***

Tubular Reabsorption

- **Glomerular filtration and tubular reabsorption are quantitatively very large relative to the amount excreted!**
- **Glomerular filtration is non-selective whereas tubular reabsorption is highly selective.**



Differences in Renal Tubular Cells Reflect Their Function in Tubular Processing



Koeppen & Stanton: Berne and Levy Physiology, 6th Edition.
Copyright © 2008 by Mosby, an imprint of Elsevier, Inc. All rights reserved.

Figure 32-3 Diagram of a nephron, including the cellular ultra-structure.

TUBULAR REABSORPTION

How does the nephron reabsorb substances


- **Reabsorption is a 2 step process:**

1. Transport of substances from tubular lumen to IF.
2. Transport from IF to blood.

- From tubular lumen to IF;
 - Transport involves **active** & **passive** mechanisms.
 - Occur through **paracellular** and/or **transcellular** routes.

- From IF to blood:


- By ultrafiltration (bulk flow).



Transport Mechanisms Across the Tubule

Active Transport

- Requires energy.
- Moves substances against their electrochemical gradient.



Primary active

Directly coupled to energy source.

e.g. Na⁺-K⁺ ATPase.

Secondary active


Indirectly coupled to energy source.

Carrier protein.

e.g. Glucose & a.a.

Passive Transport

- Does not need energy.
- Moves substances down their electrochemical gradient.




Passive diffusion Osmosis

Water
Solute like Cl⁻
Urea

TUBULAR REABSORPTION IN EACH PART OF THE NEPHRON

Proximal Convoluted Tubule

- Most of the reabsorption occurs in the PCT.. **Why?**
 - Highly metabolic cells.
 - Extensive brush border.
 - Lots of mitochondria.



nephron

Sodium Reabsorption

Basolateral $\text{Na}^+\text{-K}^+$ ATPase pumps 3Na^+ out and 2K^+ into the cell




Results in low $[\text{Na}^+]_i$




This gradient favours Na^+ entry across the apical membrane via transporter proteins

(a) Early proximal convoluted tubule



Animal Physiology 2e, Figure 28.16 (Part 1)

Glucose Reabsorption




Transport Maximum for Glucose

What is meant by transport maximum?

Why does it occur?

What happens if blood glucose level increased to 400mg/dl?



Sodium Reabsorption


Early part of PCT

- Mainly coupled to;
 - Glucose.
 - Amino acids.
 - Lactate.
 - Phosphate
 - Hydrogen
- Symporters
- Antipporter

Late part of PCT


- Mainly coupled to Cl⁻
- *Why??*

Sodium Chloride Reabsorption in the 2nd half of the PCT



e.g.
Formate
Oxalate
Sulfate

Water Reabsorption in the PCT



Steps for water and urea reabsorption:

- ① Solute (Na⁺, X, Y) are actively reabsorbed, increasing the osmolarity of peritubular fluid and plasma.
- ② Water is reabsorbed by osmosis.
- ③ Urea (permeating solute) is reabsorbed passively.

Organic Anion/Cation Secretion


Organic Anions

- Endogenous:
 - Bile salts.
 - Oxalate.
 - Urate.
 - Vitamins (ascorbate, folate).
- Exogenous:
 - Acetazolamide.
 - Furosemide.
 - Salicylates.
 - Penicillin.


Organic cations

- Endogenous;
 - Creatinine.
 - Dopamine.
 - Epinephrine.
 - Norepinephrine.
- Exogenous;
 - Atropine.
 - Morphine.
 - Amiloride.
 - Procainamide.

Summary of PCT Transport Mechanisms




Summary of PCT Filtrate Modification



LOOP OF HENLE

Loop of Henle




- Highly permeable to water
- Moderate permeability to solutes

- **Impermeable** to water
- Reabsorption of solutes in the thick segment

Loop of Henle

Mechanism of transport in the thick ascending loop of Henle



Loop diuretics


- Furosemide
- Ethacrynic acid
- Bumetanide

Distal Convoluted Tubule

Resembles the thick ascending loop of Henle

Known as *the diluting segment*

Reabsorbs 5% of NaCl



Early part

Late part

Resembles the cortical collecting tubule

Impermeable to urea

Principal cells

Intercalated cells Type A

Reabsorb Na⁺ & H₂O
Secrete K⁺

Reabsorb K⁺
Secrete H⁺

Controlled by Aldosterone


Acid-base regulation

Permeability to H₂O depends on *ADH*

nephron


Early Distal Tubule

DISTAL CONVOLUTED TUBULE




Thiazide diuretics

Late Distal Tubule & Collecting Tubule






Koepfen & Stanton: Berne and Levy Physiology, 6th Edition.
Copyright © 2008 by Mosby, an imprint of Elsevier, Inc. All rights reserved

LATE DISTAL TUBULE AND COLLECTING DUCT



Medullary Collecting Duct




Medullary Collecting Ducts


Permeability to water is under ADH control

Permeable to urea

Secrete H^+




Summary of the Concentrations of the different Solutes in the Different Tubular Segments



Regulation of Tubular Reabsorption


- ***Regulation of tubular reabsorption depends on:***
 1. Physical forces that govern reabsorption.
 2. Hormonal and neural mechanisms.
- Tubules can increase their reabsorption in response to increased tubular load → ***glomerulotubular balance.***
- ***What are the physical forces that govern tubular reabsorption?***

Physical Forces that Govern Tubular Reabsorption




Pressure Natriuresis & Pressure Diuresis

- = increasing urinary excretion of Na^+ & H_2O in response to increases in ABP.
- Autoregulation should limit this! *What happens if autoregulation is impaired?*



Hormonal Regulation of Tubular Reabsorption



REGULATION OF POTASSIUM

Body Potassium Balance

- **Input:**

- Dietary intake = 80-120 mmol/day.
- The ECF K⁺ content ≈ 70 mmol.

- **Output:**

- GI loss ≈ 5-10%
- Renal excretion ≈ 90-95%.

***Control External
balance***

Between body and
environment

- ***Internal balance*** (within the body) is regulated by modifying the distribution of K⁺ between the ICF & ECF.

Body Potassium Balance




Figure 29-1


Normal potassium intake, distribution of potassium in the body fluids, and potassium output from the body.

The Importance of Regulating K⁺

Table 37-1 Physiological Role of K⁺ Ions

A. Roles of Intracellular K⁺	
Cell-volume maintenance	Net loss of K ⁺ → cell shrinkage Net gain of K ⁺ → cell swelling
Intracellular pH regulation	Net loss of K ⁺ → cell acidosis Net gain of K ⁺ → cell alkalosis
Cell enzyme functions	K ⁺ dependence of enzymes (e.g., some ATPases, succinic dehydrogenase)
DNA/protein synthesis, growth	Lack of K ⁺ → reduction of protein synthesis, stunted growth
B. Roles of Transmembrane [K⁺] Ratio	
Resting cell membrane potential	Reduced [K ⁺] _i /[K ⁺] _o → membrane depolarization Increased [K ⁺] _i /[K ⁺] _o → membrane hyperpolarization
Neuromuscular activity	Low plasma K ⁺ : muscle weakness, muscle paralysis, intestinal distention, respiratory failure High plasma K ⁺ : increased muscle excitability; later, muscle weakness (paralysis)
Cardiac activity	Low plasma K ⁺ : slowed conduction of pacemaker activity, arrhythmias High plasma K ⁺ : conduction disturbances, ventricular arrhythmias, and ventricular fibrillation

Renal Potassium Handling




- $\approx 90\%$ of K^+ is reabsorbed in the PCT & TAL.
- $\approx 10\%$ enters the distal portions of the nephron.
 - **If K^+ intake is low** only 1-3% of filtered K^+ will be excreted.
 - **If K^+ intake is normal/high**, 10-15% of filtered K^+ will be excreted

Potassium Handling by the kidney

It is the sum of filtration – reabsorption + secretion

In the PCT → K^+
reabsorption is a passive
process.. **How?**


Water reabsorption
through the paracellular
route drags K^+ with it
(**solvent drag**).



Potassium Handling by the TAL

By secondary active transport using the apical triple transporter (NKCC2).


B THICK ASCENDING LIMB (TAL)



Potassium Handling by the CT

Alpha-Intercalated cells


C CORTICAL COLLECTING TUBULE (CCT):
 α INTERCALATED CELL



Secrete H^+ and reabsorb K^+

Principal cells

D CORTICAL COLLECTING TUBULE (CCT):
PRINCIPAL CELL



Reabsorb Na^+ and water &
secrete K^+

Factors affecting Potassium

- Aldosterone.
- Insulin
- Epinephrine
- ECF pH
- Luminal flow (diuresis).

THANK YOU
