


ACID-BASE BALANCE

Dr. Maha Saja

MBBS, MSc Physiology, PhD

Contents



- What are acids and bases?
- What is meant by a weak and a strong acid or base?
-
- What is the normal pH of body fluids?
- Why is it important to keep body fluid pH within certain limits?
- What are the body's defense mechanisms against changes in blood pH: body buffers, the lungs and the kidney.
- Understand the role of the kidney in regulating pH of body fluids.
- Acid-base disturbances.

Acid-Base Balance

- Acid-base balance is concerned with the precise regulation of free (unbound) hydrogen ion (H^+) concentration in body fluids.
- Normally, $[H^+] = 0.00004 \text{ mEq/L}$ (40 nEq/L).
- ***Why is it important to control $[H^+]$?***

Why Should $[H^+]$ be Tightly Controlled?

- Slight deviations in $[H^+]$ have profound effects on enzyme and protein activity and thus the body's metabolic activity in general.
- Changes in $[H^+]$ affects K^+ levels in the body.

Why is the Body's $[H^+]$ Constantly Changing?


A number of processes can alter $[H^+]$ concentration in the body, such as;

1. Metabolism of ingested food.
2. GI secretions.
3. Generation of acids & bases from amino acid/protein metabolism.
4. Changes in CO_2 production.

Acid-Base Fundamentals

- **An Acid** = a molecule that can release H^+ in a solution.
 - H_2CO_3 (carbonic acid)
 - HCl (hydrochloric acid)
- **A base** = a molecule that accepts H^+ in a solution.
 - Bicarbonate ions (HCO_3^-).
 - Hydrogen phosphate (HPO_4^{-2})
- ***What is the difference between carbonic & hydrochloric acid?***

Strong vs Weak Acids & Bases



Strong acids dissociate rapidly and release large amounts of H^+ in solution

Weak acids dissociate incompletely and less strongly releasing small amounts of H^+ in solution

Weak Acids



Acid \leftrightarrow *Conjugate base* + H^{+}

The extent to which a given acid dissociates in solution is constant. And is known as the **dissociation constant (K)**.

$$K = \frac{[H^{+}][A^{-}]}{[AH]}$$




[H⁺] & the pH

- H⁺ ion concentrations are expressed as pH.
- $\text{pH} = -\text{Log} [\text{H}^+]$
 - If the [H⁺] increase → pH will decrease (more acidic)
 - If the [H⁺] decrease → pH will increase (more alkaline)


- ***What is the normal pH of the ECF?***

Normally pH= 7.35-7.45



The Body and pH

pH of arterial blood



pH and H⁺ Concentration of Body Fluids

	[H ⁺] (mEq/L)	pH
Extracellular fluid		
Arterial blood	4.0 X 10⁻⁵	7.4
Venous blood	4.5 X 10⁻⁵	7.35
IF	4.5 X 10⁻⁵	7.35
Intracellular fluid	1 X 10⁻³ to 4 X 10⁻⁵	6-7.4
Urine	3 X 10⁻² to 1 X 10⁻⁵	4.5-8
Gastric HCl	160	0.8

Acid Production by the Body

- The body produces large amounts of acids on daily basis as by products of metabolism.
 - Metabolism of dietary proteins.
 - Anaerobic metabolism of carbs and fat.
- Acids in the body are of two kinds:
 1. Volatile (CO_2)
 2. Non-volatile “fixed” (sulfuric acid, lactic acid) (daily acid load \approx 50-100 mEq/day) (0.8 mEq/kg/d).

The Body's Defense Against Changes in $[H^+]$

Three main systems:

1. *Body fluid buffers.*

Works within seconds.

2. *Lungs*

Works within minutes.

3. *Kidneys*


Works within hours-days.

The most powerful of the three.

BODY FLUID BUFFERS

What is a Buffer?

A buffer = a solution that resists changes in pH upon addition of small amount of acids or bases.



How do Buffers work?


- A buffer is a mixture of a weak acid and a weak base that are in equilibrium.
- To be more accurate, its either made of:
 - A weak acid and its conjugated base (H_2CO_3 & NaHCO_3^-).
 - A weak base and its conjugated acid (NH_3 & NH_4^+).
- ***How does a buffer do its job?***

Chemical Buffer Systems in the Body

- ***There are 3 chemical buffers in the body;***
 1. The Bicarbonate buffer system.
 2. The phosphate buffer system.
 3. Proteins.
- They are the 1st line of defence against changes in pH i.e. $[H^+]$, act within seconds.
- Some are more powerful extracellularly and others are more powerful intracellularly.


The Bicarbonate Buffer System

- Composed of:
 - A weak acid (H_2CO_3).
 - Its conjugated base (NaHCO_3).




The Bicarbonate Buffer System


- H_2CO_3 forms in the body by the reaction of CO_2 & H_2O***



- H_2CO_3 ionizes weakly to form small amounts of H^+ & HCO_3^-***




- The second component is $NaHCO_3$ which dissociates to form Na^+ & HCO_3^-***



The Bicarbonate Buffer System

Putting it all together;



The Carbonic Acid-Bicarbonate Buffer System

This is the main ECF buffer system

What happens if you add a base or an acid to the system?

The Henderson-Hasselbalch Equation

What is the HHE?

- It is an equation that enables the calculation of pH of a solution.

What is it?


$$pH = pK + \log \frac{HCO_3^-}{0.03 \times PCO_2}$$

K = dissociation constant, pK = 6.1

0.03 = solubility of CO₂

The Henderson-Hasselbalch Equation

How was it derived?



1. H_2CO_3 and its dissociated ions are always in equilibrium → the products of the reaction on one side of the equation are proportional to the product on the other side.

$$[\text{H}_2\text{CO}_3] \propto [\text{H}^+] \times [\text{HCO}_3^-]$$

2. Since H_2CO_3 is a weak acid, it will not dissociate completely and the concentration of its products will depend on its dissociation constant (K)

$$K \times [\text{H}_2\text{CO}_3] = [\text{H}^+] \times [\text{HCO}_3^-]$$

The Henderson-Hasselbalch Equation

3. Based on the previous equation, $[H^+]$ can be expressed as follows;

$$[H^+] = K \times \frac{[H_2CO_3]}{[HCO_3^-]}$$

4. Because H_2CO_3 can rapidly dissociate into CO_2 and H_2O . And since CO_2 is much easier to measure it can replace H_2CO_3 in the equation;

$$[H^+] = K \times \frac{[CO_2]}{[HCO_3^-]}$$

*This is Henderson's equation
(1908)*

It means that;

$\uparrow [CO_2] \rightarrow \uparrow [H^+]$

$\uparrow [HCO_3^-] \rightarrow \downarrow [H^+]$

The Henderson-Hasselbalch Equation

5. In 1909, Sorensen created the pH scale to express $[H^+]$

$$pH = -\log[H^+]$$

6. In 1916, Hasselbalch decided to merge Henderson's equation with Sorensen's pH scale creating what we now know as the *"Henderson-Hasselbalch equation"*.

$$[H^+] = K \times \frac{[CO_2]}{[HCO_3^-]} \quad \Longrightarrow \quad -\log[H^+] = -\log\left(K \times \frac{[CO_2]}{[HCO_3^-]}\right)$$

The Henderson-Hasselbalch Equation

$$-\log[H^+] = -\log\left(K \times \frac{[CO_2]}{[HCO_3]}\right)$$



$$pH = pK + \log\frac{[HCO_3]}{[CO_2]}$$




*This is Henderson-Hasselbalch equation
(1908)*

7. Since it is much easier to measure PCO_2 rather than dissolved $[CO_2]$ and because dissolved CO_2 is proportional to PCO_2 multiplied by the solubility of CO_2 (0.03 mmol/mmHg) $\rightarrow [CO_2]$ was replaced by $PCO_2 \times 0.03$

$$pH = pK + \log\frac{[HCO_3]}{0.03 \times PCO_2}$$



The Bicarbonate Buffer System



$$\text{pH} = \text{pK} + \log \frac{[\text{HCO}_3^-]}{0.03 \times \text{PCO}_2}$$

• *What do we understand from this equation?*

1. $\text{pH} \propto \frac{\text{HCO}_3^-}{\text{PCO}_2}$

Regulated by kidneys (referring to HCO₃⁻)

Regulated by lungs (referring to PCO₂)

Each element of the buffer system is regulated


- ↑↑ HCO₃⁻ will ↑↑ pH
- ↑↑ PCO₂ will ↓↓ pH

Ratio of $\frac{HCO_3^-}{PCO_2}$ is $\approx 20:1$

Acid-Base Balance

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Figure 24.12



Other Buffering Systems

The phosphate buffer:

- Plays a major role in buffering intracellular & renal tubular fluid.
- Composed of;
 - H_2PO_4^- (dihydrogen phosphate)
 - HPO_4^{2-} (Hydrogen phosphate)

Proteins:

- Contributes to buffering inside cells.
- E.g. Hb.

Summary of Body's Buffering Systems



- Buffer systems do not work independently in body fluids but actually work together.
- A change in the balance in one buffer system, changes the balance of the other systems.
- Buffers do not reverse the pH change, they only limit it.
- Buffers do not correct changes in $[H^+]$ or $[HCO_3^-]$, they only limit the effect of change on body pH until their concentration is properly adjusted by either the lungs or the kidney.

RESPIRATORY REGULATION OF ACID-BASE BALANCE

Respiratory Regulation of A/B

- 2nd line of defence against acid-base disturbances in the body.
- **HOW?**
 - By modulating CO₂ excretion.
- $\uparrow\uparrow [H^+] \rightarrow \uparrow\uparrow \text{ventilation (RR)} \rightarrow \downarrow\downarrow \text{PCO}_2$
- $\downarrow\downarrow [H^+] \rightarrow \downarrow\downarrow \text{ventilation (RR)} \rightarrow \text{accumulation of CO}_2 \rightarrow \uparrow\uparrow \text{PCO}_2$.
- Normally, PCO₂ = 40 mmHg (35-45 mmHg)

Respiratory Regulation of CO₂




RENAL REGULATION OF ACID-BASE BALANCE

Renal Regulation of A/B Balance


- 3rd line of defence against acid-base disturbances and the most powerful.
- It regulates by excreting either an acidic or basic urine.
- **HOW?**
 1. Secreting H⁺
 2. Reabsorbing HCO₃⁻
 3. Generating “new” bicarbonate ions.

Overview HCO_3^- Reabsorption by the Renal Tubules



How is HCO_3^- Reabsorbed by the tubules?


What happens at the PCT?




The PCT reabsorbs, "reclaims", 80-90% of the filtered HCO_3^- .

HCO_3^- reabsorption is linked to H⁺ secretion.

HCO_3^- Reabsorption by the PCT



What happens at the DCT & CT?



What happens at the late DCT & CT?

- The filtrate arriving at the DCT & CT is low in HCO_3^- .
- The distal segments of the nephron are characterised by the presence of “intercalated cells” capable of **actively secreting H^+** through H^+ -ATPase and H^+ - K^+ ATPase present on their apical membrane (**Type-A intercalated cells**).
- Only a limited number of H^+ can be excreted in its free form in urine.
- Lowest possible urine $\text{pH}=4.5 \rightarrow \approx 0.04 \text{ mmol/L}$ of free H^+ .
- **How does the kidney excrete the extra H^+ ?**

Non-Bicarbonate Buffers in the Tubular Lumen?

The extra H⁺ secreted will need to be buffered in the tubular lumen

2 main non-bicarbonate buffers in the tubule

Filtered

Synthesized

Phosphate buffer system




Ammonia buffer system




Excretion of H⁺ and Generation of New HCO₃⁻

The phosphate buffer system



Excretion of H⁺ as phosphate is capable of handling a limited amount of H⁺ and will not be enough to rid the body of its daily acid load nor if there is unusually high acid production.



① CO_2 combines with water within the type A intercalated cell, forming H_2CO_3 .


② H_2CO_3 is quickly split, forming H^+ and bicarbonate ion (HCO_3^-).

③a H^+ is secreted into the filtrate by a H^+ ATPase pump.

③b For each H^+ secreted, a HCO_3^- enters the peritubular capillary blood via an antiport carrier in a HCO_3^- - Cl^- exchange process.

④ Secreted H^+ combines with HPO_4^{2-} in the tubular filtrate, forming H_2PO_4^- .

⑤ The H_2PO_4^- is excreted in the urine.



→ Primary active transport
- - - Secondary active transport
→ Simple diffusion
- - - Facilitated diffusion


● Transport protein
● Ion channel
CA Carbonic anhydrase

Excretion of H^+ and Generation of New HCO_3^-

The Ammonia Buffer System

- Renal tubular cells, especially PCT, are capable of generating ammonium (NH_4^+) “*ammoniogenesis*” which is then excreted in urine carrying with it H^+ .
- The rate of ammoniogenesis can be modified according to the needs of the body.
- Quantitatively, the ammonia buffer system is more important than the phosphate buffer system for H^+ excretion in urine.
- It is the most important system in case of acidosis.


Ammonia buffer system



The Overall Scheme of Renal Excretion of Acids & Bases

To excrete acid:

1. Freely filter HCO_3^-
2. Reabsorb the majority of filtered HCO_3^-
3. Reabsorb some additional HCO_3^-
4. Secrete H^+ (titrate filtered bases, i.e. HPO_4^{-2}) and secrete NH_4^+
5. Excrete acidic urine containing NH_4^+



To excrete base:

1. Freely filter HCO_3^-
2. Reabsorb the majority of filtered HCO_3^-
3. Reabsorb some additional HCO_3^-
4. Secrete some HCO_3^-
5. Excrete alkaline urine containing HCO_3^-

Factors Affecting H^+ Secretion and HCO_3^- Reabsorption

Table 30-2

Factors That Increase or Decrease H^+ Secretion and HCO_3^- Reabsorption by the Renal Tubules

Increase H^+ Secretion and HCO_3^- Reabsorption

↑ PCO_2

↑ H^+ , ↓ HCO_3^-

↓ Extracellular fluid volume

↑ Angiotensin II

↑ Aldosterone

Hypokalemia

Decrease H^+ Secretion and HCO_3^- Reabsorption

↓ PCO_2

↓ H^+ , ↑ HCO_3^-

↑ Extracellular fluid volume

↓ Angiotensin II

↓ Aldosterone

Hyperkalemia

THANK YOU
