

AMINO ACIDS

(Foundation Block)

Dr. Sumbul Fatma


LEARNING OUTCOMES

- What are the amino acids?
- General structure.
- Classification of amino acids.
- Optical properties.
- Amino acid configuration.
- Non-standard amino acids.
- Derivatives of amino acids.


WHAT ARE AMINO ACIDS?

- ◉ Amino acids are the chemical units that combine to form proteins.
- ◉ Amino acids are a type of organic acid that contain both a carboxyl group (COOH) and an amino group (NH₂).
- ◉ Amino acids play central roles: as building blocks of proteins and as intermediates in metabolism.
- ◉ Humans can produce about half of amino acids. The others must be supplied in the food.
- ◉ When proteins are digested or broken down, amino acids are left.

GENERAL STRUCTURE




ZWITTERION



Net charge is zero
on the molecule

ISOELECTRIC POINT

- The pH at which the molecule carries no net charge.
 - In acidic solution- cationic.
 - In alkaline solution- anionic.




PK VALUE

- It is the ability of an acid to donate a proton (dissociate).
- Also known as **pKa** or acid dissociation constant.
- The pK values of α -carboxylic group is in the range of 2.2.
- The pK values of α -amino group is in the range of 9.4.

TITRATION CURVE OF GLYCINE

- ⦿ **pK1**- pH at which 50% of molecules are in cation form and 50% are in zwitterion form.
- ⦿ **pK2**- pH at which 50% of molecules are in anion form and 50% are in zwitterion form.
- ⦿ Buffering action is maximum around pK values and minimum at pI.



CLASSIFICATION OF AMINO ACIDS

○ **Based on the body requirement**


- **Essential amino acids:** cannot be made by the body.
e.g. histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine.
- **Nonessential amino acids:** produced by the body.
e.g. alanine, asparagine, aspartic acid, and glutamic acid.
- **Conditional amino acids:** not essential, except in time of illness or stress.
e.g. arginine, cysteine, glutamine, tyrosine, glycine, proline, and serine.

CONTINUED ...


- ◉ **According to the properties of the side chains:**
 - Nonpolar amino acids.
 - Uncharged amino acids.
 - Polar amino acids.

NONPOLAR AMINO ACIDS


- Each amino acid does not bind or give off protons or participate in hydrogen or ionic bonds.
- These amino acids promote hydrophobic interactions.
- In proteins found in aqueous solution, the side chains of the nonpolar amino acids tend to cluster together in the interior of the protein.
- The nonpolar R-group fill up the interior of the folded protein and help give it its 3D shape.
- In proteins located in hydrophobic environment, such as a membrane, the nonpolar R-groups are found on the outside surface of the protein, interacting with lipid environment to stabilize protein structure.




Glycine




Alanine




Valine




Leucine




Isoleucine




Phenylalanine



Tryptophan




Methionine




Proline

CONTINUED ...


- ◉ The structure of the proline amino acid differs from other nonpolar amino acids that the side chain of proline and its α -amino group form a ring structure (an imino group).




UNCHARGED AMINO ACIDS




Serine




Asparagine




Threonine



Glutamine



Tyrosine



Cysteine

CONTINUED ...


- ◉ These amino acids have zero net charge at neutral pH.

However

- ◉ The side chains of cysteine and tyrosine can lose a proton at an alkaline pH.
- ◉ Serine, threonine and tyrosine each contain a polar hydroxyl group that can participate in hydrogen bond formation.
- ◉ The side chains of asparagine and glutamine each contain a carbonyl group and an amide group, both of which can also participate in hydrogen bonds.

POLAR AMINO ACIDS


- Amino acids with acidic side chains:



- Aspartic and glutamic acids are proton donors.
- At neutral pH, these amino acids are fully ionized (negatively charged). So, they are called aspartate and glutamate.

CONTINUED ...


○ Amino acids with basic side chains:




- Histidine, Lysine and Arginine are proton acceptors.
- At neutral pH, lysine and arginine are fully ionized (positively charged).

OPTICAL PROPERTIES

- The α -carbon of most of the amino acids is attached to four different chemical groups- **asymmetric**.
- Asymmetric molecules are optically active, and symmetric molecules are optically inactive.
- All mammalian amino acids are optically active except glycine.
 - They rotate the plane of polarized light in a polarimeter.




AMINO ACID CONFIGURATION




CONTINUED ...

- ◉ L-Amino acids rotate polarized light to the left.
- ◉ D-Amino acids rotate polarized light to the right.
- ◉ Both L and D forms are chemically same.
- ◉ All mammalian amino acids are found in L-configuration.
- ◉ D-amino acids are found in antibiotics, plants and in the cell wall of microorganisms.


NON-STANDARD AMINO ACIDS




4-Hydroxyproline




ε-N,N,N-Trimethyllysine




3-Methylhistidine




5-Hydroxylysine




O-Phosphoserine




γ-Carboxyglutamate




ε-N-Acetyllysine




ω-N-Methylarginine



N-Acetylserine



N,N,N-Trimethylalanine



N-Formylmethionine

AMINO ACIDS DERIVATIVES

- ◉ **Gamma amino butyric acid** (GABA, a derivative of glutamic acid) and **dopamine** (from tyrosine) are neurotransmitters.
- ◉ **Histamine** (Histidine) is the mediator of allergic reactions.
- ◉ **Thyroxine** (Tyrosine) is an important thyroid hormone.

TAKE HOME MESSAGES

- ◉ Each amino acid has an α -carboxyl and a primary α -amino group (except for proline, which is an imino acid).
- ◉ At physiological pH, the α -carboxyl is dissociated.
- ◉ Each amino acid also contains twenty distinctive side chains and the chemical nature of this side chain determines the function of the amino acid.
- ◉ All free amino acids and charged amino acids in peptide chains, can serve as buffers.

TAKE HOME MESSAGES

- ◉ Buffering action of proteins is maximum around pK values and minimum at isoelectric point.
- ◉ All mammalian amino acids are optically active except glycine.
- ◉ All mammalian amino acids are found in L-configuration

REFERENCES

Lippincott's Illustrated reviews: Biochemistry 4th edition, Unit 1,
Chapter 1, Pages 1-12.