
LARYNX TRACHEA BRONCHI

Dr. Sameerah Shaheen





Objectives

- By the end of the lecture, you should be able to:
- Describe the Extent, structure and functions of the larynx.
- Describe the Extent, structure and functions of the trachea.
- Describe the bronchi and branching of the bronchial tree.
- Describe the **functions** of **bronchi and their divisions**.

LARYNX


- The larynx is the part of the respiratory tract which contains the vocal cords.
- In adult it is 2-inch-long tube.
- It opens <u>above</u> into the laryngeal part of the pharynx.
- <u>Below,</u> it is continuous with the trachea
- <u>The larynx has functions in</u>:
 - Respiration (breathing).
 - Phonation (voice production).
 - Deglutition (swallowing).







Relations


- The larynx is related to major <u>critical</u> structures in the neck.
- <u>Arteries</u>:
- <u>Carotid arteries:</u> (common, external and internal).
- <u>Thyroid arteries</u>: (superior & inferior thyroid arteries).
- Veins:
- Jugular veins, (external & internal)
- <u>Nerves:</u>
- Laryngeal nerves: (Superior laryngeal & recurrent laryngeal).
- vagus nerve.









- The cartilages are:
 - Connected by joints, membranes & ligaments.
 Moved by muscles
 - Moved by muscles.




MEMBRANES & LIGAMENTS



The **thyrohyoid membrane** is thickened in the median plane to form **median thyrohyoid ligament** and on both sides to form **lateral thyrohyoid ligaments**.





- Quadrangular membrane:
 - Or aryepiglottic membrane,
 - It extends <u>between</u> the arytenoid and epiglottis.
 - Its <u>lower free margin</u> forms the vestibular ligament which forms the <u>vestibular</u> fold (false vocal cord).
- Cricothyroid membrane (conus elasticus): ★
 - Its lower margin is attached to the upper border of cricoid cartilage.
 - Upper free margin forms
 Vocal ligament which
 forms the true vocal cord



Laryngeal Inlet


- It is the upper opening of the larynx.
- It faces upward and backward and <u>opens into</u> the laryngeal part of the pharynx, (laryngopharynx).
- Bounded by:
 - Anteriorly: by the upper margin of epiglottis (E)
 - Posteriorly & below by arytenoid cartilages (A)
 - Laterally by the Aryepiglottic folds (AEF)





Laryngeal Cavity

- Extends from laryngeal inlet to lower border of the cricoid cartilage
- Narrow in the region of the vestibular folds (rima vestibuli)
- Narrowest in the region of the vocal folds (rima glottidis)
- Divided into <u>three parts</u>:
 - A. Supraglottic part or vestibule: it is the part above the vestibular folds.
 - **B.** Ventricle: it is the part between the vestibular folds & the vocal folds.
 - **C.** Infraglottic part, the part below the vocal folds.
 - NB. The ventricle has an upward invagination called saccule which is rich in goblet cells.



<u>Mucous Membrane</u>

- The cavity is lined with ciliated columnar epithelium except the surface of the vocal cords.
- The surface of vocal folds, is covered with stratified squamous epithelium because of exposure to continuous trauma during phonation.
- It contains many mucous glands, more numerous in the region of the saccule (for lubrication of vocal folds).

<u>Muscles</u>

Laryngeal muscles are


divided into two groups:

- Extrinsic muscles: subdivided into two groups:
 - Elevators of the larynx.
 - Depressors of the larynx.
- Intrinsic muscles: subdivided into two groups:
 - Muscles controlling the laryngeal inlet.
 - Muscles controlling the movements of the vocal cords.

Extrinsic muscles of Larynx

Elevators of the Larynx

- A- The Suprahyoid Muscles: (MSGD)
 - 1. Mylohyoid.
 - 2. Stylohyoid.
 - 3. Geniohyoid.
 - 4. Digastric.
- B- The Longitudinal Muscles of the <u>Pharynx</u>:
 - Stylopharyngeus.
 - Salpingopharyngeus.
 - Palatopharyngeus.




Depressors of the Larynx


- The Infrahyoid Muscles:
 - Sternohyoid.
 - Sternothyroid.
 - Omo<mark>hyoid</mark>.


Intrinsic muscles of Larynx

Muscles Controlling the Laryngeal Inlet

- Oblique arytenoid.
- Aryepiglottic muscle.









- Muscle increasing the Length & Tension of Vocal Cords.
 - Cricothyroid.
 - NB. It is the only intrinsic muscle which found outside the larynx.

Muscles controlling the vocal cords


- Muscle <u>decreasing</u> the Length & Tension of Vocal Cords (relax vocal cords).
 - Thyroarytenoid (vocalis).



Movements of the Vocal Cords



Adduction



Abduction


Muscles controlling the vocal cords


Adductors (close rima glottis)

- Lateral cricoarytenoid.
- <u>Transverse</u> arytenoid.

Abductor (open rima glottis)

• **Posterior** cricoarytenoid






Nerve Supply (very important)

<u>Sensory</u>

- Above the vocal cords: Internal laryngeal nerve, branch of the superior laryngeal of the vagus nerve.
- Below the vocal cords: Recurrent laryngeal nerve, of the vagus nerve.


<u>Motor</u>

- All intrinsic muscles, are supplied by the recurrent laryngeal nerve except the cricothyroid.
- The <u>cricothyroid</u> is supplied by the <u>external laryngeal</u> <u>nerve</u> of <u>superior laryngeal</u> <u>of vagus</u>.



TRACHEA (windpipe)

- Mobile, fibrocartilginous tube, **5** inches long, **1** inch in diameter
- <u>Begins</u>: In the neck below the cricoid cartilage of the larynx (at lower border of cricoid cartilage at (C6).
- <u>Ends: In the thorax at the level of sternal angle (lower border of T4), by dividing into right and left principal (main, primary) bronchi.</u>
- The ridge at the bifurcation from inside is called <u>carina.</u>
- It is the most sensitive part of the respiratory tract and is associated with the cough reflex.




Relations in the Superior Mediastinum

<u>Anterior</u>

- Sternum.
- Thymus, (remains of thymus gland).
- Left brachiocephalic vein.
- Arch of aorta.
- Origin of:
- Brachiocephalic artery.
- left common carotid artery.

Posterior

- Esophagus.
- Left recurrent laryngeal nerve.



<u>Left side</u>

- Arch of aorta.
- Left common carotid artery.
- left subclavian artery.
- Left vagus nerve.
- Left phrenic nerve.
- Pleura.

۲

<u>Right side</u>

- Azygos vein
- Right vagus nerve.
- Pleura

Nerve Supply


- Branches of the vagus nerve and recurrent laryngeal nerve give <u>sensory fibers</u> to supply the <u>mucous membrane</u>.
- Branches from the <u>sympathetic trunks</u> supply the <u>trachealis muscle</u> and the <u>blood vessels</u>.

Lymphatic Drainage

Into the pretracheal and paratracheal lymph nodes.


Blood Supply

- <u>Arteries:</u> Branches from the <u>inferior thyroid</u> and <u>bronchial arteries</u> (from descending thoracic aorta)
- <u>Veins</u>: Drain to inferior thyroid veins.



Right Principal Bronchus

- About one inch long.
- Wider, shorter and more vertical than the left.
- Gives superior lobar bronchus before entering the hilum of the right lung.
- On entering the hilum it divides into middle and inferior lobar bronchi.



Left Principal Bronchus

- About two inches long.
- Narrower, longer and more horizontal than the right.
- Passes to the left below the aortic arch and in front of esophagus.
- On entering the hilum of the left lung it divides into superior and inferior lobar bronchi.

Bronchial Divisions

Within the lung, <u>each bronchus</u> divides and redivides into number of branches that <u>can be</u> <u>divided into two groups:</u>

Conduction zone branches -

- 1. Primary (main) bronchi.
- 2. Secondary (lobar) bronchi.
- Tertiary (segmental) bronchi. (supply the bronchopulmonary segment).
- 4. Smaller bronchi.
- 5. Bronchioles.
- 6. Terminal bronchioles.

Structures of the Respiratory Zone

