Lecture Title: <u>Respiratory Fungal Infections</u>

(Respiratory Block, Microbiology)

Lecturer name: Dr. Ahmed M. Albarrag



RESPIRATORY FUNGAL INFECTIONS

- > Respiratory System
- Rout of infection?
- Respiratory fungal infections are less common than viral and bacterial infections.

> Invasive diseases have significant difficulties in diagnosis and treatment.

RESPIRATORY FUNGAL INFECTION - ETIOLOGY



> YEAST

- Candidiasis \succ
- Opportunistic Cryptococcosis (Cryptococcus neoformans, C. gattii)
- Mould fungi \triangleright
 - Aspergillosis (Aspergillus species) \triangleright
 - Zygomycosis (Zygomycetes, e.g. Rhizopus, Mucor) \geq
 - Other mould \triangleright
- **Dimorphic fungi** \succ
 - \triangleright Histoplasma capsulatum
 - Blastomyces dermatitidis \triangleright

- Primary infections > Paracoccidioides brasiliensis
- Coccidioides immitis \triangleright

Primary Systemic Mycoses



- Infections of the respiratory system, (Inhalation)
- Dissemination seen in immunecompromised hosts
- Common in North America and to a lesser extent in South America. Not common in other parts of the World.

Etiologies are dimorphic fungi

In nature found in soil of restricted habitats. Primary pathogens They are highly infectious

They include:

Histoplasmosis,

Blastomycosis,

Coccidioidomycosis,

Paracoccidioidomycosis

Aspergillosis



Aspergillosis is a spectrum of diseases of humans and animals caused by members of the genus *Aspergillus*.

- These include
- (1) Mycotoxicosis
- (2) Allergy
- (3) Colonization (without invasion and extension) in preformed cavities
- (4) Invasive disease of lungs
- (5) Systemic and disseminated disease.

Aetiological Agents: Aspergillus species,

common species are:

A. fumigatus, A. flavus, A. niger, A. terreus and A. nidulans.

Classification of aspergillosis



Airways/nasal exposure to airborne Aspergillus

Persistence without disease - colonisation of the airways or nose/sinuses

Invasive aspergillosis

<u>Chronic aspergillosis</u> Aspergilloma of lung Maxillary (sinus) aspergilloma

<u>Allergic</u>

Allergic bronchopulmonary (ABPA) Allergic Aspergillus sinusitis

Risk factors



- Bone marrow/ organ transplantation
- Cancer: Leukemia, lymphoma,.. etc
- > AIDS
- > Drugs: Cytotoxic drugs, steroids,.. etc
- Diabetes



Aspergillosis



<u>Chronic Aspergillosis</u> (Colonizing aspergillosis)

(Aspergilloma OR Aspergillus fungus ball)

Signs include: Cough, hemoptysis, variable fever

Radiology will show mass in the lung , radiolucent crescent


Invasive pulmonary Aspergillosis


Signs: Cough , hemoptysis, fever, Leukocytosis Radiology will show lesions with halo sign

Invasive pulmonary aspergillosis




Note the Halo sign





Aspergilloma







Allergic bronchopulmonary (ABPA)

- Symptoms of Asthma
- Bronchial obstruction
- Eosinophilia
- > Wheezing +/-
- Also:
 - Skin test reactivity to Aspergillus
 - Serum antibodies to Aspergillus
 - Serum IgE > 1000 ng/ml

Common airborne Fungi





Aspergillus niger

Aspergillus fumigatus



FUNGAL SINUSITIS

Fungal sinusitis



Clinical:

- Nasal polyps and other symptoms of sinusitis
- In immunocompromised, Could disseminate to eye craneum (Rhinocerebral)
- > The most common cause in KSA is Aspergillus flavus
- > In addition to Aspergillus, there are other fungi that can cause fungal sinusitis
- >Aspergillus sinusitis has the same spectrum of Aspergillus disease in the lung

Diagnosis

- Clinical and Radiology
 Histology
- Culture
- Precipitating antibodies useful in diagnosisMeasurement of IgE level, RAST test

Treatment : depends on the type and severity of the disease and the immunological status of the patient

Diagnosis of aspergillosis

Specimen:

Respiratory specimens: Sputum, BAL, Lung biopsy,

Saud

- Other samples:
- Blood, etc.

Lab. Investigations:

➢ Direct Microscopy:

Giemsa Stain, Grecott methenamine silver stain (GMS)

Will show fungal septate hyphae

≻<u>Culture</u> on SDA

≻<u>Serology</u>:

Test for Antibody ELISA test for galactomannan Antigen


PCR: Detection of Aspergillus DNA in clinical samples

Diagnosis of aspergillosis





Cultures of Aspergillus



Smear: Septat fungal hyphae. Aspergillosis

Treatment of aspergillosis



Voriconazole

Alternative therapy Amphotericin B, Itraconazole, Caspofungin





Pulmonary zygomycosis Rhinocerebral zygomycosis

Risk factors

Transplant patients Malignancy AIDS Diabetic ketoacidosis

Many others

Pumonary zygomycosis



Acute

- Consolidation , nodules, cavitation, pleural effusion, hemoptysis
- Infection may extend to chest wall, diaphragm, pericardium.
 - Pulmonary infractions and hemorrhage
 - Rapid evolving clinical course

Early recognition and intervention are critical

≻<u>Etiology:</u>


Zygomycetes , Non-septate hyphae e.g. Rhizopus,

Diagnosis



Specimen:

- Respiratory specimens: Sputum, BAL, Lung biopsy,
- Other samples
- Lab. Investigations:
 - Direct Microscopy:
 Giemsa, Grecott methenami
 Will show broad no
 - Culture on SDA (no cyclohexii
- Serology: Not available



Treatment: Amphotericin B Surgery

Pneumocystosis (PCP)

Ring Saub Church

Pneumocystis pneumonia (PCP)

- It is interstitial pneumonia of the alveolar area.
- Affect compromised host
- Especially common in AIDS patients.
- **Etiology:**

Pneumocystis jiroveci

- Previously thought to be a protozoan parasite, but later it has been proven to be a fungus
- Does not grow in laboratory media e.g. SDA
- Naturally found in rodents (rats), other animals (goats, horses), Humans may contract it during childhood

Pneumocystosis



Laboratory Diagnosis:

Patient specimen: Bronchoscopic specimens (Bronchoalveolar lavage), Sputum, Lung biopsy tissue.

Histological sections or smears stained by GMS stain.


Immunuofluorescence (better sensitivity)

If positive will see <u>cysts</u> of hat-shape,

cup shape, crescent

Treatment:

Trimethoprim – sulfamethoxazole Dapsone



Thank You ③

(Respiratory Block, Microbiology)

Dr. Ahmed M. Albarrag

