Pneumonia

Community acquired pneumonia (CAP)

Objectives

- Discuss the epidemiology and pathophysiology of pneumonia and CAP
- Explain the different classifications of pneumonia
- Recognize clinical presentations associated with CAP
- Discuss the diagnosis and treatment of CAP
- Identify common etiological agents causing CAP and discuss their laboratory work up
- Discuss virulence factors and prevention of Streptococcus pneumoniae

Definition

- Pneumonia is an infection that leads to inflammation of the parenchyma of the lung (the alveoli) (consolidation and exudation)
- It may present as acute, fulminant clinical disease or as a chronic disease with a more prolonged course

Epidemiology

- Overall the rate of CAP 5-6 cases per 1000 persons per year
- Mortality 23%
 High, especially in <u>old people</u>
- Almost 1 million annual episodes of CAP in adults ≥ 65 yrs in the US


<u>Risk factors</u>

- Age ≤ 2 yrs, ≥ 65 yrs
- Alcoholism
- Smoking
- Asthma and COPD
- Aspiration
- Dementia
- Prior influenza
- HIV
- Immunosuppression
- Institutionalization
- Recent hotel : *Legionella*
- Travel, pets, occupational exposures- birds (*C. psittaci*)

Etiological agents

Infectious:

- Bacterial
- Fungal
- Viral
- Parasitic
- Non-infectious like: – Chemical – Allergen related




Pathogenesis

Two factors involved in the formation of pneumonia

– Pathogens

– Host defenses.



Defense mechanism of respiratory tract

- Filtration and deposition of environmental pathogens in the upper airways
- Cough reflux
- Mucociliary clearance
- Alveolar macrophages
- Humoral and cellular immunity
- Oxidative metabolism of neutrophils

Pathophysiology

- 1. Inhalation or aspiration of pulmonary pathogenic organisms into a lung segment or lobe.
- 2. Results from secondary bacteraemia from a distant source, such as Escherichia coli urinary tract infection and/or bacteraemia (less commonly).
- 3. Aspiration of oropharyngeal contents (multiple pathogens).


Classification


Pneumonia classified according to:

- 1. Pathogen
 - Bacterial
 - Typical
 - Atypical
 - Viral
 - Fungal
 - Parasite
- 2. Anatomy
- 3. Acquired environment

Classification by anatomy

- 1. Lobar: entire lobe
- 2. Lobular: (bronchopneumonia).
- 3. Interstitial





Lobar pneumonia

Classification by acquired environment

♦ Community acquired pneumonia (CAP)

♦ Hospital acquired pneumonia (HAP)


◆ Nursing home acquired pneumonia (NHAP)

CAP- fever+ productive cough + infiltrate

• <u>CAP</u>: pneumonia acquired outside of hospitals or extended-care facilities

Typical

- Strept. pneumoniae
 - (lobar pneumonia)
- Haemophilus influenzae
- Moraxella catarrhalis
- S. aureus
- Gram-negative organisms



Atypical

- Mycoplasma pneumoniae
- Chlamydia pneumoniae
- Legionella pneumophila


Community acquired pneumonia					
• Strep pneumonia	48%				
• Viral	23%				
• Atypical orgs (MP,LG,CP)	22%				
• Haemophilus influenza	7%				
• Moraxella catharralis	2%				
• Staph aureus	1.5%				
• Gram –ive orgs	1.4%				
• Anaerobes					

Typical pneumonia Clinical manifestation

- The onset is acute
- Prior viral upper respiratory infection
- Respiratory symptoms
 - Fever
 - Shaking chills
 - Cough with sputum production (rusty-sputum)
 - Chest pain- or pleurisy
 - Shortness of breath


Diagnosis

- Clinical
 - History & physical
- X-ray examination
- Laboratory
 - CBC- leukocytosis
 - Sputum
 - Gram stain- 15%
 - Culture
 - Blood culture- 5-14%
 - Pleural effusion gram + culture










Streptococcus pneumoniae


- Gram positive diplococci
- Alpha hemolytic streptococci
- Catalase negative
- Normal flora of upper respiratory tract in 20-40% of people
- Causes:
 - Resp infections
 - pneumonia, sinusitis, otitis,
 - Non resp infections
 - bacteremia, meningitis

Streptococcus pneumoniae

- Virulence factors:
 - Capsule
 - More than 90 capsular types
 - Pneumolysin
 - Autolysin
 - Neuraminidase

Prevention: vaccination






Streptococcus pneumoniae

Sensitive to OptochinLysed by bile (bile soluble)







Atypical pneumonia

 \bullet

- Chlamydia pneumonia
- Mycoplasma pneumonia
- Legionella spp
- Psittacosis (*Chlamydia psittaci*)
- Q fever (*Coxiella burnettii*)

- Approximately 15% of all CAP
- Not detectable on gram stain
- Won't grow on standard media
- Some don't have a bacterial cell wall → Don't respond to βlactams

Atypical pneumonia

Symptoms

- Insidious onset
- Mild to severe
- Headache
- Malaise
- Fever
- Dry cough
- Arthralgia / myalgia

Signs

• Minimal

- Low grade fever
- Few crackles
- Rhonchi

Diagnosis & Treatment


- Diagnosis:
 - X-ray
 - CBC
 - Mild elevation WBC
 - U&Es
 - Low serum Na (Legionalla)
 - LFTs
 - **†** ALT
 - † Alk Phos
 - Sputum Culture on special media (BCYE) for *Legionella*
 - Urine antigen for *Legionella*
 - Serology for detecting antibodies
 - DNA detection

- Treatment:
 - Macrolide
 - Quinolones
 - Tetracycline
 - B lactams have no activity
- Treat for 10-14 days

Mycoplasma pneumonia

- Eaton's agent (1944)
- No cell wall
- Common
- Rare in children and in > 65
- People younger than 40.
- Crowded places like schools, homeless shelters, prisons.
- Can cause URT symptoms
- Usually mild and responds well to antibiotics.
- Can be very serious

- May be associated with extra pulmonary findings:
 - skin rash, hemolysis, myocarditis, pancreatitis, encephalitis
- Diagnosis:
 - Serology
 - NAAT
 - Culture can be done but requires special media and slow grower (weeks)




Mycoplasma pneumonia Cx-ray

Chlamydia pneumonia

- Obligate intracellular organism
- 50% of adults sero-positive
- Mild disease
- Sub clinical infections common
- 5-10% of community acquired pneumonia
- Diagnosis:
 - Serology
 - NAAT

Psittacosis



- Chlamydia psittaci
- Exposure to birds
- Bird owners, pet shop employees, vets
- Parrots, pigeons and poultry
- Birds often asymptomatic

Q fever (Coxiella burnetti)

- Exposure to farm animals mainly sheep
 - Spread by inhalation of infected animal birth products
- Pneumonia is acute form of infection
- Diagnosis: serology



Legionella pneumophila

- Legionnaire's disease
- Serious outbreaks linked to exposure to cooling towers
- Can be very severe and lead to ICU admission.

- Can cause
 - Hyponatraemia common
 - (<130mMol)
 - Bradycardia
 - WBC < 15,000
 - Abnormal LFTs
 - Raised CPK
 - Acute Renal failure

Legionella pneumophila

• Diagnosis:

- Specimen: sputum
- Culture on specialized media (BCYE)
- > DFA (low sensitivity)
- ≻ NAAT
- Urine antigen testing

• Pontiac fever:

- Non pneumonic
- Influenza like illness
- Self limiting
- Related to exposure to environmental aerosols containing Legionella (potentially reaction to bacterial endotoxins)





Antibiotic Treatment of CAP

- Factors to consider in selection of antibiotic:
 - Co morbidities
 - Previous antibiotic exposure in last 3 months
 - Severity
 - Out patient management vs requiring inpatient admission vs requiring ICU

		Macrolides	Doxycycline	Levofloxacin	B-lactam And Macrolide B-lactam And Levo
Outpatient, healthy patient with no exposure to antibiotics in the last 3 months	- <i>S. pneumoniae</i> -Atypical pathogens -Viral				
Outpatient, patient with comorbidity or exposure to antibiotics in the last 3 months	As above + Anaerobes S. aureus				
Inpatient : Not ICU	Same as above + coliforms				
Inpatient : ICU	Same as above + <i>Pseudomonas</i>				

References

- Ryan, Kenneth J.. Sherris Medical Microbiology, Seventh Edition. McGraw-Hill Education.
 - Lower respiratory tract infections, part of the chapter on Infectious Diseases: Syndromes and Etiologies
 - Streptococci, chapter 25
 - Legionella and Coxiella, chapter 34
 - Mycoplasma, chapter 38
 - Chlamydia, chapter 39