


ACID BASE BALANCE

Acid-Base Balance



24-46

DR SYED SHAHID HABIB
MBBS DSDM FCPS
Professor & Consultant Clinical Physiology
Dept. of Physiology
College of Medicine & KKHU

OBJECTIVES

At the end of this lecture you should be able to:

- **Define: acid and base.**
- **Explain what is meant by strong and weak acids and bases**
- **List and identify the names/formulas for the common strong acids and strong bases.**
- **To explain the role of Henderson-Hasselbalch equation in acid-base regulation**



ACIDS

Acids dissociate in solution to liberate free H^+ ions

- STRONG acids (eg. Hydrochloric acid i.e. HCl) completely dissociate (to H^+ and Cl^-)**
- WEAK acids (H_2CO_3) have more limited dissociation**

BASES

- Bases are ions or molecules that bind free H^+ and remove it from solution


eg. HCO_3^- combines with H^+ to form H_2CO_3

- Alkali is a molecule formed by one of the alkaline metals. (Na, K, Li) with a highly basic ion such as a hydroxyl ion (OH^-).

pH SCALE

20	7.70
30	7.52
40	7.40
50	7.30
60	7.22

H⁺ = 80 -last two digits of pH



pH

- pH is the log of the reciprocal of the H⁺ ion concentration

$$\text{pH} = \log (1 / [\text{H}^+])$$

OR

$$\text{pH} = - \log([\text{H}^+])$$

WHY WE EXPRESS IT AS pH?

pH

- The normal H ion concentration in blood is 40 nmol/l or 0.00004 mmol/l or 0.000000004 Eq/L).
- For example for Na it is 140 mmol/l
- Because H ion concentration in blood is so low that it is expressed in negative log to the base 10 of H ion concentration

40 nmol/l or 0.00004 mmol/l is equal to pH 7.4

pH and H⁺ ion concentration

pH	H ⁺ ion in nmol/lit
• 6.0	• 1000
• 7.0	• 100
• 8.0	• 10
• 9.0	• 1.0

One point change in pH results in a ten fold change in H⁺ ion conc.

WHAT IS THE NORMAL BODY pH?


7.35 – 7.45

	H ⁺ Concentration (mEq/L)	pH
Extracellular fluid		
Arterial blood	4.0×10^{-5}	7.40
Venous blood	4.5×10^{-5}	7.35
Interstitial fluid	4.5×10^{-5}	7.35
Intracellular fluid	1×10^{-3} to 4×10^{-5}	6.0 to 7.4
Urine	3×10^{-2} to 1×10^{-5}	4.5 to 8.0
Gastric HCl	160	0.8

IMPORTANCE

- H^+ ions are deadly because they can affect cell function by altering the charge of functional proteins including enzymes
- H ions are very reactive cations and bind to protein anions strongly if they are in high amounts and impair their activity

**ACTIVITIES OF ALL ENZYME SYSTEMS
IN THE BODY IS INFLUENCED BY
HYDROGEN IONS**



ACID PRODUCTION

H⁺ is continually produced by metabolic activity:

❖ **Volatile acids: (e.g. carbonic acid, H₂CO₃; formation catalyzed by carbonic anhydrase)**




ACID PRODUCTION (Cont.)


- **Non-volatile acids:** ingested acids and products of fat, amino acid, and sugar metabolism:
 - e.g. phosphoric acid, lactic acid, butyric acid
- **Incomplete Carbohydrate and Fat Metabolism Produces Nonvolatile Acids (strenuous exercise, hemorrhagic or cardiogenic shock, uncontrolled diabetes mellitus, starvation, and alcoholism)**

ACID LOAD

- **Amino Acid Metabolism yields about 50 meq/day for example H_2SO_4 , HCl , and H_3PO_4**
- **CO_2 production yields 12,500 meq/day or mmol/day 300 L of CO_2**
- **Normal daily diet yields 80 meq/day**



The 7 Most Alkaline Foods



HENDERSON-HASSELBACH EQUATION


Relates pH to the Ratio of the Conc. of Conjugate Base and Acid

$$\text{pH} = \text{pK} + \log \frac{\text{Base}}{\text{Acid}}$$

The ratio of dissociated to undissociated forms of an acid is **CONSTANT (K)** and shows the **Strength of an Acid**

$$K = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

eg: $K = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$



$$K' = \frac{\text{H}^+ \times \text{HCO}_3^-}{\text{H}_2\text{CO}_3}$$

$$\text{H}^+ = K' \times \frac{\text{H}_2\text{CO}_3}{\text{HCO}_3^-}$$

$$\text{H}^+ = K \times \frac{0.03 \times \text{CO}_2}{\text{HCO}_3^-}$$

$$\text{H}^+ = K \times \frac{0.03 \times \text{CO}_2}{\text{HCO}_3^-}$$

$$H^+ = K \times \frac{0.03 \times CO_2}{HCO_3^-}$$

$$-\log H^+ = -\log K \times -\log \frac{0.03 \times CO_2}{HCO_3^-}$$

$$pH = pK \times \log \frac{HCO_3^-}{CO_2}$$

The Henderson-Hasselbalch Equation Relates pH to the Ratio of the Concentrations of Conjugate Base and Acid

pK

Dissociation Constant

pK (also a log) is where concentration of both components of the buffer are equal.

(REMEMBER to maintain plasma pH at 7.4, there needs to be much more HCO_3^- than H_2CO_3)

pK

- $\text{pH} = \text{pK} + \log \frac{\text{Base}}{\text{Acid}}$
- $\text{pH} = \text{pK} + \log \frac{50}{50}$
- $\text{pH} = \text{pK}$






FIGURE 25.1

A titration curve for a phosphate buffer. The pK_a for H₂PO₄⁻ is 6.8. A strong acid (HCl) (right to left) or strong base (NaOH) (left to right) was added and the resulting solution pH recorded (y-axis). Notice that buffering is best (i.e., the change in pH upon the addition of a given amount of acid or base is least) when the solution pH is equal to the pK_a of the buffer.

NORMAL OPERATING POINT FOR BICARBONATE/CARBONIC ACID BUFFER SYSTEM



Normal
7.35 – 7.45

ACIDOSIS

ALKALOSIS

Death

6.8

Normal

7.8

Death




1

1.2 meq/L



2

24 meq/L





OBJECTIVES

At the end of this lecture you should be able to:

- **To define buffer system and discuss the role of blood buffers and to explain their relevant roles in the body**
- **To describe the role of kidneys in the regulation of acid-base balance**
- **To describe the role of lungs in the regulation of acid-base balance**



BUFFER SYSTEMS


- Buffer is a solution which minimizes pH changes when acid or base is added to a solution (any substance that can reversibly bind H^+)
- It consists of a WEAK ACID and its conjugate base (or a weak base and its conjugate acid)
- For example in Bicarbonate buffer system H_2CO_3 is the weak acid and $NaHCO_3$ is its conjugate base.

Buffers Promote the Stability of pH

BUFFER SYSTEMS




ADD STRONG ACID



ADD STRONG BASE



BUFFER SYSTEMS



ADD STRONG ACID



ADD STRONG BASE



BUFFER POWER

- **Depends on relative amount of Acid and Base in a Buffer solution**
- **It is maximum when both are in equal amounts**
- **Absolute concentration of Buffers in body fluids is also important**
- **If the pH of medium is near pK of buffer system it becomes more effective**

$$\text{pH} = \text{pK} + \log [\text{Base}] / [\text{Acid}]$$

$$\text{pH} = \text{pK} + \log \text{HCO}_3^- / \text{H}_2\text{CO}_3$$

$$\text{pH} = 6.1 + \log 20 / 1$$

It is not only the amount of base and acid that is important but the ratio between them must remain constant

TABLE 25.1**Major Chemical pH Buffers in the Body**

Buffer	Reaction
Extracellular fluid	
Bicarbonate/CO ₂	$\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3 \rightleftharpoons \text{H}^+ + \text{HCO}_3^-$
Inorganic phosphate	$\text{H}_2\text{PO}_4^- \rightleftharpoons \text{H}^+ + \text{HPO}_4^{2-}$
Plasma proteins (Pr)	$\text{HPr} \rightleftharpoons \text{H}^+ + \text{Pr}^-$
Intracellular fluid	
Cell proteins (e.g., hemoglobin, Hb)	$\text{HHb} \rightleftharpoons \text{H}^+ + \text{Hb}^-$
Organic phosphates	$\text{Organic-HPO}_4^- \rightleftharpoons \text{H}^+ + \text{organic-PO}_4^{2-}$
Bicarbonate/CO ₂	$\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3 \rightleftharpoons \text{H}^+ + \text{HCO}_3^-$
Bone	
Mineral phosphates	$\text{H}_2\text{PO}_4^- \rightleftharpoons \text{H}^+ + \text{HPO}_4^{2-}$
Mineral carbonates	$\text{HCO}_3^- \rightleftharpoons \text{H}^+ + \text{CO}_3^{2-}$



pH DEFENCE MECHANISMS IN THE BODY

- **Chemical buffering (First Line)**
Acid-Base buffer systems of the body fluids
- **Respiratory response (Second Line)**
Respiratory center
- **Renal response (Third Line)**
Kidneys [slow to respond & powerful]

BUFFER SYSTEMS

Weak Acid

Conjugate Base



BODY BUFFER SYSTEMS

– BICARBONATE/CARBONIC ACID:



- $\text{pK} = 6.1$
- major plasma buffer

– PHOSPHATE: $\text{HPO}_4^- / \text{H}_2\text{PO}_4$


- $\text{pK} = 6.8$
- major intracellular and urine buffer
- conc. in ECF is only 8 % of bicarbonate buffer

IMPORTANT NOTE: A pK_a of 6.8 Makes Phosphate a Good Buffer in ECF however, its plasma conc. is low (about 1 mmol/L) unlike HCO_3^- which is 24 mmol/L

BODY BUFFER SYSTEMS

- **AMMONIA: $\text{NH}_3 / \text{NH}_4^+$**
 - **pK = 9.0**
 - **used to buffer the urine**
- **PROTEINS (Amphoteric) : Prot / H Prot**
 - **important in ICF**
- **HEMOGLOBIN: Hb / HHb**
 - **important in ICF**

Respiratory Regulation of Acid-Base Balance




Doubling the ventilation rate raises the pH 7.4→7.63


Normal

Increasing alveolar ventilation to about twice normal raises the pH of the extracellular fluid by about 0.23.

Effect of blood pH on rate of alveolar ventilation.




RENAL MECHANISMS TO REGULATE BODY pH



About 80 to 90 per cent of the bicarbonate reabsorption (and H⁺ secretion) occurs in the proximal tubule


HYDROGEN ION SECRETION



HCO_3^- Is "Titrated" Against H^+ in the Tubules

HYDROGEN ION SECRETION

Late DCT & CT




- α intercalated cell \rightarrow acid-secretion
- β intercalated cell \rightarrow bicarbonate-secretion


Remember that I Cells have H-ATPase and H/K-ATPase

PHOSPHATE BUFFER SYSTEM


PCT, LOH (A), DCT



AMMONIA BUFFER SYSTEM



AMMONIA BUFFER SYSTEM




OBJECTIVES

At the end of this lecture you should be able to:

- **To explain the principles of blood gas and acid-base analysis**
- **To interpret blood gas analysis and diagnose various acid base disorders**
- **Describe causes of acid base disorders**
- **Understand use of acid base nomograms**




BASE/ACID RATIO




ARTERIAL BLOOD ANALYSIS

ANALYTE	REF. RANGE
pH	7.4 ± 0.05
PO₂	75-100 mmHg (10.0-13.3 kpa)
PCO₂	36.0-46.0 mmHg (4.8-6.1 kpa)
HCO₃⁻	22.0-26.0 mmol/L
O₂ Saturation	95-100 %
Base Excess	± 2.5 (Normal)




DISORDER	IMPORTANT CAUSES
Respiratory Acidosis	<ul style="list-style-type: none">• Inadequate ventilation
Respiratory Alkalosis	<ul style="list-style-type: none">• Hyperventilation
Metabolic Acidosis	<ul style="list-style-type: none">• Diabetic ketoacidosis,• Lactic acidosis• Ethylene glycol or salicylate poisoning (elevated anion gap)• Renal tubular acidosis & CRF• Diarrhea, ileostomy (normal anion gap)
Metabolic Alkalosis	<ul style="list-style-type: none">• Excessive alkali ingestion (antacids)• H⁺ loss (vomiting)

ACIDOSIS AND ALKALOSIS




COMPENSATION OF ACIDOSIS



COMPENSATION OF ALKALOSIS

Normal

H^+ ↓



UNCOMPENSATED

ACIDOSIS	
RESPIRATORY	METABOLIC
$H^+ \uparrow$ $pH \downarrow$ $CO_2 \uparrow$ $HCO_3 \text{ N}$	$H^+ \uparrow$ $pH \downarrow$ $CO_2 \text{ N}$ $HCO_3 \downarrow$
ALKALOSIS	
RESPIRATORY	METABOLIC
$H^+ \downarrow$ $pH \uparrow$ $CO_2 \downarrow$ $HCO_3 \text{ N}$	$H^+ \downarrow$ $pH \uparrow$ $CO_2 \text{ N}$ $HCO_3 \uparrow$

COMPENSATED

ACIDOSIS	
RESPIRATORY	METABOLIC
$H^+ \uparrow$ $pH \downarrow$ $CO_2 \uparrow$ $HCO_3 \uparrow \uparrow \uparrow$	$H^+ \uparrow$ $pH \downarrow$ $CO_2 \downarrow \downarrow \downarrow$ $HCO_3 \downarrow$
ALKALOSIS	
RESPIRATORY	METABOLIC
$H^+ \downarrow$ $pH \uparrow$ $CO_2 \downarrow$ $HCO_3 \downarrow \downarrow \downarrow$	$H^+ \downarrow$ $pH \uparrow$ $CO_2 \uparrow \uparrow \uparrow$ $HCO_3 \uparrow$

ACIDOSIS AND ALKALOSIS

	pH	H ⁺	Pco ₂	HCO ₃ ⁻
Normal	7.4	40 mEq/L	40 mm Hg	24 mEq/L
Respiratory acidosis	↓	↑	↑↑	↑
Respiratory alkalosis	↑	↓	↓↓	↓
Metabolic acidosis	↓	↑	↓	↓↓
Metabolic alkalosis	↑	↓	↑	↑↑

$$\text{ANION GAP} = \{[\text{Na}^+] + [\text{K}]\} - \{[\text{HCO}_3^-] + [\text{Cl}^-]\}$$

16-18 mmol/L
12-16 if K⁺ is not included

High anion gap metabolic acidosis

Methanol intoxication

Uremia

Lactic acid

Ethylene glycol intoxication

p-Aldehyde intoxication

Ketoacidosis

Salicylate intoxication

Normal anion gap metabolic acidosis


Diarrhea

Renal tubular acidosis

Ammonium chloride ingestion

Table 31-2 Plasma or Extracellular Fluid Factors That Increase or Decrease H⁺ Secretion and HCO₃⁻ Reabsorption by the Renal Tubules

Increase H ⁺ Secretion and HCO ₃ ⁻ Reabsorption	Decrease H ⁺ Secretion and HCO ₃ ⁻ Reabsorption
↑ P _{CO} ₂	↓ P _{CO} ₂
↑ H ⁺ , ↓ HCO ₃ ⁻	↓ H ⁺ , ↑ HCO ₃ ⁻
↓ Extracellular fluid volume	↑ Extracellular fluid volume
↑ Angiotensin II	↓ Angiotensin II
↑ Aldosterone	↓ Aldosterone
Hypokalemia	Hyperkalemia



SUMMARY

