



Autonomic Nervous System

DR JAMILA EL MEDANY

OBJECTIVES

At the end of the lecture, students should be able to:


- *Define the autonomic nervous system.*
- *Describe the structure of autonomic nervous system*
- *Trace the preganglionic & postganglionic neurons in both sympathetic & parasympathetic nervous system.*
- *Enumerate in brief the main effects of sympathetic & parasympathetic system*

Autonomic Nervous System

- Concerned with the innervation and control of Involuntary structures: visceral organs, smooth & cardiac muscles and glands
- Function: maintain homeostasis of the internal environment along with the Endocrine system
- Located: both in the central and peripheral nervous systems.
- Regulated (controlled) by Hypothalamus.


Autonomic Nervous System

- Unlike the somatic nervous system, the **Efferent** pathway of the autonomic nervous system is made up of two neurons called as **preganglionic** and **postganglionic** neurons
- The cell bodies of the **preganglionic** neurons are located in the **brain and spinal cord**. Their axons synapse with the **postganglionic** neurons whose cell bodies are located in the **autonomic ganglia**



■ Based on the **anatomical**, **physiological** and **pharmacological** characteristics, the autonomic nervous system is divided into:


- **Sympathetic**: Activated during **exercise, excitement, and emergencies**. **“fight, flight, or fright”**
- **Parasympathetic**: Concerned with **conserving energy**. **“rest and digest”**



Both divisions operate in conjunction with one another (have **antagonistic** control over the viscera) to maintain a **stable internal environment**


Para sympathetic

Sympathetic



Sympathetic Division



- Preganglionic Neurons: located in the **lateral gray horn** of **T₁-L₂** segments of spinal cord (**Thoracolumbar outflow**)



Post Ganglionic Ganglia


■ Located nearer the central nervous system as:

- **Prevertebral:** celiac & mesenteric
- **Paravertebral** forming sympathetic chain




Paravertebral Ganglia

- ❑ They are interconnected to form **2 sympathetic chains**, one on each side of vertebral column.
- ❑ Number of ganglia:
 - **Three** in **cervical** part of chain
 - **Eleven to twelve** in thoracic part
 - **Four** in **lumbar & sacral** parts each.
- ❑ The chains end into a common '**ganglion impar**' in front of coccyx




Preganglionic fibers

- Run in the **ventral roots** of the spinal nerve
- Travel through the **spinal nerve**, and then join the sympathetic chain via the **White Rami Communicans. (WRC)**




■ **Within the sympathetic chain, these fibers may:**

1. **Ascend, descend or remain at the same level to synapse with neurons (postganglionic) of paravertebral ganglia located in sympathetic chain.**
2. **Leave the sympathetic chain (without synapse) to reach coeliac & mesenteric ganglia (around branches of abdominal aorta) to synapse with their neurons (postganglionic).**



Postganglionic fibers

- From the sympathetic chain ganglia enter again into the spinal nerve through **Grey Rami Communicantes (GRC)** to supply *structures in head & thorax + blood vessels & sweat glands*
- From the cells of coeliac & mesenteric ganglia supply *abdominal & pelvic viscera.*



Parasympathetic Division


Preganglionic neurons

Located in:

■ Nuclei of the 3rd, 7th, 9th & 10th cranial nerves, in the brain stem (Cranial outflow)


&

■ The lateral gray horn of S₂-S₄ segments of spinal cord (Sacral outflow)




Cranial Outflow

- Preganglionic fibers from **cranial outflow** are carried by **3rd, 7th, 9th & 10th cranial nerves** and terminate in **ciliary, pterygopalatine, submandibular, otic & peripheral ganglia**
- Postganglionic fibers innervate **organs of the head, neck, thorax, and abdomen**



Sacral Outflow

- Preganglionic fibers from sacral outflow are carried by **pelvic splanchnic nerves** to peripheral ganglia in pelvis where they synapse.
- Postganglionic fibers innervate organs of the pelvis and lower abdomen



Autonomic nervous system

Structure	Sympathetic effect	Parasympathetic effect
Iris of eye	Dilates pupil	Constricts pupil
Ciliary muscle of eye	Relaxes	Contracts
Salivary glands	Reduces secretion	Increases secretion
Lacrimal gland	Reduces secretion	Increases secretion
Heart	Increases rate and force of contraction	Decreases rate and force of contraction
Bronchi	Dilates	Constricts
Gastrointestinal tract	Decreases motility	Increases motility
Sweat glands	Increases secretion	
Erector pili muscles	Contracts	

Thank U & Good Luck

