

Glycolysis



Color index :







Main text

IMPORTANT

Extra Info

Drs Notes


Objectives:

-  Recognize glycolysis as the major oxidative pathway of glucose.
-  List the main reactions of glycolytic pathway.
-  Discuss the rate-limiting enzymes/Regulation.
-  Assess the ATP production (aerobic/anaerobic).
-  Define pyruvate kinase deficiency hemolytic anemia.
-  Discuss the unique nature of glycolysis in RBCs.

Glycolysis

- Glycolysis means breaks down glucose and forms pyruvate with the production of two molecules of ATP. The pyruvate is the end product of glycolysis.
- Glycolysis, the major pathway for glucose oxidation, **occurs in the cytosol of all cells.**
- It is unique, in that it can function **either aerobically or anaerobically**, depending on the availability of oxygen and intact mitochondria.
- It allows tissues to survive in presence or absence of oxygen, e.g., skeletal muscle.
- **RBCs**, which lack mitochondria, are completely reliant on glucose as their metabolic fuel, and metabolizes it by **anaerobic glycolysis**.

Types of DNA	Aerobic	Anaerobic
Presence of mitochondria and oxygen	Must have mitochondria and oxygen .	In absent of mitochondria or oxygen or both.
Number of steps	10 steps (reactions).	11 steps (reactions).
End product	End product is Pyruvate.	End product is Lactate or lactic acid .
Byproducts	Produce 2 ATP and 2 NADH = (8 ATP).	produce 2 ATP




Glycolysis , an example of metabolic pathway


- ★ A product of first reaction becomes a substrate for second reaction.

Aerobic glycolysis (overview)

↔ reversible
→ irreversible




Aerobic glycolysis (1)




- Kinase means phosphorylation enzyme " an enzyme that adds a phosphate group ",
- Glucokinase is a hexokinase isozyme.
- ATP In.
- ADP Out.
- Regulation of hexokinase and glucokinase , [click here](#) .



- isomerase means it changes the configuration.
- Isomerization from aldose form to ketos form.



- Bisphosphate means 2 phosphate groups not close to each others while Biphosphate means 2 phosphate groups close to each others.
- ATP in.
- ADP out.
- can be regulated (inhibited or activated) , more details [click here](#) :
 - + AMP and Fructose 2,6-bisphosphate.
 - - ATP and citrate.



Aerobic glycolysis (2)


Reactions 1-5 consume 2 molecules of ATP



- Aldolase breaks down carbon-carbon bond (*split the molecule*).




- *isomerase* means it changes the configuration.
- Isomerization from ketos form to aldose form.
- When the body needs fats it will convert glyceraldehyde 3-phosphate to dihydroxyacetone phosphate but in glycolysis the body needs energy so it convert dihydroxyacetone phosphate to glyceraldehyde 3-phosphate. By the end of reaction 5 we will have 2 molecule of glyceraldehyde 3-phosphate one from reaction 4 and the other from reaction 5 .



- 2x P_i and NAD⁺ in.
- 2x NADH and H⁺ out.

"About NAD⁺ and NADH , keep in mind that the reverse can happen "




Aerobic glycolysis (3)

Reactions 6-10 produce two molecules in each reaction.



- **2x** ATP in.
- **2x** ADP out.


"Keep in mind that the reverse can happen"



- **2x** H₂O out.



- **2x** ATP in.
- **2x** ADP out.
- can be regulated (activated) :
- **+** Fructose 1,6-Bisphosphate .
- Regulation of pyruvate kinase A , [click here](#) .



Summary of the aerobic glycolysis In order for you to gain a better understanding In tables

Reaction 1	
Reactant	D-Glucose
Product	Glucose 6-phosphate
Enzyme	1- Hexokinase in most tissue 2- glucokinase in hepatocyte
Action	adding phosphate
Consume	1 ATP

Reaction 4	
Reactant	Fructose 1,6-bisphosphate
Product	Dihydroxyacetone phosphate and Glycerdehyde 3-phosphate
Enzyme	Aldolase
Action	Break down carbon-carbon bond (split the molecule)
Consume	-

Reaction 2	
Reactant	Glucose 6-phosphate
Product	Fructose 6-phosphate
Enzyme	Phosphoglucose isomerase
Action	isomerization from aldose form to ketose form
Consume	-

Reaction 5	
Reactant	Dihydroxyacetone phosphate
Product	Glycerdehyde 3-phosphate
Enzyme	Triose phosphate isomerase
Action	Isomerization from ketose form to aldose form
Consume	-

Reaction 3	
Reactant	Fructose 6-phosphate
Product	Fructose 1,6-bisphosphate
Enzyme	Phosphofruktokinase-1 (PFK-1)
Action	adding phosphate
Consume	1 ATP

Reaction 6	
Reactant	2x Glycerdehyde 3-phosphate
Product	2x 1,3 Bisphosphoglycerate
Enzyme	Glycerdehyde 3-phosphate Dehydrogenase.
Action	Adding Phosphate group to the molecule
Byproduct	NADH

Summary of the aerobic glycolysis In order for you to gain a better understanding In tables ... contd

Reaction 7	
Reactant	2x 1,3 Bisphosphoglycerate
Product	2x 3-Phosphoglycerate
Enzyme	Phosphoglycerate kinase
Action	Phosphate group add to ADP to become ATP.
Produce	2x ATP

Reaction 8	
Reactant	2x 3-Phosphoglycerate
Product	2x 2-Phosphoglycerate
Enzyme	Phosphoglycerate mutase
Action	The P group change position from O in carbon-3 to O in carbon-2
Consume	-

Reaction 9	
Reactant	2x 2-Phosphoglycerate
Product	2x Phosphoenolpyruvate
Enzyme	Enolase
Action	Remove water
Consume	-


Reaction 10	
Reactant	2x Phosphoenolpyruvate
Product	2x Pyruvate
Enzyme	Pyruvate kinase A
Action	Phosphate group add to ADP to become ATP
Produce	2x ATP

Regulation of enzymes

“ Hexokinase / Glucokinase “

- Regulation of hexokinase (in most cells) and glucokinase (in liver or we can say hepatocyte “ hepato means liver and cyte means cell).
- **Hexokinase** : when Glucose 6-phosphate is abundant it will indicate to the cell that it doesn't need hexokinase anymore and it will be **inhibited directly** .
- **Glucokinase** : is an enzyme that facilitates phosphorylation of glucose to glucose-6- phosphate. It is an isozyme of hexokinase.
 1. In the presence of high Fructose-6- phosphate, glucokinase translocates and binds tightly to **GKRP** (Glucokinase regulatory protein) in the **nucleus**, making it **inactive** “ it will not work “ .
 2. When glucose levels are high in blood and **hepatocytes (GLUT-2)**, glucokinase is released from **GKRP** (Glucokinase regulatory protein) and enters the cytosol .

	Hexokinase	Glucokinase
Site	All tissues (Hexokinase have 4 isoforms all of them in all tissues except glucokinase)	Hepatocytes and pancreas cells
Regulation by: (inhibition)	Glucose 6-phosphate	Fructose-6- phosphate
Regulation by: (stimulation)	-	Glucose
Function	Phosphorylates “ adds a phosphate group on “ glucose inside the body cells thus glucose concentration in blood is higher than glucose concentration in the cell.	After meals, it removes sugar “ glucose “ coming in the portal circulation, converting it into Glucose 6-phosphate




Regulation of enzymes

“ PFK-1 / Pyruvate kinase A “

Phosphofruktokinase-1 (PFK-1) “Enzyme” :


Rate limiting enzyme

“ يعني ياتر باله glycolysis كله “



- Reaction number 3 in glycolysis it's **irreversible** reaction
 ⊕ **stimulated by AMP and Fructose 2,6-bisphosphate .**
 ⊖ **inhibited by ATP and citrate.**
- They are not involved in the chemical reaction, they have an allosteric effect.


Pyruvate kinase A Covalent Modification “Enzyme” :



- Reaction number 10 in glycolysis it's **irreversible** reaction. once glucagon (hormone) bind to the receptor it convert the enzyme pyruvate kinase A from active form to an inactive form by phosphorylation .
 ⊕ **Activation of enzyme can be done by phosphatase (removes a phosphate group)**
- When it's in the active form It catalyzes the transfer of a phosphate group from phosphoenolpyruvate (PEP) to adenosine diphosphate (ADP), yielding one molecule of pyruvate and one molecule of ATP.

Pyruvate kinase deficiency: hemolytic anemia


- Pyruvate kinase deficiency is due to genetic mutation (it affects the survival of red blood cells).
- ◆ The enzyme may show an abnormal K_m or V_{max} for substrates or coenzymes.
- ◆ The enzyme may show an abnormal response to the activator fructose 1,6 - bisphosphate.
- ◆ Enzyme activity or stability may be altered , or the amount of enzyme may be decreased .
- Pyruvate kinase mutation may lead to :
 1. Altered enzyme Kinetics . (mutation in the **allosteric binding site** and its goal to inhibit enzyme activity).
 2. Altered response to activator . (mutation in the **active site** and its goal to stop enzyme activity).
 3. Altered enzyme stability.
 4. Decreased amount of the enzyme.




Regulation of enzymes

“ long term regulation “ hormonal “


1



3



10



- Long term regulation of glycolysis **in reactions 1 , 3 and 10** :

1

Glucagon: **Repression** .
- inhibits the glycolysis .

2

Insulin: **Induction** .
- Activities the uptake of glucose (GLUT 4).

- Regulatory Mechanisms:

- Rapid, short-term: **Allosteric, Covalent modifications** are the enzymes regulation in the three irreversible steps.
- Slow, long-term: **Induction/repression** (insulin and glucagon respectively).

Substrate-level phosphorylation Vs. Oxidative phosphorylation


- Phosphorylation is the metabolic reaction of introducing a **phosphate group** into an **organic molecule**.

Oxidative phosphorylation	Substrate-level phosphorylation
<p>“Production of ATP molecules from the redox reactions of an electron transport chain”.</p> <p>Formation of high-energy phosphate bonds by phosphorylation ADP to ATP.</p>	<p>“ Production of ATP molecules via transfer of a phosphate group from an intermediate high-energy substrate directly to ADP”.</p> <p>Formation of high-energy phosphate bonds by phosphorylation ADP to ATP (or GDP to GTP).</p>
<p>Coupled to the transfer of electrons from reduced coenzyme to molecular oxygen via ETC</p>	<p>Coupled to cleavage of a high-energy metabolic intermediate (substrate)</p>
<p>In mitochondria (ETC).</p>	<p>In cytosol or mitochondria (glycolysis and Krebs cycle).</p>

Aerobic glycolysis : ATP production

- Phosphorylation is the metabolic reaction of introducing a **phosphate group** into an **organic molecule**.

ATP Consumption	<p>Reaction 1 + Reaction 3 " From glycolysis steps" 2 ATP</p>
NADH Consumption	Conversion of pyruvate to lactate
NADH Production	<p>Reaction 6 " From glycolysis steps" 2 NADH " 1 NADH means 3 ATP" 2 X 3 = 6 ATP " From ETC (oxidative level)" AND Reaction 7 + Reaction 9 " From substrate level (glycolysis steps)" Keep in mind these 2 reactions was multiplied by 2 2 ATP X 2 = 4 ATP</p>
ATP Production	
Net	10 - 2 + 8 ATP



Aerobic glycolysis : ATP production


ATP consumed	2 ATP	
ATP produced	Substrate-level (directly)	2 ATP x 2 = 4 ATP
	Oxidative-level (in ETC)	2 NADH x 3 = 6 ATP
Total	10	
Net	10 " produced " - 2 " consumed " = 8 ATP	

Anaerobic glycolysis


- Anaerobic glycolysis is the transformation of glucose to lactate when limited amounts of oxygen (O_2) are available. Anaerobic glycolysis is only an effective means of energy production during short, intense exercise, providing energy for a period ranging from 10 seconds to 2 minutes.
- NADH is produced cannot be used by ETC for ATP production (No O_2 and/ or no mitochondria).
- Less ATP production, as compared to aerobic glycolysis.
- The end product of anaerobic glycolysis is **lactate** "**obligatory output**". (why?) Because if it is not formed, All cellular NAD^+ will be converted to NADH, with no means to replenish (fill again) the cellular NAD^+ Glycolysis stops death of the cell.
- Anaerobic Glycolysis is important in RBCs because they don't have mitochondria .

Anaerobic enzyme : Lactate dehydrogenase

11



- Reversible reaction.
- **Lactate dehydrogenase** enzyme is used for both direction .
- **2x** $NADH + H^+$ in .
- **2x** NAD^+ out .



★ To summarize

Anaerobic glycolysis : ATP production

ATP consumed	2 ATP	
ATP produced	Substrate-level (directly)	2 ATP x 2 = 4 ATP
	Oxidative-level (in ETC) ماقدروا يطلقون من الـ NADH طاقة لأنه ما دخل الـ ETC NADH can <u>NOT</u> be used by ETC (oxidative-level) because: there is no O ₂ and/or no mitochondria.	2 NADH X 3 = 6 ATP
Total	4	
Net	4 " produced " - 2 " consumed " = 2 ATP	

Anaerobic glycolysis in RBC (2,3 BPG shunt)




- **Mutase enzyme** It is important for association and dissociation between O_2 and hemoglobin.
- Increase in "2,3-BPG" will help with loss of association between O_2 and hemoglobin and will release more O_2 .
- It usually occurs with people who live in high altitude.



- H_2O in .
- P_i out .

- Shunt = diversion (تحويل):
 - 2,3-BPG " 2,3-bis-phosphoglycerate " Shunt is responsible for loading and unloading of oxygen .
 - No ATP production .
 - Present in both fetal and adult hemoglobin .
 - in fetal hemoglobin, it interacts less with hemoglobin .
 - Presence of 2,3-BPG shunt reduces hemoglobin's oxygen affinity **i.e.** fetal hemoglobin has higher affinity to oxygen than adult hemoglobin.



Anaerobic glycolysis in RBC (2,3 BPG shunt) : ATP production

ATP consumed	2 ATP	
ATP produced	Substrate-level (directly)	2 ATP x 2 = 4 ATP (without shunt) 1 ATP x 2 = 2 ATP (with shunt)
	Oxidative-level (in ETC) ماقدروا يطلعون من الـ NADH طاقة لانه ما دخل الـ ETC NADH can <u>NOT</u> be used by ETC (oxidative-level) because: there is no O ₂ and/or no mitochondria.	2 NADH x 3 = 6 ATP
Total	4 or 2	
Net	4 " produced " - 2 " consumed " = 2 ATP (without shunt) 2 " produced " - 2 " consumed " = 0 ATP (with shunt)	

Anaerobic glycolysis in RBC (2,3 BPG shunt)

“ summary “

End product	<ol style="list-style-type: none">1. Lactate .2. No net production or consumption of NADH .
Energy yield	<ul style="list-style-type: none">- If no shunt is formed = 2 ATP .- If shunt occurs = 0 ATP .
PK deficiency hemolytic anemia depends on	<ul style="list-style-type: none">- Degree of PK deficiency.- Compensation 2,3-BPG

Take home messages



Glycolysis is the major oxidative pathway for glucose.



Glycolysis is employed by all tissues.



Glycolysis is a tightly-regulated pathway.



PFK-1 is the rate-limiting regulatory enzyme.



Glycolysis is mainly a catabolic pathway for ATP production, but it has some anabolic features (amphibolic).



Pyruvate kinase deficiency in RBCs results in hemolytic anemia.

Quiz

Q1 : Oxidative phosphorylation happens in

- | | | | |
|-------------|-------------|------------------|--------------|
| A) Nucleus | B) Cytosol | C) Mitochondria | D) Rough ER |
|-------------|-------------|------------------|--------------|

Q2 : The net ATP production in Aerobic Glycolysis is

- | | | | |
|-----------|-----------|-----------|-----------|
| A) 8 ATP | B) 4 ATP | C) 7 ATP | D) 5 ATP |
|-----------|-----------|-----------|-----------|

Q3 : Which of the following biochemical pathways does NOT require oxygen ?

- | | | | |
|-----------------|----------------|---------|------------------|
| A) Krebs cycle | B) Glycolysis | C) ETC | D) Calvin cycle |
|-----------------|----------------|---------|------------------|

Q4 : The final product of Anaerobic Glycolysis is

- | | | | |
|--------------|-------------|-------------|-----------------------|
| A) Pyruvate | B) Lactate | C) Citrate | D) Acetyl coenzyme A |
|--------------|-------------|-------------|-----------------------|

Q5 : There are steps reaction in aerobic glycolysis

- | | | | |
|--------|--------|--------|--------|
| A) 13 | B) 20 | C) 10 | D) 15 |
|--------|--------|--------|--------|

Q6 : The first 3 steps in glycolysis are

- | | | | |
|------------------|----------------|------------------|------------------|
| A) Irreversible | B) Reversible | C) Both of them | D) None of them |
|------------------|----------------|------------------|------------------|

SAQs :

Q1: What is the function of enolase enzyme ?

Q2: How many NADH are produced by glycolysis per glucose ?

★ MCQs Answer key:

1) C 2) A 3) B 4) B 5) c 6) A

★ SAQs Answer key:

- 1) Remove water, so it will convert phosphoglycerat to Phosopheonolpyruvate
- 2) 2 NADH



Girls team:

Alia Zawawi
Nada Babilli

Rania Aqil
Reem alamri

Reema Alomar
Reem Alqahtani

Renad Alhumaidi
Shaden Alobaid

Noura Alsalem
Lama Alahmadi

Sadem Alhazmi
Somow Abdulrahman

Budoor Almubarak
Samar Almohammedi

Nuha Alkudsi
Norah Alsheikh
Muneerah Alssdhan
Mayasem Alhazmi
Noura alshathri
Duaa Alhumoudi



Boys team:

Mansour albawardi

📍 Hassan alshuraf

📍 Abdulrahman almbki

Mohammed alsayari

Abdullaziz alomar


Abdulaziz alrabiah

Saud alrasheed

Abdullah almazro

Hamad almousa

Ahmad alkhayat



A goal should scare you a little and excite you a lot!.

📍 Shatha Aldhohair

📍 Mishal Althunayan

Made by 📍



Bio Chem 439



Biochemistry439@gmail.com



@Biochemistry439