MUSCLES OF BACK

Department of Anatomy King Saud University



Dr. Sameerah Shaheen Assistant Professor of Anatomy College of Medicine sshahee@ksu.edu.sa

OBJECTIVES

At the end of the lecture, students should be able to:

- Distinguish between the different groups of back muscles.
- Compare between groups of back muscles as regard their <u>nerve supply</u> and <u>action</u>.
- List the back <u>muscles of each group</u>.
- Describe the <u>attachments</u> of each muscle of the superficial group, as well as, its <u>nerve supply</u> and action.
- Describe the triangles of back and their clinical significance.

BACK MUSCLES

They are organized into 3 groups:

Deep group: attached to & involved in the movement of vertebral column & head.

□ Intrinsic muscles:

☐ Develop in the back

☐ Supplied by posterior rami of spinal nerves

Superficial group: attached to & involved in the movements of upper limb (shoulder).

Intermediate group: attached to ribs & associated with of the thoracic cage movements.

Serve respiratory functions.

□ Extrinsic muscles:


☐ Not developed in the back.

☐ Supplied by anterior rami of spinal nerves.

DEEP GROUP OF BACK MUSCLES


- ☐ They extend from sacrum to skull.
- They include <u>extensors</u> and <u>rotators</u> of head & vertebral column. So, It is a set of muscles that straighten and rotate the back.
- ☐ Their tone is <u>responsible for</u> maintenance of normal curvature of vertebral column.
- The largest muscle of this group is "erector spinae" which is formed of 3 vertical columns (from lateral to medial: iliocostalis, longissimus & spinalis.

(Note the length and attachment of the muscle fibers)



INTERMEDIATE GROUP OF BACK MUSCLES

- ☐ It is separated from the deep group by <u>thoracolumbar</u> <u>fascia</u>.
- ☐ It includes:
 - Serratus posterior superior (rib elevator).
 - Serratus posterior inferior (rib depressor).
- Nerve supply: anterior rami of thoracic spinal nerves (intercostal nerve).
- Serratus posterior superior contributes in deep inspiration.
- Serratus posterior inferior contributes in <u>forced expiration</u>.




SUPRERFICIAL GROUP OF BACK MUSCLES

Includes two sets of muscles:

They originate from the vertebral column and attach to the bones of the shoulder – the clavicle, scapula and humerus.

- Muscles connecting vertebral column to scapula (move scapula through shoulder girdle joints) & include:
 - 1. Trapezius.
 - 2. Levator scapulae.
 - 3. Rhomboid minor.
 - 4. Rhomboid major.
- Muscle connecting vertebral column to humerus (move humerus through shoulder joint), & include:
 - Latissimus dorsi.



TRAPEZIUS

- Origin: Skull, ligamentum nuchae & spinous processes of cervical & thoracic vertebrae (C7-T12).
- Insertion: lateral 1/3 of clavicle, acromion& spine of scapula.
- ☐ Action:
 - Upper fibers: elevates scapula & rotates it during abduction of the arm.
 - 2. Middle fibers: retract scapula
 - 3. Lower fibers: depress scapula.
- Nerve supply: Motor innervation is the accessory (11th cranial) nerve. Also, proprioceptor fibers from C3 and C4 spinal nerves.

To test the accessory nerve, trapezius function can be assessed by shoulders shrug




Latissimus dors

LEVATOR SCAPULAE RHOMBOID MINOR & MAJOR


☐ Origin:

- 1. Levator scapulae: cervical transverse processes C1-C4, Insertion: medial border of scapula.
- 2. Rhomboid major: thoracic spinous processes of T2-T5 vertebrae, Insertion: medial border of the scapula, between the scapula spine and inferior angle
- 3. Rhomboid minor: the spinous processes of C7-T1 vertebrae, Insertion: medial border of scapula at the level of the spine of scapula.
- Nerve supply: dorsal scapular nerve. Actions:
- 1. Levator scapulae: elevates scapula.
- 2. Rhomboid minor & major: retract & rotate scapula.



LATISSIMUS DORSI

- ☐ Origin:
- Spinous processes of T6-T12
- Iliac crest
- Thoracolumbar fascia
- inferior 3 or 4 ribs
- Insertion: tendon attaches to the intertubercular sulcus of the humerus.
- Nerve supply: thoracodorsal nerve. (C6,7,8) From posterior cord of brachial plexus
- Actions: extension, adduction & medial rotation of upper limb.
- ☐ It is also called the climbing muscle.



☐ Auscultatory Triangle:



- **Boundaries:** latissimus dorsi, trapezius, and medial border of scapula.
- Site where breath sounds are most easily heard with a stethoscope.


□Lumbar Triangle:

(Triangle of Petit)



- **Boundaries:** latissimus dorsi, posterior border of external oblique muscle of the abdomen, and iliac crest.
- Site of an abdominal hernia; or where pus may emerge from the abdominal wall in extraabdominal lumbar abscess.

MUSCULAR TRIANGLES OF BACK



SUMMARY

□ BACK MUSCLES:

- 1. <u>Deep group</u>: attached to & moves vertebral column, supplied by posterior rami of spinal nerves.
- 2. <u>Intermediate group</u>: attached to & moves ribs, supplied by anterior rami of spinal nerves.
- 3. Superficial group:
 - Origin: vertebral column.
 - Insertion: scapula (EXCEPT latissimus dorsi: humerus).
 - <u>Action</u>: moves scapula (<u>EXCEPT</u> latissimus dorsi: moves humerus).
 - Nerve supply: anterior rami of spinal nerves through brachial plexus (EXCEPT trapezius: 11th cranial nerve).



QUESTION 1

- □Which one of the following muscles of back that <u>rotates the humerus medially</u>?
- 1. Trapezius.
- 2. Latissimus dorsi.



- 3. Rhomboid major.
- 4. Serratus posterior superior.

QUESTION 2

- □ Regarding <u>back muscles</u>, which one of the following statements is <u>correct</u>?
- 1. All back muscles are supplied by posterior rami of spinal nerves.
- 2. Muscles of intermediate group move vertebral column.
- 3. Muscles of superficial group are involved in upper limb movements.
- 4. Muscles of deep group serve respiratory functions.