


Physiology of Bone



Dr. Aida Korish, PhD

Physiology Department, KSU

Objectives

2

By the end of the lecture you will be able to:


- 1- Define bone and differentiate cortical & trabecular bone (sites and function of each).
- 2- Identify the bone cells and the function of each.
- 3- Define bone remodelling and explain the mechanism of bone formation.
- 4-State the normal levels and forms of Ca^{++} in the ECF and its relation to PO_4 .
- 5- Interpret the importance of the exchangeable calcium.
- 6- Discuss the effect of different hormones on calcium homeostasis.
- 7- Define osteoporosis and state its causes.

Physiology of Bone

3

- Bone is a special form of connective tissue.
- It is well vascularized with total blood flow of 200–400 mL/min in adult humans.
- The ends of each long bone (**epiphyses**) are separated from the shaft of the bone by a plate of actively proliferating cartilage, the **epiphyseal plate**.
- Linear bone growth can occur as long as the epiphyses are separated from the shaft of the bone, but such growth ceases after the epiphyses unite with the shaft (**epiphyseal closure**) at puberty.


Dr.Aida Korish akorish@ksu.edu.sa



Functions of bone



4

- Is involved in the overall Ca^{++} and PO_4^- homeostasis.
- Protects the vital organs.
- Permits locomotion and support against gravity.
- Contains the bone marrow (blood cells formation).



Types and structure of bone

5






Types and structure of bone

6

Compact or cortical bone: in the outer layer of most bones is (80%) of the bones in the body.

Trabecular or spongy bone inside the cortical bone, is 20% of the body bone.

- In compact bone, the bone cells lie in lacunae. They receive nutrients by way of canaliculi from **haversian canals** vessels.
- Collagen is arranged in concentric layers, around the haversian canals forming cylinders called **osteons** or **haversian systems**.
- Trabecular bone is made up of spicules or plates. Nutrients diffuse from bone extracellular fluid (ECF) into the trabeculae.



Composition of Compact Bone

7

A- Matrix

(30%) is organic Matrix : composed of


a-Collagen fibers 90-95%: extend primarily along the lines of tensional force and give bone its powerful tensile strength.

b- Ground substance 5-10% of ECF and proteoglycans, hyaluronic acid

(70%) is inorganic bone salts :

- ❑ **Crystalline salts of Ca^{++} & PO_4 (Hydroxyapatite) the Ca/P ratio is 1.5-2).**
- ❑ **Mg^+ , Na^+ , K^+ , Carbonate ions are also present.**


NB: newly formed bone have a considerably higher percentage of organic matrix in relation to salts.



B- Bone cells

8


- **Osteoblasts** are the bone forming cells that secrete collagen forming a matrix around themselves which then calcifies.
- Osteoblasts regulate Ca and Phosphate concentration in bone fluid.
- When the osteoblasts got surrounded by calcified matrix, they are called **Osteocytes** and send processes into the canaliculi that ramify throughout the bone.
- **Osteoclasts** are multinuclear cells that erode and resorb previously formed bone. They phagocytose bone, digesting it in their cytoplasm.



Mechanism of Bone Calcification.

9

- **Osteoblasts** secrete *collagen* (monomers) and *ground substance* (proteoglycans).
- The collagen monomers polymerize to collagen fibers.
- The resultant tissue becomes **osteoid**, a cartilage-like material differing from cartilage in that calcium salts readily precipitate in it.
- Osteoblasts become entrapped in the osteoid and are now called **osteocytes**.
- After the osteoid is formed, calcium salts begin to precipitate on the collagen fibers forming the **hydroxyapatite crystals**.



Tensile and Compressional Strength of Bone

10



- The collagen fibers of bone, like those of tendons, have great tensile strength, whereas the calcium salts have great compressional strength.
- These combined properties plus the degree of bondage between the collagen fibers and the crystals provide a bony structure that has both extreme tensile strength and extreme compressional strength.
- **N.B:** hydroxyapatite crystals fail to be formed in normal tissues except in bone despite the high levels of Ca & P ions due to the presence of an inhibitor of precipitation called *pyrophosphate*.

Deposition and Absorption of Bone

Remodeling of Bone

11


- Bone is continually deposited by ***osteoblasts***, and absorbed where ***osteoclasts*** are active.
- Osteoblasts are found on the outer surfaces of the bones and in the bone cavities.
- A small amount of osteoblastic activity occurs on about 4% of all bone surfaces at any given time in an adult), so that at least some new bone is being formed constantly.
- The renewal rate is about 4% per year for compact bone and 20% per year for trabecular bone.



Bone remodeling

12

The bone remodelling process



Download from
Dreamstime.com

This watermarked comp image is for previewing purposes only.

ID 61582324



© Designua | Dreamstime.com

Cont.....Bone resorption

13

- **Osteoclasts** are large phagocytic multinucleated cells
- They are normally active on less than 1% of the bone surfaces of an adult.
- The osteoclasts secrete two types of substances:
 - (1) proteolytic enzymes from the lysosomes
 - (2) several acids from the mitochondria and secretory vesicles.
- The enzymes dissolve the organic matrix, and the acids cause solution of the bone salts.


The osteoclastic cells also phagocytose minute particles of bone matrix and crystals, dissolve them and release the products into the blood.



Importance of Continual Bone Remodeling

14

- 1- **Bone adjusts its strength** in proportion to the degree of **bone stress** and it thickens when subjected to heavy loads.
 - 2- The **shape of the bone** can be rearranged for **proper support** of mechanical forces.
 - 3- Because **old bone** becomes relatively brittle and weak, **new organic matrix** is needed to maintain the normal toughness of bone .
- The **bones of children** are less brittle due to more remodeling in the children.



Control of the Rate of Bone Deposition by Bone “Stress”


15

- ❑ Bone is **deposited** in proportion to the **load** that it must carry.
- ❑ Continual **physical stress** stimulates osteoblastic **deposition** and **calcification** of bone.
- ❑ The bones of **athletes** become considerably **heavier** than those of non athletes. Also, the bone of the leg in the cast becomes thin and up to 30 % decalcified within a few weeks.
- ❑ **Bone stress** also determines the **shape of bones** under certain circumstances. (e.g. Healing of fractures may start angulated in children then become straight).

Repair of a Fracture Activates Osteoblasts

16

- ❑ **Fracture** of a bone activates all the periosteal and intraosseous **osteoblasts** involved in the break.
- ❑ Large numbers of new osteoblasts are formed from *osteoprogenitor cells*, which are bone stem cells in the surface tissue lining bone, called the "bone membrane.
- ❑ Shortly a large bulge of osteoblastic tissue and new organic bone matrix, develops between the two broken bone ends followed shortly by the deposition of calcium salts. This is called a *callus*.



Body Calcium levels


17

- 1.5% of body weight is Calcium,
- about 1100 -1300 g.
- 99% is in the skeleton.

TABLE 36.1

Body Content and Tissue Distribution of Calcium and Phosphorus in a Healthy Adult

	Calcium	Phosphorus
Total Body Content	1,300 g	600 g
Relative Tissue Distribution (% of total body content)		
Bones and teeth	99%	86%
Extracellular fluid	0.1%	0.08%
Intracellular fluid	1.0%	14%





Plasma calcium

Plasma calcium level: (9 -11 mg/dl) average :9.4 mg/dl

59% (diffusible)= Ionized + Complexed

41% (non diffusible)= protein bound

Serum calcium and Phosphate



© Elsevier. Costanzo: Physiology 3E www.studentconsult.com

[Add to My Slides](#) [Go to My Slides](#)

Figure 9-34 Ca^{2+} homeostasis in an adult eating 1000 mg/day of elemental Ca^{2+} . Hormonal effects on Ca^{2+} absorption from the gastrointestinal tract, bone remodeling, and Ca^{2+} reabsorption in the kidney are shown. PTH, Parathyroid hormone.

Calcium Exchange Between Bone and ECF

20


- The bone contains a type of *exchangeable* calcium that is always in equilibrium with the Ca^{++} ions in the ECF
- It normally amounts to about (0.4-1%) of the total bone calcium.
- This calcium is a form of readily mobilizable salt such as dicalcium phosphate (CaHPO_4) and other amorphous calcium salts.
- The importance of exchangeable calcium is that it provides a rapid *buffering* mechanism to keep the Ca^{++} ions concentration in ECF from rising to excessive levels or falling to very low levels under transient conditions of excess or decreased availability of calcium.

Hormonal Control of Calcium Metabolism & Physiology of Bone

21

Three major hormones are concerned :


- **1, 25 dihydroxycholecalciferol**: a steroid hormone formed from Vitamin D.
- **Parathyroid hormone (PTH)**: secreted by parathyroid gland
- **Calcitonin**: secreted by c-cells in the thyroid gland.
- To a lesser extent ; Glucocorticoids, GH, estrogens & various growth factors also affect Calcium Metabolism.



Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.

Hormonal control of plasma calcium


22



Vitamin D

23

VITAMIN D – ACTION OF CALCITRIOL



➤ Increases the intestinal absorption of calcium and phosphate by increased synthesis of calcium binding protein (calbinding D28k)



➤ Mineralization of bone at low doses
 ➤ Mobilization of calcium from bone at high doses




➤ Increased reabsorption of calcium and phosphorus
 ➤ Decreased excretion of calcium and phosphorus

Osteoporosis

24

- ❑ **Osteoporosis:** Is caused by a relative excess of osteoclastic function. Loss of bone matrix is marked. Matrix and mineral are both lost and there is a loss of bone mass. Due to :
- ❑ Lack of physical stress
- ❑ Malnutrition, lack of vitamin C
- ❑ Old age, Postmenopausal lack of estrogen.
- ❑ Cushing's syndrome.

Dr.Aida Korish
akorish@ksu.edu.sa



A B
● Difference between normal bone (A) and osteoporotic bone (B).




FIGURE 21-12 Total body calcium, an index of bone mass, at various ages in men and women. Note the rapid increase to young adult levels (phase I) followed by the steady loss of bone with advancing age in both sexes (phase III) and the superimposed rapid loss in women after menopause (phase II). (Reproduced with permission from Evans TG, Williams TF (eds): Oxford Textbook of Geriatric Medicine. Oxford University Press; 1992.)

Complications of Osteoporosis

25

- ❑ The incidence of fractures is increased particularly in the **distal forearm** (Cole's fracture), vertebral body, and hip. These areas have a high content of trabecular bone, which is more active metabolically, it is lost more rapidly.
- ❑ Fractures of the **vertebrae** with kyphosis produces “widow’s hump” in elderly women with osteoporosis.
- ❑ Fractures of the **hip** in elderly are associated with a mortality rate of 12–20%, and half of those who survive require prolonged expensive care.
- ❑ Increased **intake of calcium** and **moderate exercise** may help prevent or slow the progress of osteoporosis.



THANK YOU

FOR LISTENING