



AXILLARY & MEDIAN NERVES






Objectives

By the end of the lecture, you should be able to:

- Describe the **origin**, **course**, relations, **branches** and **distribution** of the axillary & median nerves.
- Describe the common causes, signs and effect of lesion to the **axillary** and **median** nerves.

Axillary Nerve

- **Origin:** (C 5 & C 6).
- Posterior cord of brachial plexus.
- **Course:**
- It passes downward and laterally along the posterior wall of the axilla, then it exit the axilla.
- Then, it passes posteriorly **around the surgical neck of the humerus**.
- It is accompanied by **posterior circumflex humeral vessels**.
- **Branches:**
- **Motor** to:
 1. **Deltoid muscle,**
 2. **Teres minor muscle.**
- **Sensory:**
- **Upper lateral cutaneous nerve of arm** which carries sensations from the skin above the deltoid muscle.



Axillary Nerve Lesion

- The axillary nerve is commonly injured due to:


1. Fracture of surgical neck of the humerus.
2. Inferior dislocation of the shoulder joint.
3. Compression from incorrect use of crutches.



Axillary Nerve Lesion


Affects:

- **Motor:**
 - Paralysis of the deltoid and teres minor muscles.
 - Impaired abduction of the shoulder (15 to 90°).
 - The paralyzed deltoid wastes rapidly.
 - As the deltoid atrophies, the rounded contour of the shoulder is lost and becomes flattened compared to the uninjured side.
- **Sensory:** Loss of sensation over the lateral side of the proximal part of the arm.




Median Nerve

- **Root value:**
- **(C 5, 6, 7, 8, and T 1).**
- The median nerve is formed **anterior to the third part of the axillary artery** by the **union of lateral and medial roots.**
- The lateral root (C5,6 &7), arises from the lateral cord of the brachial plexus.
- The medial root (C8 & T1), arises from the medial cord of the brachial plexus.



- It enters the arm from the axilla at the inferior margin of the **teres major** muscle.
- It passes vertically along the medial side of the arm and is related to the **brachial artery** throughout its course:
 - In upper ½ of the arm, it lies lateral to the brachial artery;
 - In the middle of the arm, it crosses the artery from lateral to medial;
 - In the lower ½ it descends on the medial side of the brachial artery.
- It descends anterior to the elbow joint.


Median Nerve in the Arm



The median nerve has **no major branches in the arm.**


Median Nerve in the Forearm

- Median nerve passes into the forearm anterior to elbow joint, between the **2 heads** of pronator teres.
- It innervates **most of the muscles** of the anterior compartment of the forearm, (6.5 muscles).
- (Except the flexor carpi ulnaris, and medial half of the flexor digitorum profundus, which are supplied by **ulnar nerve**).



Median Nerve in the Hand

- The median nerve continues into the hand by passing deep to **flexor retinaculum**.
- It innervates:
 - 3 muscles of the thenar eminence, (thumb), (abductor pollicis brevis, flexor pollicis brevis and opponens pollicis).
 - Lateral 2 lumbrical muscles associated with movement of the index and middle fingers; and
 - Skin over the palmar surface of the lateral three and one-half fingers.
 - Skin of the lateral 2/3rd of the palm of the hand.



Median Nerve Lesion

- Injury of the median nerve at different levels cause different syndromes.
- In the arm and forearm the median nerve is usually not injured by trauma because of its relatively deep position.
- **Median nerve can be damaged:**
 1. In the elbow region, (supracondylar fracture of the humerus).
 2. At the wrist above the flexor retinaculum.
 3. In the carpal tunnel.




- The most serious disability of median nerve injuries is:

- **Loss of opposition of the thumb.**


The delicate **pincer-like** action is not possible

- **Loss of sensation from lateral 3 ½ fingers & lateral ⅔ of the palm.**





Median Nerve Lesion in the Elbow Region

- Damaged in **supracondylar fracture** of humerus.
- **Muscles affected are:**
 - Pronator muscles of the forearm.
 - All long flexors of the wrist and fingers, (except flexor carpi ulnaris and medial half of flexor digitorum profundus).



- **Motor:**
- Loss of **pronation**. Hand is kept in supine position
- Wrist shows weak flexion, with ulnar deviation
- **No** flexion possible on the interphalangeal joints of the index and middle fingers
- **Weak** flexion of ring and little finger.
- **Thumb** is adducted and **loss of flexion** of terminal phalanx and **loss of opposition**
- **Wasting** of thenar eminence
- Hand looks flattened and ***“apelite”***, and presents an inability to flex the three most radial digits when asked to make a fist.

Wasting of thenar eminence



- Sensory:
- Loss of sensation from:
 - The radial 2/3rd of the palm.
 - Palmer aspect of the lateral 3½ fingers.
 - Distal part of the dorsal surface of the lateral 3½ fingers.
- Trophic Changes:
 - Dry and scaly skin.
 - Easily cracking nails.
 - Atrophy of the pulp of the fingers.




Median Nerve Lesion at the Wrist

- Often injured by penetrating wounds (stab wounds or broken glass) of the forearm.
- Motor:
 - Thenar muscles are paralyzed and atrophy, so that the thenar eminence becomes flattened.
 - Opposition and abduction of thumb are lost, and thumb and lateral two fingers are arrested in adduction and hyperextension position. “**Apelike hand**”.
- Sensory & trophic changes are the same as in the elbow region injuries.

Carpal Tunnel Syndrome

- The most common neurological problem associated with the median nerve is compression beneath the flexor retinaculum at the wrist.
- **Motor:**
- Weak motor function of thumb, index & middle finger.
- **Sensory:**
- Burning pain, '**pins and needles**' along the distribution of median nerve to the lateral 3½ fingers.




NB. **No** sensory changes over the palm as the **palmer cutaneous branch of the median nerve** is given before the median nerve enters the carpal tunnel.

Summary

- **Axillary Nerve**
- **Origin:**
Posterior cord.
- Spinal segments:
- C5, & C6.
- **Function:**
 - **Motor:**
 - Deltoid, &
 - Teres minor.
 - **Sensory:**
 - Skin over upper lateral part of arm.

- **Median Nerve**
- **Origin:** Medial and lateral cords.
- Spinal segments: (C5, C6, C7, C8 & T1).
- **Function:**
 - **Motor**
All muscles in the anterior compartment of the forearm (except flexor carpi ulnaris and medial half of flexor digitorum profundus), three thenar muscles of the thumb and two lateral lumbrical muscles.
 - **Sensory**
Skin over the palmar surface of the lateral three and one-half digits and over the lateral 2/3rd of the palm of the hand.



**Thank You
&
Good Luck**