

Development of Respiratory System

*BY : Dr.SANA
ALSHAARAWY*

OBJECTIVES

At the end of the lecture the students should be able to:

- Identify the development of the laryngeotracheal (respiratory) diverticulum.
- Identify the development of the larynx.
- Identify the development of the trachea.
- Identify the development of the bronchi & Lungs.
- Describe the periods of the maturation of the lung.
- Identify the most congenital anomaly.


Respiratory System

■ Upper respiratory tract:

- ◆ Nose
- ◆ Nasal cavity & paranasal sinuses
- ◆ Laryngo-pharynx
- ◆ Larynx.


■ Lower respiratory tract:

- ◆ Trachea
- ◆ Bronchi
- ◆ Lungs




Development of the Respiratory Tract

- **Begins** during the 4th week of development.
- **Begins as** a median outgrowth (**laryngo-tracheal groove**) from the **caudal part** of the **ventral wall** of the **primitive pharynx** (**foregut**)
- The groove **envaginates** and forms the **laryngotracheal (respiratory) diverticulum**.




■ A longitudinal **tracheo-esophageal septum** develops and divides the diverticulum into a:

- ◆ **Dorsal portion:** primordium of the oropharynx and esophagus
- ◆ **Ventral portion:** primordium of Respiratory Diverticulum (larynx, trachea, bronchi and lungs).



- **The proximal part** of the **respiratory diverticulum** remains tubular and forms **larynx & trachea.**
- **The distal end** of the diverticulum dilates to form **lung bud**, which **divides** to give rise to **2 lung buds (primary bronchial buds)**




- The **endoderm** lining the **laryngotracheal diverticulum (Respiratory diverticulum)**
gives rise to the:
 - ◆ **Epithelium & Glands** of the respiratory tract.

- The **surrounding splanchnic mesoderm**
gives rise to the:
 - ◆ **Connective tissue, Cartilage & Smooth muscles** of the respiratory tract


Development of the Larynx

- The opening of the laryngotracheal diverticulum into the primitive foregut becomes the laryngeal orifice.
- The epithelium & glands are derived from endoderm.
- Laryngeal muscles & the cartilages of the larynx except Epiglottis develop from the mesoderm of 4th & 6th pairs of pharyngeal arches.



Epiglottis


- It develops from the **caudal part** of the **hypopharyngeal eminence**, a swelling formed by the **proliferation of mesoderm in the floor of the pharynx.**



- Growth of the **larynx and epiglottis is rapid** during **the first three years** after birth. By this time the epiglottis has reached its adult form.



Recanalization of larynx

- The **laryngeal epithelium** proliferates rapidly resulting in **temporary occlusion** of the laryngeal lumen
- **Recanalization** of larynx normally occurs by the **10th week**.
- Laryngeal **ventricles**, **vocal folds** and **vestibular folds** are formed during **recanalization**.





Development of the Trachea

- The **endodermal lining** of the **laryngotracheal tube distal to the larynx** differentiates into the **epithelium and glands** of the trachea and pulmonary epithelium
- The **cartilages, connective tissue, and muscles** of the trachea are derived from the mesoderm.





Development of the Bronchi & Lungs


- **The 2 primary bronchial buds** grow laterally into the **pericardio-peritoneal canals** which the **primordia of pleural cavities**
- **Bronchial buds** divide and re-divide to give the **bronchial tree**.



- **The right main bronchus is slightly shorter & wider than the left one and is oriented more vertically**
- **The embryonic relationship persists in the adult.**
- **The main bronchi subdivide into secondary and tertiary (segmental) bronchi which give rise to further branches.**






- **The segmental bronchi** 10 in right lung and 8 or 9 in the left lung begin to form by the **7th week**
- The surrounding mesenchyme also divides.
- **Each segmental bronchus with its surrounding mass of mesenchyme is the primordium of a bronchopulmonary segment.**





Development of the pleura

- As the lungs develop they acquire a layer of **visceral pleura from splanchnic mesenchyme.**
- The thoracic body wall becomes lined by a layer of **parietal pleura derived from the somatic mesoderm.**




Maturation of the Lungs

- Maturation of lung is divided into 4 periods:
 - ◆ **Pseudoglandular** (5 - 17 weeks)
 - ◆ **Canalicular** (16 - 25 weeks)
 - ◆ **Terminal sac** (24 weeks - birth)
 - ◆ **Alveolar** (late fetal period - childhood)




Pseudoglandular Period (5-17 weeks)

- Developing lungs somewhat resembles an exocrine gland during this period.
- By **17 weeks** all major elements of the lung have formed except those involved with gas exchange (alveoli).
- Respiration is **NOT** possible.
- **Fetuses** born during this period are **unable to survive**.



Canalicular Period (16-25 weeks)



- Lung tissue becomes **highly vascular**.
- Lumina of **bronchi** and **terminal bronchioles** become larger.
- By **24 weeks** each **terminal bronchiole** has given rise to two or more **respiratory bronchioles**.
- The respiratory bronchioles divide into 3 to 6 tubular passages called **alveolar ducts**.
- Some thin-walled **terminal sacs** (**primordial alveoli**) **develop** at the end of respiratory bronchioles.



- **Respiration is possible** at the end of this period.
- Fetus born at the end of this period may survive if given intensive care (but usually die because of the immaturity of respiratory as well as other systems)

Terminal Sac Period (24 weeks - birth)

- Many more terminal sacs develop.
- Their epithelium becomes very thin forming the developing alveoli.
- Capillaries begin to bulge into developing alveoli.
- The epithelial cells of the **alveoli** and the endothelial cells of the **capillaries** come in intimate contact and establish the blood-air barrier.
- Adequate gas exchange can occur which allows the prematurely born fetus to survive






- **Surfactant** production **begins** by **24 weeks** and **increases** during the terminal stages of pregnancy.
- **Sufficient terminal sacs, pulmonary vasculature & surfactant** are present to permit survival of a prematurely born infants
- Fetuses born prematurely at 24-26 weeks may suffer from respiratory distress due to surfactant deficiency but may survive if given intensive care.

Alveolar Period (32 weeks – 8 years)

- At the beginning of the alveolar period, **each respiratory bronchiole terminates in a cluster of thin-walled terminal saccules (future alveoli)** separated from one another by loose connective tissue.
- **These terminal saccules or alveoli represent future alveolar sac.**




PRETERM LUNGS



32 WEEKS GESTATIONAL AGE


FULL-TERM LUNGS



40 WEEKS GESTATIONAL AGE

- Characteristic mature alveoli do not form until after birth, so; 95% of alveoli develop postnatally.

- About 50 million alveoli, one sixth of the adult number are present in the lungs of a **full-term newborn infant.**





- From 3-8 **year** or so, **the number of alveoli continues to increase,** forming additional primordial alveoli.
- **By** about the **eighth year,** the **adult complement of 300 million alveoli is present.**

Developmental anomalies

Tracheo-esophageal Fistula

- An **abnormal passage** between the trachea and esophagus.
- **Results from incomplete division** of the cranial part of the foregut into respiratory and esophageal parts by the **tracheo-esophageal septum**.
- Occurs once in 3000 to 4500 live births.
- Most affected infants are **males**.
- In more than **85% of cases**, the fistula is associated with esophageal atresia (esophagus ends in a blind-ended pouch rather than connecting normally to the stomach).



Thank You