Diseases of the Respiratory System

Dr. Ammar C. Al-Rikabi Dr. Maha Arafah

Respiratory block Pathology Lec 2 2020

Diseases of the Respiratory System

Chronic Obstructive Pulmonary Diseases (COPD)



Objectives:

- Give introduction for diffuse lung disease
- Compare and contrast the major clinical and functional differences between predominant chronic bronchitis versus predominant emphysema in patients with COPD
- Define Bronchiectasis, its causes, presentation, morphology and significant.

Obstructive Lung Diseases

(diffuse)

I) Bronchial Asthma

2) Chronic obstructive pulmonary disease (COPD) They are of two types:
a) Chronic bronchitis
b) Emphysema

3) Bronchiectasis

Common symptoms in lung disease

- Dyspnea: difficulty with breathing
- Cough
- Hemoptysis

Diseases of the Respiratory System

Chronic Bronchitis

Objectives:


- a. Define chronic bronchitis.
- b. Describe the causes, pathogenesis and the morphology of chronic bronchitis.
- c. Describe the mechanism of airway obstruction in a patient with chronic bronchitis.
- d. Understand that when severe obstruction is present in chronic bronchitis, significant emphysema is nearly always present

Chronic Bronchitis: defined as persistent productive cough for at least 3 consecutive months in at least 2 consecutive years

Causes


- Cigarette smoking is the most important risk factor
- Air pollutants
- Cystic fibrosis

Chronic Bronchitis: morphology




Chronic bronchitis

Chronic Bronchitis: morphology



Chronic bronchitis

Chronic Bronchitis: morphology



Chronic bronchitis. In chronic bronchitis the main abnormality is secretion of abnormal amounts of mucus, causing plugging of the airway lumen (P).

Clinical features and compilcations

- Persistent reproductive cough
- Dyspnea on exertion
- Hypercapnia, hypoxemia, cyanosis
- Emphysema
- Cor pulmonale
- Death due to further impairment of respiratory functions after superimposed acute baterial infections.

Chronic Bronchitis

Definition: Persistent productive cough (with sputum) for at least 3 months in at least 2 consecutive years

Causes

• Cigarette smoking is the most important risk factor; air pollutants also contribute

Features

 enlargement of mucous-secreting glands, goblet cell hyperplasia, chronic inflammation, and bronchiolar wall fibrosis.

Complications

- Persistent reproductive cough, dyspnea on exertion, hypercapnia, hypoxemia, cyanosis, cor pulmonale with edema (blue bloater)
- Death may result from further impairment of respiratory function due to superimposed acute infections.

Diseases of the Respiratory System

Emphysema

Objectives:

- a. Define emphysema.
- b. Describe the gross and microscopic changes in emphysema.
- c. Discuss the typical clinical presentation and causes of death.
- d. Describe the most likely mechanism of emphysema (the protease-antiprotease mechanism).
- e. Describe the pathophysiologic mechanisms of emphysema

Emphysema

Is abnormal permanent enlargement of all or part of the respiratory unit accompanied by destruction of their walls


> Associated with loss of elastic recoil and support of small airways leading to tendency to collapse with obstruction




Diseases of Lung




Emphysema




Bullous emphysema with large apical and subpleural bullae



Distal acinar /paraseptal: forming multiple cyst-like structures with spontaneous pneumothorax.



Paraseptal emphysema, microscopic destruction of alveolar walls



Pursed lip expiration is a common maneuver adopted by patients with severe chronic obstructive pulmonary disease including emphysema. The patient starts to breathe out closed or nearly closed lips to keep the intrabronchial pressure high and prevent collapse of the bronchial wall and expiratory obstruction. Later in expiration the lips are blown forwards and open, often with a grunt ("fish-mouth breathing).




Barrel-shaped chest in a patient with emphysema. The hyperinflation result from


- air-trapping with inflammatory changes
- 2. hypersecretion of viscid contraction in the small airways.

Note the associated indrawing of the intercostal muscles. Similar changes are seen in patients with chronic bronchitis and asthma.


Emphysema: **Pathogenesis**

Pathogenesis of Emphysema






Diseases of Lung




Classification of emphysema

Emphysema: morphology

Diseases of Lung




Generalized emphysema. (a) Normal distal lung acinus. (b) Centriacinar emphysema. (c) Centriacinar emphysema. (d) Panacinar emphysema. (e) Panacinar emphysema (Gough-Wentworth section).



Destruction of some alveolar septi

Emphysema: SUMMARY Emphysema:	Dilated air spaces beyond respiratory arteriols
Types	 Centriacinar: Smoking Panacinar: deficiency of α1 AT Paraseptal: Occurs adjacent to areas of fibrosis or atelectasis.
	 Irregular: scar
Clinical features	 Cough and wheezing. Respiratory acidosis Weight loss Pulmonary function tests reveal low FEV1
	Pneumothorax
Complications	 Prieumotriorax Death from emphysema is related to: Pulmonary failure with respiratory acidosis, hypoxia and coma. Pulmonary hypertension. Right-sided heart failure (Cor pulmnale)



Bronchiectasis

Objectives:

Definition

Causes


Presentation

morphology and significant

Bronchiectasis

Bronchiectasis is permenat dilatation of bronchi with destruction of their walls.

It is a result of chronic inflammation associated with an inability to clear mucoid secretions.



Causes of bronchiectasis

1. Bronchial obstruction

Localized:

 tumor, foreign bodies or mucous impaction <u>Generalized:</u>

- bronchial asthma
- chronic bronchitis


2. Congenital or hereditary conditions

- Congenital bronchiectasis
- Cystic fibrosis.
- Intralobar sequestration of the lung.
- Immunodeficiency status.
- Immotile cilia and kartagner syndrome

3. Chronic or severe infection / necrotizing pneumonia


Caused by TB, staphylococci or mixed infection

Diseases of Lung



Cilial dysmotility syndrome. Electron micrograph of cilia from a person with recurrent chest infections since childhood. The outer dynein arms are absent and there are abnormal single microtubules (M), which prevent normal motility.

Presentation




Bronchiectasis, chest radiograph



Bronchiectasis, gross

Diseases of Lung




Bronchiectasis

 Dilatation of bronchi with destruction of bronchial walls

Diseases of Lung



Bronchiectasis. This is a lower lobe of lung surgically resected for bronchiectasis.



Bronchiectasis, micropscopic

Bronchiectasis

Bronchiectasis:

Dilatation and destruction of bronchi and bronchioles secondary to chronic inflammation and obstruction

- Infection/ Necrotizing pneumonia
- Obstruction
 - **Congenital** (Cystic fibrosis, Kartagener's Syndrome)


Clinical features


Complications

Causes

- Sever persistent cough with sputum (mucopurulent sputum) sometime with blood.
- Clubbing of fingers.
- If sever, obstructive pulmonary function
- Lung Abscess
- Rare complications: metastatic brain(cerebral) abscess and amyloidosis.

Diseases of Lung





Lung abscess.

Diseases of Lung

Key Facts Chronic obstructive pulmonary disease

. Definition: a disease state characterized by airflow limitations that are not fully reversible. The airflow limitation is usually both progressive and associated with an abnormal inflammatory response of the lungs to noxious particles or gases.

. Cigarette smoking remains the most important cause of COPD. Other risks are recurrent chest viral infections in childhood, atopy, asthma, and occupational exposure to dusts (especially mining).

. Respiratory bronchiolitis is one of the earliest lesions seen in smokers.

. Chronic bronchitic airways show mucous hypersecretion with mucous gland hyperplasia.

. Chronic bronchitis and bronchiolitis cause airway narrowing.

. Emphysema causes loss of elastic recoil in lungs and contributes to functional airways obstruction.

. Generalized emphysema is defined as permanent dilatation of any part of the respiratory acinus, with destruction of tissue in the absence of scarring.

.There are two patterns of generalized emphysema: centrilobular and panacinar.

. Many patients with COPD have a reversible component to functional airways obstruction.

. Pulmonary hypertension and right-sided heart failure are common in long-standing chronic obstructive pulmonary disease.

Acute deterioration in COPD is usually caused by viral or bacterial infection.

Chronic injury (e.g., smoking)

Small airway disease

EMPHYSEMA Alveolar wall destruction Overinflation

CHRONIC BRONCHITIS Productive cough Airway inflammation

ASTHMA Reversible obstruction

Bronchial hyperresponsiveness triggered by allergens, infection, etc.

