


Effects of low and high gas pressure on the body



Dr.Thamir Al-khlaiwi
Physiology department
talkhlaiwi@KSU.EDU.SA

Objectives

By the end of this lecture you should be able to:

- 1-Describe the **effects of exposure** to low and high barometric pressures on the body.
- 2- Describe the **body acclimatization** to low barometric pressure.
- 3-Define **decompression sickness** and explain **how it can be avoided**.
- 4-Understand the effects of high nitrogen pressure, and **nitrogen narcosis**.

Effect of increased barometric pressure (Deep sea diving)

- When human descend below the sea, the pressure around them increased.
- To prevent the lungs from collapse air must be supplied also under high pressure.
- This exposes the blood in the lungs to extremely high alveolar gas pressure (hyperbarism).
- Under certain limits these high pressures cause tremendous alterations in the physiology of the body.




Figure 44-4

Open-circuit demand type of SCUBA apparatus.

Cont..

- The surrounding pressure increases by 1 atmosphere for every 10 meters (33 feet) of depth in sea water. So the diver is exposed to 2 atmospheric pressures.
- Therefore, at a depth of 31 meters (100 feet) in the ocean the diver is exposed to a pressure of 4 atmospheres.
- These problems confront SCUBA (self contained under water breathing apparatus..)



➤ **Effect of depth on the volume of the gases:**

At depth, there is compression of gases to smaller and smaller volumes. i.e 1L (sea level) → 1/2 L at 33 feet and so on.

➤ **Effect of depth on density of gases:**

There is increase in the density of gas and hence increased work of breathing.

➤ **Nitrogen effect at high nitrogen pressure:**

Nitrogen will have 2 principle effects:

- * Nitrogen narcosis (anesthetic effect)
- * Decompression sickness.




Figure 44-1
Effect of sea depth on pressure (top table) and on gas volume (bottom).

Oxygen toxicity when breathing hyperbaric air (Effect of Very High PO₂ on Blood Oxygen Transport)

When the Po₂ in the blood rises above 100 mm Hg, the amount of oxygen dissolved in the water of the blood increases markedly.

Acute Oxygen Poisoning:

The extremely high tissue Po₂ that occurs when oxygen is breathed at very high alveolar oxygen pressure can be detrimental to many of the body's tissues.

At 4 atmospheres pressure of oxygen (Po₂ = 3040 mm Hg) will cause brain *seizures followed by coma* in most people within 30 to 60 minutes.

Other symptoms include nausea, muscle twitchings, dizziness, disturbances of vision, irritability, and disorientation.




Figure 44-2

Quantity of oxygen dissolved in the fluid of the blood and in combination with hemoglobin at very high PO₂s.

Cont. Oxygen toxicity at a high PO₂.

- Molecular oxygen (O₂) has little capability of oxidizing other chemical compounds. Instead, it will first be converted into an “active” form of oxygen called: oxygen free radicals.

For example: superoxide and hydrogen peroxide.

- At high levels, these oxygen free radicals can have serious destructive and even lethal effects on the cells.

Nitrogen narcosis


- Nitrogen like most other anesthetic gases, dissolve freely in the fats of the body including the membranes and other lipid structures of the neurons (after one hour).
 - This leads to alteration of the electrical conductance of the membranes, reduces their excitability and subsequent narcosis develops.
 - At 120 feet: the diver loses many of his cares.
 - At 150 feet: there is a feeling of euphoria and drowsiness and impaired performance.
- At higher pressure: loss of coordination and finally coma might develop.

Foot	Liters
0	1
33	2
100	4
200	7
300	10



Decompression Sickness (Bends, Compressed Air Sickness, Caisson Disease, Diver's Paralysis, Dysbarism).

- ❑ During descending, the high partial pressure of nitrogen (encountered when breathing compressed air at depth) forces this gas into solution in body tissue particularly in fat (it has a high N₂ solubility).
- ❑ On ascending, this inert gas comes out of physical solution forming a gaseous phase (bubbles), leading to symptoms and signs.




Cont.. Decompression sickness

□ During slow ascending:

N₂ is slowly removed from the tissues since the partial pressure is higher than that in the arterial blood and alveolar gas.

□ If decompression is rapid:

bubbles of gaseous nitrogen are released, in tissues and blood, causing the symptoms of decompression sickness (the bends or caisson disease).





Symptoms & signs of decompression sickness

- ✓ The mildest form of DS is fatigue or drowsiness after decompression.
- ✓ Locally there is skin itch.
- ✓ other severe symptoms may occur e.g.:bubbles in the tissues cause severe pains particularly around the joints.
- ✓ Neurological symptoms including paresthesia, itching, paralysis, and inner ear disturbances.



Cont.. Decompression Sickness (S&S)

- ✓ **Thoracic pains:** dyspnea, substernal pain, cyanosis, and cough.
- ✓ **Bubbles in the coronary arteries** may cause myocardial damage.
- ✓ **Decompression sickness shock**, capillaries become permeable to plasma and hypovolemia rapidly develop.
- ✓ **Edema** may be prominent and shock is also usually complicated by pulmonary edema.



Treatment of decompression symptoms

- Rapid recompression in a pressure chamber followed by slower decompression.
- This reduces the volume of the bubbles and forces them back into solution.
- In a very deep dive, the risk of decompression sickness can be reduced if a helium-O₂ mixture is breathed during the dive.
- Also it is important to reduce the oxygen concentration in the gaseous mixture to avoid oxygen toxicity that would cause seizures.




Cont.. Treatment of Decompression sickness

- Helium is more desirable than nitrogen in deep dives because it has:
- $\frac{1}{4}$ - $\frac{1}{5}$ the narcotic effect of nitrogen on CNS.
- $\frac{1}{7}$ the molecular weight of nitrogen.
- low density leading to decreased air way resistance of diver.
- Helium is about $\frac{1}{2}$ as soluble as nitrogen in body fluids. This reduces the quantity of bubbles that can form in tissues when the diver is decompressed after diving.
- Diffuses out of the tissues during decompression several times as rapidly as does nitrogen, thus reducing the problem of decompression sickness.

Effects of low oxygen pressure on the body

(Aviation-ascend to high altitude)

- At the sea level the barometric pressure is 760 mmHg.
- At 10,000 feet is 523 mmHg
- At 50,000 feet 87 mmHg.
- This decrease in barometric pressure is the basic cause of all the hypoxia problems in high altitude.



Alveolar PO₂ at different altitudes

- As the barometric pressure decreases, the oxygen partial pressure decreases proportionally, remaining less than 21 % of the total barometric pressure.
- At sea level PO₂= 159 mmHg.
- At 20,000 feet PO₂= 40 mmHg.
- At 50,000 feet PO₂= only 18 mmHg.

Effects of acute hypoxia

- Some of the important acute effects of hypoxia beginning at an altitude of approximately 12,000 feet, are:
Drowsiness, lassitude, mental and muscle fatigue, sometimes headache, occasionally nausea and sometimes euphoria.
- All these progress to a stage of twitching or convulsions above 18,000.
- Above 23,000 feet the un acclimatized person can enter into coma.

Chronic Breathing of Low O₂ Stimulates Respiration — Phenomenon of “Acclimatization”

- Mountain climbers have found that when they ascend a mountain slowly, over a period of days rather than a period of hours, they breathe much more deeply and therefore can withstand far lower atmospheric O₂ concentrations than when they ascend rapidly. This phenomenon is called *acclimatization*.
- The reason for acclimatization is that, within 2 to 3 days, the respiratory center in the brain stem loses about four fifths of its sensitivity to changes in PCO₂ and hydrogen ions.
- It causes fewer deleterious effects on the body and it becomes possible for the person to work harder without hypoxic effects or to ascend to still higher altitude.

Principles of acclimatization

- 1- Increase in pulmonary ventilation.
- 2- Increased red blood cells.
- 3-Increased diffusing capacity of the lungs.
- 4- Increased vascularity of the tissues.
- 5-Increased ability of the cells to utilize oxygen despite the low PO₂ through increased number of mitochondria and oxidative enzymes activity.

TABLE 5.4 Summary of Adaptive Respiratory Responses to High Altitude

Parameter	Response to High Altitude
Alveolar PO ₂	↓ (due to decreased barometric pressure)
Arterial PO ₂	↓ (hypoxemia)
Ventilation rate	↑ (hyperventilation due to hypoxemia)
Arterial pH	↑ (respiratory alkalosis due to hyperventilation)
Hemoglobin concentration	↑ (increased red blood cell concentration)
2,3-DPG concentration	↑
O ₂ -hemoglobin dissociation curve	Shifts to right; increased P ₅₀ ; decreased affinity
Pulmonary vascular resistance	↑ (due to hypoxic vasoconstriction)
Pulmonary arterial pressure	↑ (secondary to increased pulmonary resistance)

2,3-DPG, 2,3-diphosphoglycerate.

Acclimatization to low PO₂

