

Prof. Ahmed Fathalla Ibrahim Professor of Anatomy

College of Medicine

King Saud University

E-mail: ahmedfathala@gmail.com

OBJECTIVES


At the end of the lecture, students should:


- Define the autonomic nervous system.
- Describe the structure of autonomic nervous system
- Trace the preganglionic & postganglionic neurons in both sympathetic & parasympathetic nervous system.
- Enumerate in brief the main effects of sympathetic & parasympathetic system


DEFINITION


- Nerve cells located in both central & peripheral nervous system that are concerned with inervation of involuntary structures: viscera, smooth & cardiac muscles, glands.
- Function: maintains homeostasis of internal environment.
- **Regulation:** by hypothalamus.

STRUCTURE OF AUTONOMIC NERVOUS SYSTEM











- □ Preganglionic sympathetic neurons: cells of the lateral horn of spinal cord in all thoracic + upper 3 lumbar segments.
- □ Preganglionic axons leave the spinal cord, join corresponding spinal nerves & reach the sympathetic chain (via the white ramus communicans). They either:
- 1. Synapse with cells of paravertebral ganglia located in sympathetic chain (postganglionic neurons are cells of paravertebral ganglia: postganglionic axons leave the sympathetic chain & join again the spinal nerve (via grey ramus communicans) to supply structures in head & thorax + blood vessels & sweat glands.


2. Leave the sympathetic chain (without synapse) to reach coeliac & mesenteric plexuses (around branches of abdominal aorta) to synapse with their cells. Postganglionic neurons are cells of coeliac & mesenteric plexuses. Postganglionic axons supply abdominal & pelvic viscera.

PARAVERTEBRAL GANGLIA



- ☐ They are interconnected to form 2 sympathetic chains, one on each side of vertebral column.
- **☐** Number of ganglia:
- 1. Three ganglia in cervical part of chain
- 2. Eleven to twelve ganglia in thoracic part
- 3. Four in lumbar & sacral parts.
- ☐ The chains end into a common 'ganglion impar' in front of coccyx





□ Preganglionic parasympathetic neurons:

- 1. Cells located in brain stem: Preganglionic axons leave the brain stem, join 3rd, 7th, 9th & 10th cranial nerves & reach ciliary, pterygopalatine, submandibular, otic & peripheral ganglia (Postganglionic neurons are cells of those ganglia). Postganglionic axons supply structures in head, thorax & abdomen.
- 2. Cells located in 2nd, 3rd & 4th sacral segments of spinal cord. Preganglionic axons leave the spinal cord, join corresponding sacral spinal nerves to reach peripheral ganglia in pelvis where they synapse. Postganglionic neurons are cells of peripheral ganglia. Postganglionic axons supply pelvic viscera.

Structure	Sympathetic 6	ef
	D.11	

Autonomic nervous system

fect Iris of eye

Dilates pupil

Relaxes

Ciliary muscle of eye Salivary glands Lacrimal gland

Heart

Bronchi

tract

Gastrointestinal

Sweat glands

Erector pili

muscles

Reduces secretion Reduces secretion Increases rate and force of contraction

Dilates Decreases motility

Parasympathetic

Constricts pupil

Increases secretion

Increases secretion

Decreases rate and

Increases motility

effect

Contracts

force of

contraction

Constricts

Increases secretion Contracts

QUESTION 1

- □ At which one of the following sites are located preganglionic neurons of the sympathetic nervous system?
- 1. Brain stem
- 2. Thoracic segments of spinal cord —
- 3. Sacral segments of spinal cord
- 4. Sympathetic chain

QUESTION 2

- □ Regarding the parasympathetic nervous system, which one of the following statements is correct?
- 1. Its preganglionic axons are short.
- 2. It supplies sweat glands.
- 3. Its preganglionic neurons are located in the sacral segments of spinal cord. —
- 4. Its postganglionic neurons are located in the coeliac & mesenteric plexuses.

