



Physical and Psychological Factors Affecting Sport Performance

Dr. Aida Korish
Assoc.Prof.Physiology
College of Medicine
KSU


Objectives

By the end of this lecture students should be able to :

1. Identify the muscle metabolic systems and the nutrients used in exercise to regenerate ATP: Phosphocreatine-creatine system, Glycogen-lactic acid system, Aerobic system.
2. Explain the recovery of the muscle metabolic systems after exercise and the phenomena of oxygen debt.
3. Discuss the effects of smoking on pulmonary ventilation in exercise.
4. Correlate between heart diseases and the athletic performance in old age.
5. Describe the changes in body fluids and salts in exercise.
6. Interpret the effects of drugs on athletes.


Metabolic pathways in skeletal muscle

- ❓ Adenosine triphosphate (ATP) is the only energy source used directly by the muscles for contractile activities.
- ❓ The demand and the mechanism of ATP production vary according to the type of work done.
 - At rest, a muscle cell contains a small store of ATP, but it cannot rely on this ATP once it begins contracting.
 - Muscle cell must get ready to ATP production to keep pace with the increased rate of utilization.



Energy for Muscle Contraction

- Mitochondria in the muscle converts glucose, fatty acids, and amino acids into ATP (**Adenosine-PO₃ ~ PO₃ ~ PO₃**).
- Each of the last 2 high energy phosphate bonds in ATP stores 12,000 calories per mole of ATP.
- All ATP stored in the muscle is sufficient for **only 1- 2 seconds** of muscle power. (Enough for half of a 50-meter dash).
- So resting muscles must have energy stored in other forms e.g Creatine Phosphate (CP), glycogen, etc. to convert ADP into ATP



ATP regeneration


- As we begin to exercise, we almost immediately use our stored ATP within few seconds and it will be changed into ADP.

ATP is regenerated from ADP by 3 pathways:


1-Direct phosphorylation of ADP by creatine phosphate (CP).

2-Anaerobic pathway (glycolysis → lactic acid) ?

3-Aerobic oxidation of fatty acids in the mitochondria



INTERACTION OF ENERGY SYSTEMS




1-Phosphocreatine-creatine system (creatine ~ P_i) (CP) (Direct phosphorylation):

CP: Contain high energy phosphate bond of 13,000 calories/mole.

Most muscle cells have 3-8 times as much CP as ATP

Energy transfer from CP to ATP occurs within a small fraction of a second .

Energy of muscle CP is immediately available for contraction just as stored energy of ATP.



Phosphagen energy system:

Formed of combined amounts of cell **ATP + CP**

- Together provide maximal muscle power for 5-10 seconds (enough for 100 meter run).
- Energy of phosphagen system is useful for **maximal short bursts of muscle power (5-10 seconds)**.



II- Anaerobic Glycolysis (Glycogen-Lactic acid system)

Is the primary energy source for peak (sever) muscular activity.



It provides 1.3-1.6 minutes of maximal muscle activity.

Produces **2 ATP** molecules per molecule of glucose coming from blood.

Glycogen \rightarrow Glucose \rightarrow 2 **pyruvic acid** (2 ATP + 2NADH)

2 Pyruvic acid \rightarrow 2 **lactic acid** (2 NAD⁺)

Lactic acid diffuses out of muscles \rightarrow blood \rightarrow taken by the liver \rightarrow Glucose (by gluconeogenesis) \rightarrow blood \rightarrow taken by the muscle again.



Anaerobic metabolism is inefficient... Why?

- Large amounts of glucose are used for very small ATP returns.
- Lactic acid is produced whose presence contributes to muscle fatigue.


Which type of sports uses anaerobic metabolism?

- Sports that requires bursts of speed and activity, e.g., basketball.



Aerobic Metabolism

- Is the primary energy source of resting muscles (to convert glucose into glycogen and to create energy storage compounds as CP).
- During **rest** and **light to moderate** exercise, aerobic metabolism contributes 95% of the necessary ATP.
- It breaks down fatty acids, pyruvic acid (made via glycolysis), and amino acids.
- Produces 36 ATP molecules per glucose molecule.



Comparing the Energy Supply of the Phosphagen System, Anaerobic and the Aerobic systems

ATP generation per minute are the following:

Moles of ATP/min	
Phosphagen system	4
Glycogen-lactic acid system	2.5
Aerobic system	1

When comparing the same systems for endurance, the relative values are the following:

Time	
Phosphagen system	8-10 seconds
Glycogen-lactic acid system	1.3-1.6 minutes
Aerobic system	Unlimited time (as long as nutrients last)

Figure 9.20 Comparison of energy sources used during short-duration exercise and prolonged-duration exercise.




Table 84-1

Energy Systems Used in Various Sports

Phosphagen system, almost entirely

- 100-meter dash
- Jumping
- Weight lifting
- Diving
- Football dashes

Phosphagen and glycogen-lactic acid systems

- 200-meter dash
- Basketball
- Baseball home run
- Ice hockey dashes

Glycogen-lactic acid system, mainly

- 400-meter dash
- 100-meter swim
- Tennis
- Soccer

Glycogen-lactic acid and aerobic systems

- 800-meter dash
- 200-meter swim
- 1500-meter skating
- Boxing
- 2000-meter rowing
- 1500-meter run
- 1-mile run
- 400-meter swim

Aerobic system

- 10,000-meter skating
- Cross-country skiing
- Marathon run (26.2 miles, 42.2 km)
- Jogging



Recovery of aerobic system after exercise

Oxygen Debt

Oxygen Debt is the amount of extra O₂ that must be taken after exercise to restore the muscles to the resting conditions.

When a person stops exercising, the rate of oxygen uptake does not immediately return to pre-exercise levels;

It returns slowly (the person continues to breathe heavily for at least a few minutes and sometimes for as long as 1 hour thereafter).

This extra oxygen is used to repay the oxygen debt acquired during exercise.


Extra Consumption of Oxygen Repays the *Oxygen Debt* After Completion of Strenuous Exercise

Oxygen Debt is about 11.5 L of O₂

This additional oxygen is used to:

- (1) reconvert the lactic acid that has accumulated during exercise back into glucose,
- (2) reconvert adenosine monophosphate and ADP to ATP,
- (3) reconvert creatine and phosphate to phosphocreatine,
- (4) re-establish normal concentrations of oxygen bound with hemoglobin and myoglobin, and
- (5) raise the concentration of oxygen in the lungs to its normal level.

This extra consumption of oxygen after exercise is called **repaying the oxygen debt**.



Recovery of muscle metabolic systems after exercise

- Energy from CP reconstitute ATP.
- Energy from glycogen-lactic acid system reconstitute the phosphagen system (CP+ATP).
- Energy from oxidative metabolism of aerobic system reconstitute all other systems:-glycogen-lactic acid system & CP&ATP

Lactic acid causes fatigue so it should be removed by:

- 1-Portion converted into pyruvic acid that is oxidized by all body tissues.
- 2-The major remaining part is changed into glucose in the liver to replenish glycogen stores of muscles.

Recovery of muscle glycogen

- Reduction of glycogen stores by heavy exercise needs days to be replenished. (unlike ATP, CP, and removal of lactic acid which takes shorter periods).
- On high CHO diet, recovery occurs in 2 days.
- On high fat, high protein or on no food all show very little recovery.

Message:

- 1- Athlete should have high CHO diet before exercise.
 - 2- Not to participate in exhausting exercise during 48 hours preceding the event.
- Read [Guyton & Hall: Textbook of Medical Physiology 13E](#)

Nutrients used during muscle activity

- During early stages of exercise the body uses CHO of muscle and liver glycogen. Also in intense muscle activity the body uses fats and very little amino acids.
- If endurance athletic events last longer than 4-5 hours & during exhaustion muscle glycogen is depleted & muscle depend on fats.
- Glucose solution given to athletes to drink during athletic event supply 30-40% of the energy required during prolonged event as marathon race.



Effects of smoking on pulmonary ventilation in exercise

- Nicotine constricts the terminal bronchioles and increases resistance of airflow into and out of the lungs.
- Smoke irritation causes increased fluid secretion into the bronchial tree and swelling of epithelial layer.
- Nicotine paralyze the cilia of the respiratory epithelial cell surface.
- All lead to fluid and waste accumulation and reduced level of performance.
- chronic smokers may develop emphysema (obstruction of bronchioles+ chronic bronchitis+ destruction of alveoli) so slight exercise cause respiratory distress.


Effects of heart disease and old age on athletic performance

- Cardiac diseases that reduce cardiac output (C.O.P) will reduce muscle power.
- Patient with congestive heart failure has little muscle power to even walk on the floor.
- There is 50% ↓ in C.O.P between age 18-80 years, ↓ in maximal breathing capacity, ↓ in muscle mass and therefore in muscle power with age.
- Read [Guyton & Hall: Textbook of Medical Physiology 13E](#)




Body fluids and salts in exercise

- Exercise for 1 hour during endurance athletic event causes 5-10 pounds of weight loss in hot humid atmosphere due to sweat loss.
- Loss of enough sweat reduces performance 5-10% and may lead to cramps, nausea & serious effects, so it should be replaced.
- Sodium tablets and supplemental fluids containing potassium in the form of fruit juice is required to athletes.
- Acclimatization to exercise by gradual increase over 1-2 weeks instead of maximal exposure is needed.



Drugs and athletes

- Caffeine increase athletes performance.
- Male sex hormone (Androgens) & other anabolic steroids increase athletes performance but they ↑ the risk of heart attacks due to hypertension, ↑LDL and ↓HDL.
- Male sex hormones ↓testicular functions & ↓natural testosterone secretion in males.
- Women develop facial hair, stoppage of menses, ruddy skin and bass voice if they take androgens.
- Amphetamine & cocaine improve performance but overuse reduce performance they are psychic stimuli.
-the action of these drugs in addition to epinephrine and norepinephrine (hormones of adrenal medulla) secreted during exercise leading to death by ventricular fibrillation. Read [Guyton & Hall: Textbook of Medical Physiology 13E](#)



Body fitness prolongs (improves) life

- Studies shows that body fitness, exercise & weight control have additional benefit of prolonged life expectancy (between 50-70).

Reasons:-

- 1-It reduces CVD, heart attacks, brain stroke and kidney disease due to low blood pressure, low blood cholesterol, low LDL, and high HDL.
- 2-It reduces insulin resistance and type 2 diabetes.
- 3-Improved fitness reduces the risk of breast, prostate, and colon cancers and reduces obesity.





Thank you