

Myocarditis and Pericarditis

Prof. Hanan A. Habib & Dr.Khalifa BinKhamis
hahabib@ksu.edu.sa & kbinkhamis@ksu.edu.sa

Department of Pathology ,
Microbiology unit ,
College Of Medicine


2021

Objectives



- Describe the epidemiology, risk factor for myocarditis.
- Explain the pathogenesis of myopericarditis.
- Differentiate between the various types of myocarditis and pericarditis.
- Name various etiological agents causing myocarditis and pericarditis.
- Describe the clinical presentation and differential diagnosis of myocarditis and pericarditis.
- Discuss the microbiological and non microbiological methods for diagnosis of myocarditis and pericarditis.
- Explain the management ,complication and prognosis of patient with myocarditis and/or pericarditis.

Myocarditis

- **Myocarditis** : an inflammatory disease of the heart muscle.
- Mild & self-limited with few symptoms **OR** severe with progression to congestive heart failure & dilated cardiac muscle.
- localized **or** diffuse
- Myocarditis can be due to a variety of **infectious** and **non infectious** causes eg. toxins, drugs and hypersensitivity immune response.
- **Viral infection is the most common cause**



Myocarditis



Epidemiology ,Etiology and Risk Factors

- **Epidemiology** : no accurate estimate of incidence as many cases are mild & brief and diagnosis is not made.
- **Etiology** : **Coxsackie virus B** is the most common cause of myocarditis.

Other virus : **Coxsackie virus A, Echoviruses, Adenoviruses ,Influenza, EBV, Rubella, Varicella, Mumps, Rabies, Hepatitis viruses and HIV.**

Bacterial causes include *Corynebacterium diphtheriae*, Syphilis ,Lyme disease or as a complication of bacterial endocarditis .

Etiology-continue

- Parasitic causes includes Chagas diseases, *Trichinella spiralis*, *Toxoplasma gondii* and *Echinococcus*.
- **Others organisms include: *Rickettsiae*, Fungi, *Chlamydia*, enteric pathogens, *Legionella* and *Mycobacterium tuberculosis*.**
- **Giant cell myocarditis** due to Thymoma, SLE (*systemic lupus erythromatosis*) or Thyrotoxicosis.

Infectious

Noninfectious

Viruses

1. Coxsackie B
2. HIV

Systemic Diseases

1. SLE
2. Sarcoidosis
3. Vasculities(Wegener's disease)
4. Celiac disease

Bacterial

1. *Corynebacterium diphtheriae* (diphtheria)

Neoplastic infiltration

Protozoan

1. *Trypanosoma cruzi* (Chagas disease)

Drugs & Toxins

1. Ethanol
2. Cocaine
3. Radiation
4. Chemotherapeutic agents - Doxorubicin

Spirochete

1. *Borrelia burgdorferi* (Lyme disease)

Clinical presentation of myocarditis

- **Highly variable** :may occur days to weeks after onset of acute febrile illness or with heart failure without any known antecedent symptoms .
- Fever, headache, muscle aches, diarrhea, sore throat and rashes similar to most viral infections
- **Chest pain, arrhythmias ,sweating , fatigue and may present with congestive heart failure.**


Differential Diagnosis

- Acute Myocarditis
- Vasculitis
- Cardiomyopathy (due to drugs **or** radiation)

Diagnosis of myocarditis

- WBCs, ESR, Troponin and CK-MB usually **elevated**
- **ECG** (nonspecific ST-T changes and conduction delays are common)
- **Blood culture**
- **Viral serology and other specific tests** for Lyme disease, diphtheria and Chagas disease may be indicated on a case by case basis.
- **Chest X-rays** : show cardiomegaly
- Radiology : **MRI and Echocardiogram**
- Heart muscle **biopsy** (for some cases)



ECGs of normal heart



Endomyocardial diagnosis

Pathologic examination is not sensitive . It may reveal lymphocytic inflammatory response with necrosis.

“Giant cells” may be seen.




Management of myocarditis

- **Often supportive:** restricted physical activity in heart failure.
- Specific antimicrobial therapy is indicated when an infecting agent is identified.
- Treatment of heart failure arrhythmia
- Other drugs indicated in special situations like anticoagulant, NSAID (non-steroidal anti-inflammatory drugs), steroid or immunosuppressive immunomodulatory agents.
- Heart transplant

Management of myocarditis

- Most cases of viral myocarditis are self limited.
- One third of the patients are left with lifelong **complications**, ranging from mild conduction defects to severe heart failure.
- Patient should be followed regularly every 1-3 months.
- **Sudden death may be the presentation of myocarditis in about 10% of cases.**

Acute Pericarditis



Pericarditis

- **Pericarditis** is an inflammation of the pericardium usually of infectious etiology (viruses, bacterial, fungal or parasitic)
- **Etiology** : (infectious and non-infectious).

Infectious causes :

Viral Pericarditis:

- **Coxsackie virus A and B, Echovirus are the most common causes.**
- **Other viruses includes Herpes viruses, Hepatitis B , Mumps, Influenza, Adenovirus ,Varicella and HIV.**

- **Bacterial Pericarditis** usually a complication of pulmonary infections (e.g. pneumonia , empyema):
organisms : *S. pneumoniae*, ***M. tuberculosis***, *S. aureus*, *H. influenzae*, *K. pneumoniae* , *Legionella pneumophila*, ***Mycoplasma pneumoniae*** & *Chlamydia pneumoniae* .

HIV patients may develop pericardial effusions caused by: *M. tuberculosis* or *M. avium* complex.

- **Disseminated fungal infection** caused by : *Histoplasma*, *Coccidioides*.
- **Parasitic infections** eg. disseminated **toxoplasmosis**, contagious spread of *Entamoeba histolytica* - are rare causes.

Non-infectious pericarditis:

Causes:

- Immune mediated : rheumatic fever & SLE
- Miscellaneous : due to myocardial infarction , malignancy and uremia.

Pathophysiology

- **Contiguous spread**
 - lungs, pleura, mediastinal lymph nodes, myocardium, aorta, esophagus, liver.
- **Hematogenous spread**
 - septicemia, toxins, neoplasm, metabolic
- **Lymphangetic spread**
- **Traumatic or irradiation**

Pathophysiology

- Inflammation provokes fibrinous exudate with or without serous effusion
- The normal transparent and glistening pericardium is turned into a **dull, opaque, and “sandy” sac**
- Can cause pericardial scarring with adhesions and fibrosis.


Types of Pericarditis

- **Caseous Pericarditis** commonly **tuberculous** in origin.
- **Serous Pericarditis** due to **autoimmune** diseases (rheumatoid arthritis, SLE), **viral** infections
 - Transudative serous fluid
- **Fibrinous Pericarditis** due to acute MI, uremia, radiation
 - Fibrinous exudative fluid


Types of Pericarditis

- **Purulent/Suppurative pericarditis** due to bacteria, fungi or parasites.
 - Purulent exudative fluid
- **Hemorrhagic pericarditis** usually caused by infection (e.g. TB) or malignancy
 - blood mixed with a fibrinous or suppurative effusion


Types of pericarditis:



Serous pericarditis



Fibrinous pericarditis



Constrictive Pericarditis

causes:

- Idiopathic
- Radiotherapy
- Cardiac surgery
- Connective tissue disorders
- Dialysis
- **Bacterial infection** (viral, TB, fungal)

Clinical presentation of pericarditis

Acute pericarditis:

- Sudden pleuritic chest pain which is positional retrosternal (relieved by sitting forward)
- Dyspnea
- Fever
- On examination : Pericardial rub, exaggerated pulses , paradoxus JVP (*jugular venous pressure*) and tachycardia.
- As the pericardial pressure increases, palpitations , presyncope or syncope may occur.

Chronic pericarditis:

- Tuberculous pericarditis has **insidious** onset .

Tuberculous Pericarditis

- Incidence of pericarditis in patients with pulmonary TB ranges from 1 – 8 %
- Clinical findings: fever, pericardial friction rub, hepatomegaly
- Tuberculin skin test usually positive
- Fluid smear for acid fast bacilli (**AFB**) often negative
- Pericardial **biopsy** more definitive


Acute Pericarditis

Differential Diagnosis

- Acute myocardial infarction
- Pulmonary embolism
- Pneumonia
- Aortic dissection

Investigations & Diagnosis

- ECG will show ST elevation, PR depression and T-wave inversion may occur later.
- Blood culture
- **Leukocytosis** and an elevated **ESR** are typical
- Other routine testing : **urea** and **creatinine**.
- **Tuberculin skin** test is usually positive in tuberculous pericarditis cases.
- **Chest x-ray** may show enlarged cardiac shadow or calcified pericardium and **CT** scan show pericardial thickening >5mm.
- Pericardial fluid or pericardial **biopsy** specimens for fungi.
- **Immunology /Serology** : Antinuclear antibody tests and Histoplasmosis complement fixation indicated in endemic area.



Concave-up ST elevation



PR segment depression



Management of pericarditis

- Management is largely supportive for cases of idiopathic and viral pericarditis including bed rest , NSAIDS and Colchicine.
- Corticosteroid use is controversial and anticoagulants usually contraindicated.
- Specific antibiotics must include activity against *S. aureus* and respiratory bacteria.

- **Antiviral:**

Acyclovir for *Herpes simplex* or *Varicella* . **Ganciclovir** for CMV .

Pericardiocentesis



Management of pericarditis

- **Pericardiocentesis** : a therapeutic procedure to remove fluid from the pericardium (to relief Tamponade) in severe cases with pericardial effusion.
- Patients who recovered should be observed for recurrence.
- Symptoms due to viral pericarditis usually subsided within one month.



Reference book

Ryan, Kenneth J. Sherris Medical Microbiology. Latest edition.

Mc Graw –Hill education

Tutor contact



For questions and queries
contact tutor :



hahabib@ksu.edu.sa