

RENAL FUNCTIONS & GFR



□ من سلك طريقا يلتمس فيه علما سهل الله له به طريقا إلى الجنة

□

Learning Objectives:


- Enumerate general functions of the kidney.
- Identify and describe that the nephron is the structural and function unit of the kidney.
- Explain glomerular filtration membrane & filtration forces.
- Describe mechanism of filtration & composition of the glomerular filtrate.
- Calculate the net filtration pressure using parameters of Starling forces.

Functions of the Kidney

Synthetic function

Excretion

Regulation




Three distinct regions

Cortex granular superficial region

Medulla exhibits cone-shaped pyramids

Pyramids are bundles of collecting tubules

Renal pelvis flat, funnel-shaped tube




Diagrammatic view of the kidney

What is the functional unit of the kidney?


- The nephron is the functional and structural unit of the kidney.
- Each kidney has 1 million nephrons, each nephron is capable of urine formation.

Structure of a nephron

- The **Glomerulus**: capillary tuft: in which large amount of fluid is filtered from blood.
- **Bwaman's capsule**: Around the glomerulus and receives the filtrate.
- **Tubules**: in which filtered fluid eventually is converted into urine.




The Glomerulus



The nephron



Bowman's capsule: Blind end of the tubule completely surrounds the glomerulus



Efferent arteriole

Capsular space

Proximal tubule cell

The structural and functional units that form urine.

Renal corpuscle

Afferent arteriole

Renal corpuscle

Glomerulus: A tuft of capillaries associated with a proximal tubule

Structure of a nephron, cont.....

- The renal tubule is divided into different sections with different structural and functional characteristics:
- Proximal tubules (in the cortex).
- Loop of Henle.
- Distal tubule (in the renal cortex).
- Connecting tubule, cortical collecting, and the cortical collecting ducts, which run downward in the medulla and become:
- Medullary collecting ducts.

RENAL BLOOD VESSELS

- **AFFERENT ARTERIOLE:**
 - ▣ DELIVERS BLOOD INTO THE GLOMERULI.
- **GLOMERULI:**
 - ▣ CAPILLARY NETWORK THAT PRODUCES FILTRATE THAT ENTERS THE URINARY TUBULES.
- **EFFERENT ARTERIOLE:**
 - ▣ DELIVERS BLOOD FROM GLOMERULI TO PERITUBULAR CAPILLARIES.
- **PERITUBULAR CAPILLARIES:**
 - ▣ VASA RECTA.

Structure of a nephron, cont.....

Types of nephrons:

1- Cortical nephrons: (85%):

Their glomeruli in the outer portion of cortex and have short loops of Henle.

- . Peritubular capelaries


2-Juxtamedullary nephrons: (15%):

Have long loops extended into the medulla.

- . Peritubular capelaries

- . Vasa recta

Maintain salt gradient, helps conserve water




NEPHRON TYPES

Cortical and
Juxtamedullary
Nephrons

1-2 % Blood
Flows
Through
Juxta Medullary
Nephrons

Nephron and Collecting Tubule

Schema




Originates
in outer 2/3
of cortex.

Originates
in inner 1/3
of cortex.

Renal blood flow:

- Renal blood flow to the kidney represents 20% of cardiac output.
- The blood flows to each kidney through a renal artery.
- **Features of renal circulation:**
 - 1- High blood flow rate (1 200 ml/min).
 - 2- Presence of two capillary beds: glomerular and peritubular.Efferent and afferent arterioles are major sites of renal resistance.



Urine formation


The primary function of the kidney is to 'clear' unneeded substances from the blood to be excreted in urine.

Steps of urine formation (basic renal processes):

- 1- **Glomerular filtration:** Filtration of fluid from glomerular capillaries into the renal tubules.
- 2- Tubular reabsorption
- 3- Tubular secretion.
- 4- Excretion.

Urinary excretion rate = Filtration rate - reabsorption + secretion.


1st Step in Urine Formation



**GLOMERULAR
FILTRATION**

4 RENAL PROCESSES


1. Filtration
2. Reabsorption
3. Secretion
4. Excretion



$$\text{Urinary Excretion Rate} = \text{Filtration Rate} - \text{Reabsorption Rate} + \text{Secretion Rate}$$

Urine Formation Preview

- ① **Glomerular filtration**
Creates a plasma-like filtrate of the blood
- ② **Tubular reabsorption**
Removes useful solutes from the filtrate, returns them to the blood
- ③ **Tubular secretion**
Removes additional wastes from the blood, adds them to the filtrate
- ④ **Water conservation**
Removes water from the urine and returns it to blood, concentrates wastes



Glomerular filtration rate (GFR)

- The first step in urine formation is glomerular filtration.
- It is the filtration of fluid from the glomerular capillaries into the renal tubules.
- It contains all substances present in plasma except proteins.
- GFR is normally $125 \text{ ml/min} = 20\%$ renal plasma flow.

- What is glomerular membrane?
- What will filter? (composition of filtrate)
- What determine GFR?
- What are the forces responsible for passage of fluid (filtrate) through this membrane?
- Regulation of GFR


Glomerular membrane

Blood in the glomerulus is separated from the fluid in the Bowman's space by a filtration barrier (glomerular membrane) consisting of three layers:

- 1 - Single layer of capillary endothelium.
- 2- Single epithelial lining of Bowman's capsule (Podocytes) During filtration the fluid moves between their foot processes (psudopodia).
- 3- Basement membrane between endothelium and epithelium.


Glomerular membrane

Here we see a glomerular capillary in longitudinal section




Capillary
endothelium

Glomerular membrane




**Capillary
endothelium**

Glomerular membrane



**Basement
membrane**

Glomerular membrane




Podocytes (cell body with nucleus)


Filtration slits

Filtration Membrane



Filter that lies between the blood and the interior of the glomerular capsule




FILTRATION MEMBRANE



Glomerular membrane



Glomerular membrane




Characteristics of glomerular membrane:


- Allow passage of molecules up to 70,000 D
- Albumin does not normally pass as they are repelled by the negative charge of the **glycoproteins** material of basement membrane.
- Blood cells don not normally pass through the membrane.

GLOMERULAR FILTRATE

Glomerular filtrate



Common components of glomerular filtrate:



The concentration of each of these substances in the glomerular filtrate is similar to its concentration in **plasma**.

Glomerular Filtration Rate (GFR)

□ The GFR is determined by:

1 - the net filtration pressure across the glomerular capillaries.


2- the glomerular capillary filtration coefficient (Kf)

$$\text{GFR} = K_f \times \text{Net filtration pressure.}$$

$$= 12.5 \times 10 = 125 \text{ ml/min}$$

Glomerular Filtration Rate (GFR)

What controls (determines) the GFR?



Factors related to the membrane itself

Permeability

Filtering surface area

Capillary filtration coefficient (**K_f**)


The sum of the forces acting across the membrane (starling forces)

Net filtration pressure (**NFP**)

GFR = 125 ml/min
OR 180 L/day

$$\text{GFR} = K_f \times \text{NFP}$$

Net Filtration Pressure (NFP)



Blood hydrostatic pressure (BHP)
Colloid osmotic pressure (COP)
Capsular pressure (CP)

Net filtration pressure (NFP)

60 mmHg out
-32 mmHg in
-18 mmHg in

10 mmHg ou

REGULATION OF GFR



Regulation of GFR

$$\text{GFR} = K_f \times (P_G - \pi_G - P_B + \pi_B)$$

- Any factor that affect the parameters in the equation will affect the GFR.
- However, physiologic regulation of the GFR involves mechanisms that affect mainly the P_G .

P_G depends on:

- Arterial BP
- Afferent arteriolar resistance.
- Efferent arteriolar resistance.

K_f

π_G

P_B

Can get affected in disease conditions causing changes in GFR

How changes in Forces determining GFR affect GFR?

- Increased **Bowman's capsule pressure** decreases GFR. It can happen in urinary obstruction e.g. stones , tumors..
- Increased **glomerular capillary colloid** osmotic pressure decreases GFR.
- Increased **glomerular capillary hydrostatic** pressure increases GFR. This pressure is affected by:
 - ABP.
 - Afferent arteriolar resistance.
 - Efferent arteriolar resistance

Physiologic Regulation of GFR

Constriction of *Afferent* arteriole

↓↓ P_G

↓↓ GFR

Sympathetic stimulation
Epinephrine
Norepinephrine
Endothelin

Constriction of *Efferent* arteriole

Moderate

↑↑ P_G


↑↑ GFR

Severe

↓↓ P_G


↓↓ GFR

Angiotensin II



Decreased GFR

Increased GFR




As vasodilation and vasoconstriction of the afferent and efferent arterioles alter the blood flow through the glomerular capillaries, there are corresponding alterations in the glomerular filtration rate (GFR).

Factors affecting Renal blood flow and GFR

- Sympathetic stimulation of renal arterioles decrease GFR & RBF.
- Norepinephrine decreases GFR & RBF.
- Angiotensin II decreases RBF. It constricts efferent arteriole more than afferent .
- High protein diet increases GFR.
- Hyperglycemia increases GFR & RBF.
- Fever increases GFR & RBF.
- Aging decreases RBF & GFR

Juxtaglomerular Apparatus



Mesangial cells
-contractile properties
-capillary filtration

Granular Cells
-smooth muscles
-secrete Renin
-mechanoreceptors

Efferent arteriole


JGA

Macula densa cells

Mesangial cells

Granular cells

Afferent arteriole



Proximal tubule cell

Mesangial cells

Macula Densa
-tubular cells
-chemoreceptors (Na⁺, K⁺ and Cl⁻)