




OVERVIEW OF MOTOR ACTIVITY CONTROL








MAIN INPUT TO THE BASAL GANGLIA

The comes from the cerebral cortex (motor area) and projects to the **NEOSTRIATUM**
(a term for the caudate nucleus and putamen)

THE MAIN OUTPUT

Is via the thalamus to the cerebral cortex
(motor area)



BASIC CIRCUITS OF BASAL GANGLIA

1. Motor loop (putamen circuit) concerned with learned movement.
2. Cognitive loop (Caudate circuit) concerned with cognitive control of sequences of motor pattern. Basically it is concerned with motor intentions. (Note: cognition means thinking process using sensory input with information already stored in memory.)
3. Limbic loop: involved in giving motor expression to emotions like, smiling, aggressive or submissive posture.
4. Oculomotor loop concerned with voluntary eye movement [saccadic movement]

10


The Putamen Circuit



- Prim Motor Cortex
- Premotor
- Suppl Motor Areas



Ventroanterior and
ventrolateral
nuclei of thalamus



- Premotor
- Suppl Motor
- Somatosensory Cortex




The Caudate Circuit



- Prefrontal
- Premotor
- Suppl Motor




Ventroanterior and
ventrolateral
nuclei of thalamus




Association
Areas



Basal Ganglial Pathways Loops





Metabolic characteristics

- High Oxygen consumption .
- High Copper content in Wilson's disease (Copper intoxication):
- Autosomal Recessive
- Copper binding protein Ceruloplasmin is low
- Lenticular degeneration occurs

BASAL GANGLIA

FUNCTIONS

- Control of movements
- Planning and programming of movements
- Cognition

The Putamen Circuit

Executes Learned Patterns of Motor Activity

- Basal ganglia function in association with the corticospinal system to control *complex patterns of motor activity*.
- Examples are:
 - writing of letters of the alphabet.
 - cutting paper with scissors,
 - hammering nails,
 - shooting a basketball through a hoop,
 - passing a football,
 - throwing a baseball,
 - the movements of shoveling dirt,
 - most aspects of vocalization,
 - controlled movements of the eyes
 - virtually any other of our skilled movements, most of them performed subconsciously.



The Caudate Circuit

Cognitive Control of Sequences of Motor Patterns

- Cognition means the thinking processes of the brain, using both sensory input to the brain plus information already stored in memory. Thoughts are generated in the mind by a process called cognitive control of motor activity.
- Example: A person seeing a lion approach and then responding instantaneously and automatically by (1) turning away from the lion, (2) beginning to run, and (3) even attempting to climb a tree.
- Thus, cognitive control of motor activity determines subconsciously, and within seconds, which patterns of movement will be used together to achieve a complex goal



The Caudate Circuit

Change the Timing and to Scale the Intensity of Movements

- Two important capabilities of the brain in controlling movement are
 - (1) to determine how rapidly the movement is to be performed and
 - (2) to control how large the movement will be.
- For instance, a person may write the letter "a" slowly or rapidly. Also, he or she may write a small "a" on a piece of paper or a large "a" on a chalkboard. Regardless of the choice, the proportional characteristics of the letter remain nearly the same

BASAL GANGLIA

DISORDERS

MOVEMENTS (ATAXIA Rate, Range, Force, Direction)

SPEECH

POSTURE

GAIT

MENTAL ACTIVITY

OTHERS

Movement Disorders

Hyperkinetic

- Hemiballismus
- Huntington's Disease
- Athetosis



Hypokinetic

- Parkinson's Disease
- Drug Induced (Neuroleptics, MPTP)

Movement Disorder	Features	Lesion
Chorea	Multiple quick, random movements, usually most prominent in the appendicular muscles	Atrophy of the striatum. Huntington Chorea
Athetosis	Slow writhing movements, which are usually more severe in the appendicular muscles	Diffuse hypermyelination of corpus striatum and thalamus
Hemiballismus	Wild flinging movements of half of the body	Hemorrhagic destruction of contralateral subthalamic n. Hypertensive patients
Parkinsonism	Pill rolling tremor of the fingers at rest, lead pipe rigidity and akinesia	Degeneration of Substantia Nigra

Parkinson's Disease

- Described by James Parkinson
- Degeneration of dopaminergic nigrostriatal neurons (60-80 %).
- Phenthiazines (tranquilizers drugs) .
- Methyl-Phenyl-Tetrahydro-Pyridine (MPTP). The oxidant MPP⁺ is toxic to SN.
- Five cardinal features
 - Tremor
 - Rigidity
 - Akinesia & Bradykinesia
 - Postural Changes
 - Speech Changes



Hemiballismus



- Injury usually to STN
- Decreased inhibition (Indirect Pathway)
- Characterized by uncontrolled flinging
- TX: Dopamine Antagonist



Huntington's Disease

- Hereditary , autosomal dominant
- Rare onset at 30-40s as early as 20s
- Disease of caudate & putamen.
- Early, Jerky movement of hands toward end of reaching an object
- Later, hyperkinetic choreiform movements appear and gradually increase until they incapacitate the patient
- Slurred speech and incomprehensive
- Progressive Dementia
- Loss of GABA nergic neurons