



Done By:


Arwa Al-Madani

mashaal Al-towairqy

Sarah Bin-Hussain

Bedoor Al-Qadrah

Reham Al-Henaki



**What is the hemoglobin (HB)?**

hemeprotein found only in red blood cells

**what is it function?**

It binds reversibly to O<sub>2</sub> , so it is O<sub>2</sub> transporter from lung to tissues

**It is formed of 2 components :**

heme as prosthetic group \* and globine that is protein

\***prosthetic group** → non-protein molecule attached to the protein to make it functioning.


**The heme group is formed of :**

**1-protoporphyrine IX (9)** that form a ring

**2-ferreus iron Fe<sup>++</sup>** that is located at the center of the heme

Ferreus iron has 6 bonds , **they are :**

4 bonds with 4 nitrogens (N) of the protophyrine ring , 1 with histidine residue of globin chain and the last to the O<sub>2</sub>



HB type	Formed of	Normal level	Found in	Relation with O <sub>2</sub>	function
<b>HB A</b>	$\alpha 2 \beta 2$	97% Male 14-16g/dl Female 13-15 g/dl	Adult	Affected by Three allosteric effectors	-Carries O <sub>2</sub> from the lungs to tissues -Carries CO <sub>2</sub> from tissues back to the lungs
<b>HB F</b>	$\alpha 2 \gamma 2$	1 %	Fetus and newborn	Higher O <sub>2</sub> affinity than HB A	Transfers O <sub>2</sub> from maternal to fetal circulation across placenta
<b>HB A<sub>2</sub></b>	$\alpha 2 \delta 2$	2 %	Appears 12 weeks after birth		
<b>HbA<sub>1c</sub></b>	Hb A undergoes non-enzymatic glycosylation (depending on plasma glucose)				HbA <sub>1c</sub> levels are high in patients with diabetes mellitus
<b>Carboxy-Hb</b>	CO carried by HB instead of O <sub>2</sub>				CO replaces O <sub>2</sub> and binds 200X tighter than O <sub>2</sub> (in smokers)
<b>Met-Hb</b>	Contains oxidized Fe <sup>3+</sup>	~2%		cannot carry O <sub>2</sub>	
<b>Sulf-Hb</b>	due to high sulfur levels in blood				irreversible reaction

**Hbs : Unable to transport O<sub>2</sub> due to abnormal structure**

## HB A

Composed of four polypeptide chains:


Two  $\alpha$  and two  $\beta$  chains

Contains two dimers of  $\alpha\beta$  subunits


$\alpha$  &  $\beta$  chains Held together by non-covalent hydrophobic interactions


the bonds between the 2 dimers  $\alpha\beta=\alpha\beta$  is weak ionic and hydrogen

Each chain is a subunit with a heme group in the center that carries oxygen, so the HB is able to carry 4  $O_2$  molecules (because it's formed of 4 chains)



HB structure	Form of HB is	movement	O <sub>2</sub> affinity
<b>T (taut) form</b>	Deoxy HB (not attached with HB)	The movement of dimers is constrained	Low affinity
<b>R (relaxed) form</b>	Oxy HB (attached with HB)	dimers have more freedom of movement	High affinity





### O<sub>2</sub> Dissociation Curve:

It is sigmoidal shaped curve → indicates the cooperation of subunits in O<sub>2</sub> binding


(so when ↑pO<sub>2</sub> → ↑saturation of HB to O<sub>2</sub> → Increasing the HB affinity to O<sub>2</sub>)

That means Binding of O<sub>2</sub> to one heme group increases O<sub>2</sub> affinity of others and that's called (heme-heme interaction) that the affinity of hemoglobin for the last oxygen bound is approximately 300 times greater than its affinity for the first oxygen bound.

So that the affinity of hemoglobin for the last oxygen bound is approximately 300 times greater than its affinity for the first oxygen bound.

So oxygenation favors → R form

Deoxygenation favors → T form



The allosteric factor	What is it	When / where increase	When / where decrease
<b>1- pO<sub>2</sub></b> <b>indicates affinity of Hb to O<sub>2</sub></b>	<p>the pressure at which Hb is 50% saturated with O<sub>2</sub></p> <p>that means the pressure needs to make the half of Hb saturated with O<sub>2</sub>.</p>	<p>In lungs will cause: High affinity → slow unloading of O<sub>2</sub>.</p> <p>Saturation in lung 100% because the PO<sub>2</sub> is 100mm.</p> <p>So O<sub>2</sub> will be taken from the lungs.</p>	<p>In tissues Will cause Low affinity → fast unloading of O<sub>2</sub> so saturation reduced because the PO<sub>2</sub> in tissues is 40mm</p> <p>So O<sub>2</sub> will be delivered to tissues.</p>
<b>2- PH+CO<sub>2</sub> = Bohr Effect</b>	<p>Bohr effect :</p> <ul style="list-style-type: none"> <li>- removes insoluble CO<sub>2</sub> from blood stream</li> <li>- Produces soluble bicarbonate</li> </ul>	<p>In the lungs :</p> <p>The free Hb (after releasing of O<sub>2</sub> into the tissues) binds to two protons</p> <p>Protons are released and react with HCO<sub>3</sub><sup>-</sup> – to form CO<sub>2</sub> gas</p> <p>The proton-poor Hb now has greater affinity for O<sub>2</sub> .</p>	<p>In tissues :</p> <p>They have ↓ PH (acidic) than the lungs Due to proton generation: CO<sub>2</sub> + H<sub>2</sub>O → HCO<sub>3</sub><sup>-</sup> + H<sup>+</sup></p> <p>Proton ↓O<sub>2</sub> affinity of Hb → easier O<sub>2</sub> release into the tissues.</p>
<b>3- 2,3bisphosphoglycerate (BPG)</b>	<ul style="list-style-type: none"> <li>-Binds to deoxy-Hb and stabilizes the T-form</li> <li>-When oxygen binds to Hb, BPG is released</li> </ul>	<p>In high altitude as a compensatory mechanism to decrease O<sub>2</sub> affinity of Hb and deliver it faster to tissues</p> <p>-High altitude → ↓O<sub>2</sub> concentration → hypoxia → ↑2,3BPG → ↓O<sub>2</sub> affinity → ↑O<sub>2</sub> delivery to tissues</p>	



# So as a conclusion:

