

Gastrointestinal Physiology

Lecture 6

Physiology of the Pancreas

Chapter 64; Pages: 799-802

Dr. Hayam Gad

*Associate Professor of
Physiology*

Learning Objectives





- * Pancreatic Acini
- * Pancreatic Secretion
- * Pancreatic Enzymes
- * Control of Pancreatic Secretion
 - o Neural Control
 - o Hormonal Control
 - Secretin
 - Choecystokinine

The exocrine pancreas


The pancreas, which lies parallel to and beneath the stomach is composed of:-

- 1) The endocrine islets of Langerhans which secrete insulin, glucagone and somatostatin.
- 2) Acinar gland tissues which produce pancreatic juice (the main source of digestive enzymes).

The cells lining the acini are serous cells containing zymogen granules.



The main pancreatic duct joins into bile duct at ampulla of Vater that is surrounded by sphincter of Oddi



Pancreatic Secretion

- ☞ Pancreatic juice is secreted in response to the presence of chyme in the upper portions of the small intestine.
- ☞ ***The major functions of pancreatic secretion:***
 - ☞ To neutralize the acids in the chyme
 - ☞ To produce enzymes involved in the digestion of dietary carbohydrate, fat, and protein.

Volume: 1.2-1.5 l/day.

The osmolarity of pancreatic fluid is equal to that of plasma (isotonic)


PH= 8 alkaline.

Composition: 1 % inorganic materials (electrolytes)
1-2 % organic materials mostly
enzymes.



The electrolytes

- They are produced from the epithelial cells of the ductules and ducts and include cations Na^+ , K^+ , Ca^{++} and anions HCO_3^- and Cl^- .
- The greater bulk of electrolytes is in the form of NaHCO_3 .
- HCO_3^- concentration increases with increasing secretion rate.

Flow Rate and pancreatic secretion



Secretion of Bicarbonate Ions into Pancreatic Juice





Pancreatic enzymes

The pancreas secretes enzymes that act on all major types of food stuffs.

1- Pancreatic proteolytic enzymes (proteases)

- ◆ Trypsin, chymotrypsin, elastase, carboxypeptidase.
- ◆ They are secreted in inactive form and activated in intestinal lumen.
- ◆ Trypsinogen is activated into trypsin by the enzyme enteropeptidase (enterokinase), secreted by duodenal mucosal cells.
- ◆ Trypsin activates chymotrypsinogen to chymotrypsin, proelastase to elastase and procarboxypeptidase into carboxypeptidase.






◎ Trypsin, chymotrypsin and elastase are endopeptidases, splitting protein into shorter peptide chains.

◎ Carboxypeptidase is an exopeptidase which splits off amino acids at the carboxyl terminus of the peptide.

◎ Trypsin inhibitor is present in cytoplasm of glandular cells. It inhibits activation of trypsin in acini and ducts of the pancreas.



2- Pancreatic amylase

It splits starch to maltose, maltotriose and dextrans.

3- Enzymes for fat digestion


- a. Pancreatic lipase is the most important fat splitting enzyme. It breaks TG into MG and FA in the presence of bile salts and colipase.
- b. Cholesterol esterase which liberates cholesterol.
- c. Phospholipase A_2 which splits phospholipids into lysophospholipids & FA.

End Products of Fat Digestion

DIGESTION OF LIPIDS


Triglyceride

lingual and pancreatic lipases




Cholesterol ester

cholesterol ester hydrolase



Phospholipid

phospholipase A₂



Pancreatic enzymes for digestion

Protein digestion

Trypsin
Chymotrypsin
Carboxypolypeptidase

Carbohydrate digestion

Pancreatic amylase,
which hydrolyzes
starches, glycogen,
and most other
carbohydrates

Fat digestion

Pancreatic lipase
Cholesterol esterase
Phospholipase

Characteristics of Pancreatic Enzymes

Enzyme	Specific Hydrolytic Activity
Proteolytic	
Endopeptidases	
Trypsin(ogen)	Cleaves peptide linkages in which the carboxyl group is either arginine or lysine
Chymotrypsin(ogen)	Cleaves peptides at the carboxyl end of hydrophobic amino acids, e.g., tyrosine or phenylalanine
(Pro)elastase	Cleaves peptide bonds at the carboxyl terminal of aliphatic amino acids
Exopeptidase	
(Pro)carboxypeptidase	Cleaves amino acids from the carboxyl end of the peptide
Amylolytic	
α -Amylase	Cleaves α -1,4-glycosidic linkages of glucose polymers
Lipases	
Lipase	Cleaves the ester bond at the 1 and 3 positions of triglycerides, producing free fatty acids and 2-monoglyceride
(Pro)phospholipase A ₂	Cleaves the ester bond at the 2 position of phospholipids
Carboxylesterhydrolase (cholesterol esterase)	Cleaves cholesteryl ester to free cholesterol
Nucleolytic	
Ribonuclease	Cleaves ribonucleic acids into mononucleotides
Deoxyribonuclease	Cleaves deoxyribonucleic acids into mononucleotides

The suffix -ogen or prefix pro- indicates the enzyme is secreted in an inactive form


Pancreatic Secretion is Under Neural and Hormonal Control

- ★ Parasympathetic stimulation (through Ach on acinar cells) results in an increase in enzyme secretion-fluid and HCO_3^-
- ★ Secretin tends to stimulate a HCO_3^- rich secretion by activating ductal cells.
- ★ Cholecystokinin (CCK) stimulates a marked increase in enzyme secretion by stimulating the acinar cells.
- ★ Pancreatic secretion normally results from the combined effects of the multiple basic stimuli, not from one alone (potentiate each other).

Phases of Pancreatic secretion

Phase	stimulus	Mediators
Cephalic phase	Smell, taste, chewing and swallowing	Release of Ach and gastrin
Gastric phase	Protein, gastric distention	Vago-vagal reflex
Intestinal phase	Acid in chyme, fatty acids	Secretin, CCK and vago-vagal reflex

REGULATION OF PANCREATIC SECRETION




Acid from stomach releases secretin from wall of duodenum; fats and amino acids cause release of cholecystokinin

Common bile duct

Vagal stimulation releases enzymes into acini


Secretin causes copious secretion of pancreatic fluid and bicarbonate; cholecystokinin causes secretion of enzymes

Secretin and cholecystokinin absorbed into blood stream



Secretin hormone

- ✓ It is a peptide released into the blood from “S” cells in upper intestinal mucosa.
- ✓ Stimuli for its release:
 - Mainly acids (pH 4 or less)
 - To a less extent AA and FA.



Functions


- ✿ It acts on pancreatic duct cells to stimulate secretion of HCO_3^- and H_2O .
- ✿ It acts on biliary duct cells to stimulate hepatic bile flow and HCO_3^- secretion.
- ✿ It augments the action of CCK in stimulating pancreatic enzyme secretion.
- ✿ It inhibits gastric acid secretion and gastrin release, but it stimulates pepsin secretion.
- ✿ It inhibits gastric motility, contracts pylorus and slows gastric emptying.
- ✿ It relaxes LES.
- ✿ It inhibits intestinal motility and contracts ileocecal sphincter.

Cholecystokinin (CCK)

- ✓ It is a peptide released from “I” cells in the upper intestine.
- ✓ Stimuli of release:-
 - Mainly by AA and FA
 - To a lesser extent by HCl.

✓ Functions

- ✿ It acts on pancreatic acinar cells to stimulate enzyme secretion. It also augments stimulation of H_2O and HCO_3^- secretion by secretin.
- ✿ It has trophic effect on pancreas.
- ✿ It contracts gall bladder, relaxes sphincter of Oddi and causes bile discharge into intestine.
- ✿ It stimulates gastric motility, contracts pylorus thus slows gastric emptying.
- ✿ It relaxes LES.
- ✿ It stimulates intestinal motility.
- ✿ It may be concerned with the mechanism of satiety.



Hormonal control of pancreatic exocrine secretion

